

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 March 2006 (16.03.2006)

PCT

(10) International Publication Number
WO 2006/028451 A1

(51) International Patent Classification⁷: **A61K 31/4745**,
31/5377, C07D 471/04, A61P 31/12, 35/00, 37/02

(21) International Application Number:
PCT/US2004/028754

(22) International Filing Date:
3 September 2004 (03.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant: **3M INNOVATIVE PROPERTIES COMPANY** [US/US]; 3M Center, P.O. Box 33427, Saint Paul, MN 55133-3427 (US).

(72) Inventors: **GRIESGRABER, George, W.**; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). **MANSKE, Karl, J.**; P.O. Box 33427, Saint Paul, MN 55133-3427 (US).

(74) Agents: **ERSFELD, Dean, A.** et al.; Office of Intellectual Property Counsel, P.O. Box 33427, Saint Paul, MN 55133-3427 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 1-AMINO 1-H-IMIDAZOQUINOLINES

(57) Abstract: 1-Amino 1<i>H</i>-imidazoquinoline compounds, pharmaceutical compositions containing the compounds, intermediates, and methods of making and methods of use of these compounds as immunomodulators, for modulating cytokine biosynthesis in animals and in the treatment of diseases including viral and neoplastic diseases are disclosed.

WO 2006/028451 A1

1-AMINO 1*H*-IMIDAZOQUINOLINES


BACKGROUND OF THE INVENTION

There has been a major effort in recent years to find compounds that modulate the immune system. Examples of such compounds, which have demonstrated cytokine inducing and immunomodulating activity, are disclosed by U.S. Patent Nos. 4,689,338; 4,929,624; 5,266,575; 5,268,376; 5,352,784; 5,389,640; 5,446,153; 5,482,936; 5,494,916; 5,756,747; 6,110,929; 6,194,425; 6,331,539; 6,376,669; 6,451,810; 6,525,064; 6,541,485; 6,545,016; 6,545,017; 6,656,938; 6,660,735; 6,660,747; 6,664,260; 6,664,264; 6,664,265; 6,667,312; 6,670,372; 6,677,347; 6,677,348; and 6,683,088.


But despite important progress in the effort to find immunomodulating compounds, there is still a critical scientific and medical need for additional compounds that have an ability to modulate aspects of the immune response, by induction of cytokine biosynthesis or other mechanisms.

SUMMARY OF THE INVENTION

It has now been found that certain 1-amino 1*H*-imidazoquinoline compounds modulate cytokine biosynthesis. In one aspect, the present invention provides compounds of the Formulas I-1 and II-1:



I-1



II-1

wherein R_1' , R_1 , R_2 , R_3 , R , R_A , n , and m are as defined below; and pharmaceutically acceptable salts thereof.

The compounds of Formulas I-1 and II-1 are useful as immune response modifiers (IRMs) due to their ability to modulate cytokine biosynthesis (e.g., induce or inhibit the biosynthesis or production of one or more cytokines) and otherwise modulate the immune response when administered to animals. Compounds can be tested per the test procedures described in the Examples Section. Compounds can be tested for induction of cytokine biosynthesis by incubating human peripheral blood mononuclear cells (PBMC) in a culture with the compound(s) at a concentration range of 30 to 0.014 μ M and analyzing for interferon (α) or tumor necrosis factor (α) in the culture supernatant. Compounds can be tested for inhibition of cytokine biosynthesis by incubating mouse macrophage cell line Raw 264.7 in a culture with the compound(s) at a single concentration of, for example, 5 μ M and analyzing for tumor necrosis factor (α) in the culture supernatant. The ability to modulate cytokine biosynthesis, for example, induce the biosynthesis of one or more cytokines, makes the compounds useful in the treatment of a variety of conditions such as viral diseases and neoplastic diseases, that are responsive to such changes in the immune response.

In another aspect, the present invention provides pharmaceutical compositions containing the immune response modifier compounds, and methods of inducing cytokine biosynthesis in animal cells, treating a viral disease in an animal, and/or treating a neoplastic disease in an animal by administering to the animal one or more compounds of the Formulas I-1 and/or II-1, and/or pharmaceutically acceptable salts thereof.


As used herein, "a," "an," "the," "at least one," and "one or more" are used interchangeably.

The terms "comprising" and variations thereof do not have a limiting meaning where these terms appear in the description and claims.


The above summary of the present invention is not intended to describe each disclosed embodiment or every implementation of the present invention. The description that follows more particularly exemplifies illustrative embodiments. Guidance is also provided herein through lists of examples, which can be used in various combinations. In each instance, the recited list serves only as a representative group and should not be interpreted as an exclusive list.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS OF THE INVENTION

In one aspect, the present invention provides compounds of the Formulas I-1 and II-1:




I-1



II-1

wherein R_1' , R_1 , R_2 , R_3 , R , R_A , n , and m are as defined below; and pharmaceutically acceptable salts thereof.

In one embodiment, the present invention provides compounds of the following Formula I-1:



I-1

wherein:

R_1' is selected from the group consisting of hydrogen, alkyl, alkoxyalkylenyl, and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R_1' is bonded;

5
 -N(R₆)-C(R₇)-aryl,
 -N(R₆)-S(O)₂-C₁₋₁₀ alkyl,
 -N(R₆)-S(O)₂-aryl,
 -C(O)-C₁₋₁₀ alkyl,
 -O-C(R₇)-C₁₋₁₀ alkyl,
 -O-C(R₇)-aryl,
 -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl,
 -O-C(R₇)-N(R₆)-aryl;
 10 -C(O)-O-C₁₋₁₀ alkyl,
 -N₃,
 aryl,
 heteroaryl,
 heterocyclyl,
 -C(O)-aryl, and
 15 -C(O)-heteroaryl;

R₃ is selected from the group consisting of:

-Z'-R₄',
 -Z'-X'-R₄',
 -Z'-X'-Y'-R₄', and
 20 -Z'-X'-R₅';

each R is independently selected from the group consisting of alkyl, alkenyl, alkoxy, halogen, fluoroalkyl, hydroxy, amino, alkylamino, and dialkylamino;


n is an integer from 0 to 4;

m is 0 or 1; with the proviso that when m is 1, then n is 0, 1, or 2;

25 R₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, carboxy, formyl, aryl, aryloxy, arylalkoxy, heteroaryl, heteroaryloxy, heteroarylalkoxy, heterocyclyl, heterocyclylalkylenyl, amino,
 30 alkylamino, (arylalkylenyl)amino, dialkylamino, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo, with the proviso that when R₄ is a substituted alkyl group and the

substituent contains a hetero atom which bonds directly to the alkyl group then the alkyl group contains at least two carbons between the substituent and the nitrogen atom to which R₁ is bonded;


R₅ is selected from the group consisting of:



X is C₂₋₂₀ alkylene;

Y is selected from the group consisting of -C(R₇)-, -C(R₇)-O-, -S(O)₂-, -S(O)₂-N(R₆)-, and -C(R₇)-N(R₉)-; wherein R₉ is selected from the group consisting of hydrogen, alkyl, and arylalkylenyl; or R₉ and R₄ together with the nitrogen atom to which R₉ is bonded can join to form the group

10



Z is selected from the group consisting of -O- and -S(O)₀₋₂-;

A is selected from the group consisting of -CH(R₆)-, -O-, -N(R₆)-, -N(Y-R₄)-, and -N(X-N(R₆)-Y-R₄)-;


15 a and b are independently integers from 1 to 4 with the proviso that when A is -O-, -N(R₆)-, -N(Y-R₄)-, or -N(X-N(R₆)-Y-R₄)- then a and b are independently integers from 2 to 4;

R₄' is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, hydroxyalkyl, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, aryl, aryloxy, arylalkyleneoxy, heteroaryl, heteroaryloxy, heteroarylalkyleneoxy, heterocyclyl, amino, alkylamino, dialkylamino, (dialkylamino)alkyleneoxy, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo;

20









25

R_5' is selected from the group consisting of:



X' is selected from the group consisting of alkylene, alkenylene, alkynylene, arylene, heteroarylene, and heterocyclylene wherein the alkylene, alkenylene, and alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene, or heterocyclylene and optionally interrupted by one or more -O- groups;

Y' is selected from the group consisting of:



Z' is a bond or -O-;

A' is selected from the group consisting of -CH₂-, -O-, -C(O)-, -S(O)₀₋₂-, and -N(R₄')-;

5 Q is selected from the group consisting of a bond, -C(R₇)-, -C(R₇)-C(R₇)-, -S(O)₂-, -C(R₇)-N(R₁₁)-W-, -S(O)₂-N(R₁₁)-, -C(R₇)-O-, and -C(R₇)-N(OR₁₂)-;

V is selected from the group consisting of -C(R₇)-, -O-C(R₇)-, -N(R₁₁)-C(R₇)-, and -S(O)₂-;

W is selected from the group consisting of a bond, -C(O)-, and -S(O)₂-;

10 c and d are independently integers from 1 to 6 with the proviso that c + d is ≤ 7, and when A' is -O- or -N(R₄')- then c and d are independently integers from 2 to 4;

each R₆ is independently selected from the group consisting of hydrogen, alkyl, and arylalkylenyl;

each R₇ is independently selected from the group consisting of =O and =S;


each R₈ is independently C₂₋₇ alkylene;

15 R₁₀ is C₃₋₈ alkylene;

each R₁₁ is independently selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₁₋₁₀ alkoxyC₂₋₁₀ alkylenyl, and arylC₁₋₁₀ alkylenyl; and

R₁₂ is selected from the group consisting of hydrogen and alkyl; or a pharmaceutically acceptable salt thereof.

20 In one embodiment, the present invention provides compounds of the following Formula II-1:



II-1

wherein:

25 each R_A is independently selected from the group consisting of:

halogen,

hydroxy,

alkyl,

alkenyl,

haloalkyl,
alkoxy,
alkylthio,
-NH₂,
-NH(alkyl), and
-N(alkyl)₂;


n is an integer from 0 to 4;

R₁' is selected from the group consisting of hydrogen, alkyl, alkoxyalkylenyl, and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R₁' is bonded;

R_1 is selected from the group consisting of:

$-R_4$,
 $-Y-R_4$,
 $-X-R_5$,
 $-X-N(R_6)-Y-R_4$,
 $-X-C(R_7)-N(R_6)-R_4$,
 $-X-O-C(R_7)-N(R_6)-R_4$,
 $-X-S(O)_2-N(R_6)-R_4$, and
 $-X-O-R_4$;

or R₁' and R₁ together with the nitrogen atom to which they are bonded can join to form a group selected from the group consisting of:



R_2 is selected from the group consisting of:

hydrogen,
alkyl,
alkenyl,
aryl,
heteroaryl,
heterocyclyl,
-alkylene-Z-alkyl,


-alkylene-Z-aryl,
 -alkylene-Z-alkenyl, and
 alkyl or alkenyl substituted by one or more substituents selected from the
 group consisting of:

- 5 -OH,
 halogen,
 -N(R₆)₂,
 -C(R₇)-N(R₆)₂,
 -S(O)₂-N(R₆)₂,
 10 -N(R₆)-C(R₇)-C₁₋₁₀ alkyl,
 -N(R₆)-C(R₇)-aryl,
 -N(R₆)-S(O)₂-C₁₋₁₀ alkyl,
 -N(R₆)-S(O)₂-aryl,
 -C(O)-C₁₋₁₀ alkyl,
 15 -O-C(R₇)-C₁₋₁₀ alkyl,
 -O-C(R₇)-aryl,
 -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl,
 -O-C(R₇)-N(R₆)-aryl;
 -C(O)-O-C₁₋₁₀ alkyl,
 20 -N₃,
 aryl,
 heteroaryl,
 heterocyclyl,
 -C(O)-aryl, and
 25 -C(O)-heteroaryl;

R₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, carboxy, formyl, aryl, aryloxy, arylalkoxy, heteroaryl, heteroaryloxy, heteroarylalkoxy, heterocyclyl, heterocyclylalkylenyl, amino, alkylamino, (arylalkylenyl)amino, dialkylamino, and in the case of alkyl, alkenyl, alkynyl,

and heterocyclyl, oxo, with the proviso that when R₄ is a substituted alkyl group and the substituent contains a hetero atom which bonds directly to the alkyl group then the alkyl group contains at least two carbons between the substituent and the nitrogen atom to which R₁ is bonded;

5 R₅ is selected from the group consisting of:



each R₆ is independently selected from the group consisting of hydrogen, alkyl, and arylalkylenyl;


each R₇ is independently selected from the group consisting of =O and =S;

10 R₈ is C₂₋₇ alkylene;

A is selected from the group consisting of -CH(R₆)-, -O-, -N(R₆)-, -N(Y-R₄)-, and -N(X-N(R₆)-Y-R₄)-;

X is C₂₋₂₀ alkylene;

Y is selected from the group consisting of -C(R₇)-, -C(R₇)-O-, -S(O)₂-,
15 -S(O)₂-N(R₆)-, and -C(R₇)-N(R₉)-; wherein R₉ is selected from the group consisting of hydrogen, alkyl, and arylalkylenyl; or R₉ and R₄ together with the nitrogen atom to which R₉ is bonded can join to form the group



Z is selected from the group consisting of -O- and -S(O)₀₋₂-; and

20 a and b are independently integers from 1 to 4 with the proviso that when A is -O-, -N(R₆)-, -N(Y-R₄)-, or -N(X-N(R₆)-Y-R₄)- then a and b are independently integers from 2 to 4;

or a pharmaceutically acceptable salt thereof.

25 In one embodiment, the present invention provides a compound selected from the group consisting of:

N-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)cyclopentanecarboxamide;

- 1-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea;
- N*-(3-[4-Amino-1-(isopropylamino)-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-7-yl]phenyl)methanesulfonamide;
- 5 *N*¹-Isopropyl-7-[3-(morpholin-4-ylcarbonyl)phenyl]-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
- N*-(3-{[4-Amino-2-(ethoxymethyl)-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-7-yl]oxy}propyl)methanesulfonamide;
- N*¹-Isopropyl-7-phenyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
- 10 *N*-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide;
- N*-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;
- N*¹-Isopropyl-2-propyl-7-(2-pyridin-3-ylethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
- 15 2-Ethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
- 1-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea;
- N*-(3-[(4-Amino-2-ethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl)acetamide;
- [4-Amino-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol;
- 20 1-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea;
- N*-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)morpholine-4-carboxamide;
- N*-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide;
- 25 *N*-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;
- 1-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea;
- 30 *N*-(3-[(4-Amino-2-ethyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl)acetamide; and

[4-Amino-1-(isopropylamino)-6,7,8,9-tetrahydro-1H-imidazo[4,5-c]quinolin-2-yl]methanol;

or a pharmaceutically acceptable salt thereof.

For certain embodiments, the compound or salt is selected from the group consisting of:

- 5 *N*-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)cyclopentanecarboxamide;
- 1-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea;
- 10 *N*-(3-[4-Amino-1-(isopropylamino)-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-7-yl]phenyl)methanesulfonamide;
- N*¹-Isopropyl-7-[3-(morpholin-4-ylcarbonyl)phenyl]-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
- N*-(3-{[4-Amino-2-(ethoxymethyl)-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-7-yl]oxy}propyl)methanesulfonamide;
- 15 *N*¹-Isopropyl-7-phenyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
- N*-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide;
- N*-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;
- 20 *N*¹-Isopropyl-2-propyl-7-(2-pyridin-3-ylethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
- 2-Ethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
- 1-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea;
- 25 *N*-(3-[(4-Amino-2-ethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl)acetamide;
- [4-Amino-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol; and
- pharmaceutically acceptable salts thereof.

For certain embodiments, the compound or salt is *N*-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide or a pharmaceutically acceptable salt thereof.

For certain embodiments, the compound or salt is selected from the group consisting of:

1-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea;

N-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)morpholine-4-carboxamide;

5 *N*-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide;

N-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;

10 1-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea;

N-(3-{[4-Amino-2-ethyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;

[4-Amino-1-(isopropylamino)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol; and pharmaceutically acceptable salts thereof.

15 In one embodiment of Formula I-1, R_1' is selected from the group consisting of alkoxyalkylenyl and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R_1' is bonded.

For certain of the above embodiments of Formula I-1, n is 0. For certain of these
20 embodiments, R_3 is selected from the group consisting of $-Z'-R_4'$, $-Z'-X'-R_4'$, and $-Z'-X'-Y'-R_4'$. For certain of these embodiments, R_3 is selected from the group consisting of hydroxy, 2-(pyridin-3-yl)ethyl, pyridinyl, (hydroxymethyl)pyridinyl, phenyl, (hydroxymethyl)phenyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 3-(methanesulfonylamino)phenyl, 2-(methanesulfonylamino)ethoxy, 3-
25 (methanesulfonylamino)propoxy, and benzyloxy. For certain of these embodiments, R_3 is selected from the group consisting of 2-(pyridin-3-yl)ethyl, pyridinyl, (hydroxymethyl)pyridinyl, phenyl, (hydroxymethyl)phenyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 3-(methanesulfonylamino)phenyl, 2-(methanesulfonylamino)ethoxy, 3-(methanesulfonylamino)propoxy, and benzyloxy.

30 For certain of the above embodiments of Formula I-1, m is 0 and n is 1. For certain of these embodiments, R is halogen or hydroxy.

For certain of the above embodiments of Formula I-1, m and n are each 0.

In one embodiment of Formula II-1, R_1' is selected from the group consisting of alkoxyalkylenyl and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R_1' is bonded.

5 For certain of the above embodiments of Formula II-1 n is 0.

For certain of the above embodiments of Formula I-1 or Formula II-1, R_1 is selected from the group consisting of $-R_4$, $-Y-R_4$, and $-X-N(R_6)-Y-R_4$ wherein Y is $-C(R_7)-$, $-S(O)_2-$, or $-C(R_7)-N(R_9)-$.


10 For certain of the above embodiments of Formula I-1 or Formula II-1, R_1 is selected from the group consisting of hydrogen, alkyl, alkenyl, arylalkylenyl, arylalkenylenyl, heteroarylalkylenyl, heteroarylalkenylenyl, aminoalkylenyl, alkoxyalkylenyl, acyl, alkylsulfonylaminoalkylenyl, arylsulfonylaminoalkylenyl, alkylcarbonylaminoalkylenyl, alkylaminocarbonyl, alkylaminocarbonylaminoalkylenyl, arylaminocarbonyl, (arylalkylenyl)aminoalkylenyl, heterocyclylcarbonylaminoalkylenyl, 15 arylaminocarbonylaminoalkylenyl, heteroarylcarbonylaminoalkylenyl, and heteroarylaminocarbonylaminoalkylenyl.

For certain of the above embodiments of Formula I-1 or Formula II-1, R_1 is selected from the group consisting of hydrogen, methyl, isopropyl, butyl, 2-methylpropyl, 1-ethylpropyl, 3-methylbutyl, cyclohexyl, benzyl, 3-phenylpropyl, cinnamyl, furan-2- 20 ylmethyl, and $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{NHR}_{13}$, wherein R_{13} is selected from the group consisting of methylcarbonyl, isopropylcarbonyl, cyclopentylcarbonyl, tetrahydropyran-4-ylcarbonyl, methanesulfonyl, phenylsulfonyl, benzyl, ethylaminocarbonyl, isopropylaminocarbonyl, morpholine-4-carbonyl, phenylaminocarbonyl, pyridin-3-ylcarbonyl, and pyridin-3-ylaminocarbonyl.

25 For certain of the above embodiments of Formula I-1 or Formula II-1, R_2 is selected from the group consisting of hydrogen, alkyl, alkoxyalkylenyl, and hydroxyalkylenyl.

For certain of the above embodiments of Formula I-1 or Formula II-1, R_2 is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, n-butyl, 30 methoxymethyl, 2-methoxyethyl, ethoxymethyl, hydroxymethyl, 2-hydroxyethyl, and 3-hydroxypropyl.

In one embodiment, the present invention provides compounds of the following Formula VII:



VII

5 wherein:

each R_B is independently selected from the group consisting of alkyl, alkoxy, halogen, hydroxy, and trifluoromethyl;

n is an integer from 0 to 4;

R_2 is selected from the group consisting of:

- 10 hydrogen,
 alkyl,
 alkenyl,
 aryl,
 heteroaryl,
 15 heterocyclyl,
 -alkylene-Z-alkyl,
 -alkylene-Z-aryl,
 -alkylene-Z-alkenyl, and
 alkyl or alkenyl substituted by one or more substituents selected from the
 20 group consisting of:
 -OH,
 halogen,
 - $N(R_6)_2$,
 - $C(R_7)-N(R_6)_2$,
 25 - $S(O)_2-N(R_6)_2$,
 - $N(R_6)-C(R_7)-C_{1-10}$ alkyl,
 - $N(R_6)-C(R_7)-aryl$,
 - $N(R_6)-S(O)_2-C_{1-10}$ alkyl,
 - $N(R_6)-S(O)_2-aryl$,


-C(O)-C₁₋₁₀ alkyl,
 -O-C(R₇)-C₁₋₁₀ alkyl,
 -O-C(R₇)-aryl,
 -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl,
 -O-C(R₇)-N(R₆)-aryl;
 -C(O)-O-C₁₋₁₀ alkyl,
 -N₃,
 aryl,
 heteroaryl,
 heterocyclyl,
 -C(O)-aryl, and
 -C(O)-heteroaryl;

each R₆ is independently selected from the group consisting of hydrogen, alkyl, and arylalkylenyl;

R₇ is selected from the group consisting of =O and =S; and
 Z is selected from the group consisting of -O- and -S(O)₀₋₂;
 or a pharmaceutically acceptable salt thereof.

In one embodiment of Formula VII, R₂ is selected from the group consisting of alkyl and alkenyl substituted by one or more substituents selected from the group consisting of -N(R₆)-C(R₇)-aryl, -N(R₆)-S(O)₂-aryl, -O-C(R₇)-C₁₋₁₀ alkyl, -O-C(R₇)-aryl, -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl, and -O-C(R₇)-N(R₆)-aryl.

In one embodiment, the present invention provides intermediate compounds of the following Formula IX:



IX

wherein:

each R_B is independently selected from the group consisting of alkyl, alkoxy, halogen, hydroxy, and trifluoromethyl;

n is an integer from 0 to 4;

R₁' is selected from the group consisting of hydrogen, alkyl, alkoxyalkylenyl, and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R₁' is bonded;

R_1 is selected from the group consisting of:

5

-R₄,

-Y-R₄,

-X-R₅,

-X-N(R₆)-Y-R₄,


-X-C(R₇)-N(R₆)-R₄,

10 -X-O-C(R₇)-N(R₆)-R₄,

-X-S(O)₂-N(R₆)-R₄, and

-X-O-R₄;

or R₁' and R₁ together with the nitrogen atom to which they are bonded can join to form a group selected from the group consisting of:



R_2 is selected from the group consisting of:

hydrogen,
alkyl,
alkenyl,
20 aryl,
heteroaryl,
heterocyclyl,
-alkylene-Z-alkyl,
-alkylene-Z-aryl,
25 -alkylene-Z-alkenyl, and


alkyl or alkenyl substituted by one or more substituents selected from the group consisting of:

30 -OH,
-halogen,
-N(R₆)₂,

5 -C(R₇)-N(R₆)₂,
 -S(O)₂-N(R₆)₂,
 -N(R₆)-C(R₇)-C₁₋₁₀ alkyl,
 -N(R₆)-C(R₇)-aryl,
 -N(R₆)-S(O)₂-C₁₋₁₀ alkyl,
 -N(R₆)-S(O)₂-aryl,
 -C(O)-C₁₋₁₀ alkyl,
 -O-C(R₇)-C₁₋₁₀ alkyl,
 -O-C(R₇)-aryl,
 10 -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl,
 -O-C(R₇)-N(R₆)-aryl;
 -C(O)-O-C₁₋₁₀ alkyl,
 -N₃,
 aryl,
 15 heteroaryl,
 heterocyclyl,
 -C(O)-aryl, and
 -C(O)-heteroaryl;

20 R₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl,
 heteroaryl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and
 heterocyclyl groups can be unsubstituted or substituted by one or more substituents
 independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy,
 halogen, nitro, hydroxy, mercapto, cyano, carboxy, formyl, aryl, aryloxy, arylalkoxy,
 heteroaryl, heteroaryloxy, heteroarylalkoxy, heterocyclyl, heterocyclylalkylenyl, amino,
 25 alkylamino, (arylalkylenyl)amino, dialkylamino, and in the case of alkyl, alkenyl, alkynyl,
 and heterocyclyl, oxo, with the proviso that when R₄ is a substituted alkyl group and the
 substituent contains a hetero atom which bonds directly to the alkyl group then the alkyl
 group contains at least two carbons between the substituent and the nitrogen atom to
 which R₁ is bonded;

30 R₅ is selected from the group consisting of



each R_6 is independently selected from the group consisting of hydrogen, alkyl, and arylalkylenyl;


each R_7 is independently selected from the group consisting of =O and =S;

5 R_8 is C_{2-7} alkylene;

A is selected from the group consisting of $-\text{CH}(R_6)-$, $-\text{O}-$, $-\text{N}(R_6)-$, $-\text{N}(\text{Y}-R_4)-$, and $-\text{N}(\text{X}-\text{N}(R_6)-\text{Y}-R_4)-$;

X is C_{2-20} alkylene;

10 Y is selected from the group consisting of $-\text{C}(R_7)-$, $-\text{C}(R_7)-\text{O}-$, $-\text{S}(\text{O})_2-$, $-\text{S}(\text{O})_2-\text{N}(R_6)-$, and $-\text{C}(R_7)-\text{N}(R_9)-$; wherein R_9 is selected from the group consisting of hydrogen, alkyl, and arylalkylenyl; or R_9 and R_4 together with the nitrogen atom to which R_9 is bonded can join to form the group



Z is selected from the group consisting of $-\text{O}-$ and $-\text{S}(\text{O})_{0-2}-$; and


15 a and b are independently integers from 1 to 4 with the proviso that when A is $-\text{O}-$, $-\text{N}(R_6)-$, $-\text{N}(\text{Y}-R_4)-$, or $-\text{N}(\text{X}-\text{N}(R_6)-\text{Y}-R_4)-$ then a and b are independently integers from 2 to 4;

or a pharmaceutically acceptable salt thereof.

20 In one embodiment of Formula IX, R_1' is selected from the group consisting of alkoxyalkylenyl and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R_1' is bonded.

25 In certain of the above embodiments of Formula IX, R_2 is selected from the group consisting of alkyl and alkenyl substituted by one or more substituents selected from the group consisting of $-\text{N}(R_6)-\text{C}(R_7)-\text{aryl}$, $-\text{N}(R_6)-\text{S}(\text{O})_2-\text{aryl}$, $-\text{O}-\text{C}(R_7)-C_{1-10}$ alkyl, $-\text{O}-\text{C}(R_7)-\text{aryl}$, $-\text{O}-\text{C}(R_7)-\text{N}(R_6)-C_{1-10}$ alkyl, and $-\text{O}-\text{C}(R_7)-\text{N}(R_6)-\text{aryl}$.

The present invention also provides intermediate compounds of the following
Formula X:



5 wherein:

each R_B is independently selected from the group consisting of alkyl, alkoxy, halogen, hydroxy, and trifluoromethyl;

n is an integer from 0 to 4;

R_1' is hydrogen or alkyl;

10 R_{1a} is selected from the group consisting of:

- R_{4a} ,

-Y- R_{4a} ,


-X- R_5 ,

-X-N(R_6)-Y- R_{4a} ,

15 -X-C(R_7)-N(R_6)- R_{4a} , and

-X-O- R_{4a} ;

or R_1' and R_{1a} together with the nitrogen atom to which they are bonded can join to form a group selected from the group consisting of:



20 R_{2a} is selected from the group consisting of:

-hydrogen,

-alkyl,

-alkenyl,

-aryl,

25 -alkylene-Z''-alkyl,

-alkylene-Z''-aryl,

-alkylene-Z''-alkenyl, and


-alkyl or alkenyl substituted by one or more substituents selected from the group consisting of:

- 5
10
15
- OH,
 - halogen,
 - N(R₆)₂,
 - C(R₇)-N(R₆)₂,
 - S(O)₂-N(R₆)₂,
 - N(R₆)-C(R₇)-C₁₋₁₀ alkyl,
 - N(R₆)-S(O)₂-C₁₋₁₀ alkyl,
 - C(O)-C₁₋₁₀ alkyl,
 - C(O)-O-C₁₋₁₀ alkyl,
 - N₃,
 - aryl,
 - heterocyclyl, and
 - C(O)-aryl;

20
25

R_{4a} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, cyano, carboxy, formyl, aryl, aryloxy, arylalkoxy, heterocyclyl, heterocyclylalkylenyl, amino, alkylamino, (arylalkylenyl)amino, dialkylamino, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo, with the proviso that when R_{4a} is a substituted alkyl group and the substituent contains a hetero atom which bonds directly to the alkyl group then the alkyl group contains at least two carbons between the substituent and the nitrogen atom to which R₁ is bonded;

R₅ is selected from the group consisting of



each R₆ is independently selected from the group consisting of hydrogen, alkyl, and arylalkylenyl;


30 each R₇ is independently selected from the group consisting of =O and =S;

R_8 is C_{2-7} alkylene;

A is selected from the group consisting of $-\text{CH}(R_6)-$, $-\text{O}-$, $-\text{N}(R_6)-$, $-\text{N}(\text{Y}-R_4)-$, and $-\text{N}(\text{X}-\text{N}(R_6)-\text{Y}-R_4)-$;

X is C_{2-20} alkylene;

5 Y is selected from the group consisting of $-\text{C}(R_7)-$, $-\text{C}(R_7)-\text{O}-$, $-\text{S}(\text{O})_2-$, $-\text{S}(\text{O})_2-\text{N}(R_6)-$, and $-\text{C}(R_7)-\text{N}(R_9)-$; wherein R_9 is selected from the group consisting of hydrogen, alkyl and arylalkylenyl, or R_9 and R_4 together with the nitrogen atom to which R_9 is bonded can join to form the group




10 Z" is selected from the group consisting of $-\text{O}-$ and $-\text{S}(\text{O})_2-$; and

a and b are independently integers from 1 to 4 with the proviso that when A is $-\text{O}-$, $-\text{N}(R_6)-$, $-\text{N}(\text{Y}-R_4)-$, or $-\text{N}(\text{X}-\text{N}(R_6)-\text{Y}-R_4)-$ then a and b are independently integers from 2 to 4;

or a pharmaceutically acceptable salt thereof.

15 In one embodiment of Formula X, R_1' is selected from the group consisting of alkoxyalkylenyl and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R_1' is bonded.

20 In one embodiment, the present invention provides intermediate compounds of the following Formula XLII:



XLII

wherein:

25 R is selected from the group consisting of alkyl, alkenyl, alkoxy, halogen, fluoroalkyl, hydroxy, amino, alkylamino, and dialkylamino;

1 is 0, 1, or 2;

R₂ is selected from the group consisting of:

- hydrogen,
- alkyl,
- alkenyl,
- 5 aryl,
- heteroaryl,
- heterocyclyl,
- alkylene-Z-alkyl,
- alkylene-Z-aryl,
- 10 -alkylene-Z-alkenyl, and
- alkyl or alkenyl substituted by one or more substituents selected from the group consisting of:
- OH,
- halogen,
- 15 -N(R₆)₂,
- C(R₇)-N(R₆)₂,
- S(O)₂-N(R₆)₂,
- N(R₆)-C(R₇)-C₁₋₁₀ alkyl,
- N(R₆)-C(R₇)-aryl,
- 20 -N(R₆)-S(O)₂-C₁₋₁₀ alkyl,
- N(R₆)-S(O)₂-aryl,
- C(O)-C₁₋₁₀ alkyl,
- O-C(R₇)-C₁₋₁₀ alkyl,
- O-C(R₇)-aryl,
- 25 -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl,
- O-C(R₇)-N(R₆)-aryl;
- C(O)-O-C₁₋₁₀ alkyl,
- N₃,
- aryl,
- 30 -heteroaryl,
- heterocyclyl,
- C(O)-aryl, and

-C(O)-heteroaryl;

each R_6 is independently selected from the group consisting of hydrogen, alkyl, and arylalkylenyl;


R_7 is selected from the group consisting of =O and =S; and

5 Z is selected from the group consisting of -O- and -S(O)₀₋₂;
or a pharmaceutically acceptable salt thereof.

In one embodiment of Formula X, R_2 is selected from the group consisting of alkyl and alkenyl substituted by one or more substituents selected from the group consisting of
10 -N(R_6)-C(R_7)-aryl, -N(R_6)-S(O)₂-aryl, -O-C(R_7)-C₁₋₁₀ alkyl, -O-C(R_7)-aryl,
-O-C(R_7)-N(R_6)-C₁₋₁₀ alkyl, and -O-C(R_7)-N(R_6)-aryl.

In certain of the above embodiments of Formula X, l is 0 or 1. In certain of these embodiments, l is 1.

15 In one embodiment, the present invention provides intermediate compounds of the following Formula XLIII:



XLIII

wherein:

20 R is selected from the group consisting of alkyl, alkenyl, alkoxy, halogen, fluoroalkyl, hydroxy, amino, alkylamino, and dialkylamino;

l is 0, 1 or 2;

R_1' is selected from the group consisting of hydrogen, alkyl, alkoxyalkylenyl, and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between
25 the hydroxy or alkoxy substituent and the nitrogen atom to which R_1' is bonded;


R_1 is selected from the group consisting of:

- R_4 ,

-Y- R_4 ,

- 5
- X-R₅,
 - X-N(R₆)-Y-R₄,
 - X-C(R₇)-N(R₆)-R₄,
 - X-O-C(R₇)-N(R₆)-R₄,
 - X-S(O)₂-N(R₆)-R₄, and
 - X-O-R₄;

or R₁' and R₁ together with the nitrogen atom to which they are bonded can join to form a group selected from the group consisting of:



10 R₂ is selected from the group consisting of:


- 15
- hydrogen,
 - alkyl,
 - alkenyl,
 - aryl,
 - heteroaryl,
 - heterocyclyl,
 - alkylene-Z-alkyl,
 - alkylene-Z-aryl,
 - alkylene-Z-alkenyl, and
 - alkyl or alkenyl substituted by one or more substituents selected from the
- 20 group consisting of:

- 25
- OH,
 - halogen,
 - N(R₆)₂,
 - C(R₇)-N(R₆)₂,
 - S(O)₂-N(R₆)₂,
 - N(R₆)-C(R₇)-C₁₋₁₀ alkyl,
 - N(R₆)-C(R₇)-aryl,
 - N(R₆)-S(O)₂-C₁₋₁₀ alkyl,
 - N(R₆)-S(O)₂-aryl,
- 30

-C(O)-C₁₋₁₀ alkyl,
 -O-C(R₇)-C₁₋₁₀ alkyl,
 -O-C(R₇)-aryl,
 -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl,
 -O-C(R₇)-N(R₆)-aryl;
 -C(O)-O-C₁₋₁₀ alkyl,
 -N₃,
 -aryl,
 -heteroaryl,
 -heterocyclyl,
 -C(O)-aryl, and
 -C(O)-heteroaryl;

R₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, carboxy, formyl, aryl, aryloxy, arylalkoxy, heteroaryl, heteroaryloxy, heteroarylalkoxy, heterocyclyl, heterocyclylalkylenyl, amino, alkylamino, (arylalkylenyl)amino, dialkylamino, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo, with the proviso that when R₄ is a substituted alkyl group and the substituent contains a hetero atom which bonds directly to the alkyl group then the alkyl group contains at least two carbons between the substituent and the nitrogen atom to which R₁ is bonded;

R₅ is selected from the group consisting of



each R₆ is independently selected from the group consisting of hydrogen, alkyl, and arylalkylenyl;

each R₇ is independently selected from the group consisting of =O and =S;


R₈ is C₂₋₇ alkylene;

A is selected from the group consisting of -CH(R₆)-, -O-, -N(R₆)-, -N(Y-R₄)-, and

-N(X-N(R₆)-Y-R₄)-;

X is C₂₋₂₀ alkylene;

Y is selected from the group consisting of -C(R₇)-, -C(R₇)-O-, -S(O)₂-,
-S(O)₂-N(R₆)-, and -C(R₇)-N(R₉)-; wherein R₉ is selected from the group consisting of
5 hydrogen, alkyl, and arylalkylenyl; or R₉ and R₄ together with the nitrogen atom to which
R₉ is bonded can join to form the group



Z is selected from the group consisting of -O- and -S(O)₀₋₂-; and

a and b are independently integers from 1 to 4 with the proviso that when A is
10 -O-, -N(R₆)-, -N(Y-R₄)-, or -N(X-N(R₆)-Y-R₄)- then a and b are independently integers
from 2 to 4;

or a pharmaceutically acceptable salt thereof.

In one embodiment of Formula XLIII, R₁' is selected from the group consisting of
alkoxyalkylenyl and hydroxyalkylenyl wherein the alkylenyl group contains at least two
15 carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which
R₁' is bonded.

In certain of the above embodiments of Formula XLIII, R₂ is selected from the
group consisting of alkyl and alkenyl substituted by one or more substituents selected from
the group consisting of -N(R₆)-C(R₇)-aryl, -N(R₆)-S(O)₂-aryl, -O-C(R₇)-C₁₋₁₀ alkyl,
20 -O-C(R₇)-aryl, -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl, and -O-C(R₇)-N(R₆)-aryl.

In certain of the above embodiments of Formula XLIII, l is 0 or 1. In certain of
these embodiments, l is 0.

For any of the compounds presented herein, each one of the following variables
(e.g., R, R₁', R₁, R₂, R₃, R_A, R_B, m, n, A, X, X', Y, Y', and so on) in any of its
25 embodiments can be combined with any one or more of the other variables in any of their
embodiments as would be understood by one of skill in the art. Each of the resulting
combinations of variables is an embodiment of the present invention.

For certain embodiments, R is selected from the group consisting of alkyl, alkenyl,
alkoxy, halogen, fluoroalkyl, hydroxy, amino, alkylamino, and dialkylamino. For certain
30 embodiments, R is halogen or hydroxy.

For certain embodiments, R_A is selected from the group consisting of halogen, hydroxy, alkyl, alkenyl, haloalkyl, alkoxy, alkylthio, $-NH_2$, $-NH(alkyl)$, and $-N(alkyl)_2$.

For certain embodiments, R_B is selected from the group consisting of alkyl, alkoxy, halogen, hydroxyl, and trifluoromethyl. For certain embodiments, R_B is halogen or hydroxy.

For certain embodiments, R_1' is selected from the group consisting of alkoxyalkylenyl and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R_1' is bonded. For certain of these embodiments, R_1' is hydroxyalkylenyl. For certain embodiments R_1' is 2-hydroxyethyl.

For certain embodiments, R_1 is selected from the group consisting of $-R_4$, $-Y-R_4$, $-X-R_5$, $-X-N(R_6)-Y-R_4$, $-X-C(R_7)-N(R_6)-R_4$, $-X-O-C(R_7)-N(R_6)-R_4$, $-X-S(O)_2-N(R_6)-R_4$, and $-X-O-R_4$.

For certain embodiments, R_1 is selected from the group consisting of $-R_4$, $-Y-R_4$, $-X-R_5$, $-X-N(R_6)-Y-R_4$, $-X-C(R_7)-N(R_6)-R_4$, and $-X-O-R_4$; and R_1' is selected from the group consisting of hydrogen and alkyl.

For certain embodiments, R_1 is selected from the group consisting of $-R_4$, $-Y-R_4$, $-X-R_5$, $-X-N(R_6)-Y-R_4$, $-X-C(R_7)-N(R_6)-R_4$, and $-X-O-R_4$.

For certain embodiments, R_1 is selected from the group consisting of $-R_4$, $-Y-R_4$, and $-X-N(R_6)-Y-R_4$ wherein Y is $-C(R_7)-$, $-S(O)_2-$, or $-C(R_7)-N(R_9)-$.

For certain embodiments, R_1 is selected from the group consisting of hydrogen, alkyl, alkenyl, arylalkylenyl, arylalkenylenyl, heteroarylalkylenyl, heteroarylalkenylenyl, aminoalkylenyl, alkoxyalkylenyl, acyl, alkylsulfonylaminoalkylenyl, arylsulfonylaminoalkylenyl, alkylaminocarbonyl, arylaminocarbonyl, (arylalkylenyl)aminoalkylenyl, and arylaminocarbonylaminoalkylenyl.

For certain embodiments, R_1 is selected from the group consisting of hydrogen, methyl, isopropyl, butyl, 2-methylpropyl, 1-ethylpropyl, 3-methylbutyl, cyclohexyl, benzyl, cinnamyl, furan-2-ylmethyl, and $-CH_2CH_2CH_2-NHR_{13}$, wherein R_{13} is selected from the group consisting of methanesulfonyl, phenylsulfonyl, benzyl, and phenylaminocarbonyl.

For certain embodiments, R_1 is selected from the group consisting of hydrogen, alkyl, alkenyl, arylalkylenyl, arylalkenylenyl, heteroarylalkylenyl, heteroarylalkenylenyl,

aminoalkylenyl, alkoxyalkylenyl, acyl, alkylsulfonylaminoalkylenyl, arylsulfonylaminoalkylenyl, alkylaminocarbonyl, alkylaminocarbonylaminoalkylenyl, arylaminocarbonyl, (arylalkylenyl)aminoalkylenyl, heterocyclylcarbonylaminoalkylenyl, and arylaminocarbonylaminoalkylenyl.

5 For certain embodiments, R_1 is selected from the group consisting of hydrogen, methyl, isopropyl, butyl, 2-methylpropyl, 1-ethylpropyl, 3-methylbutyl, cyclohexyl, benzyl, 3-phenylpropyl, cinnamyl, furan-2-ylmethyl, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{-NHR}_{13}$, wherein R_{13} is selected from the group consisting of methanesulfonyl, phenylsulfonyl, benzyl, isopropylaminocarbonyl, morpholine-4-carbonyl, and phenylaminocarbonyl.

10 For certain embodiments, R_1 is selected from the group consisting of hydrogen, alkyl, alkenyl, arylalkylenyl, arylalkenylenyl, heteroarylalkylenyl, heteroarylalkenylenyl, aminoalkylenyl, alkoxyalkylenyl, acyl, alkylsulfonylaminoalkylenyl, arylsulfonylaminoalkylenyl, alkylcarbonylaminoalkylenyl, alkylaminocarbonyl, alkylaminocarbonylaminoalkylenyl, arylaminocarbonyl, (arylalkylenyl)aminoalkylenyl, 15 heterocyclylcarbonylaminoalkylenyl, arylaminocarbonylaminoalkylenyl, heteroarylcarbonylaminoalkylenyl, and heteroarylaminocarbonylaminoalkylenyl.

For certain embodiments, R_1 is selected from the group consisting of hydrogen, methyl, isopropyl, butyl, 2-methylpropyl, 1-ethylpropyl, 3-methylbutyl, cyclohexyl, benzyl, 3-phenylpropyl, cinnamyl, furan-2-ylmethyl, and 20 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{-NHR}_{13}$, wherein R_{13} is selected from the group consisting of methylcarbonyl, isopropylcarbonyl, cyclopentylcarbonyl, tetrahydropyran-4-ylcarbonyl, methanesulfonyl, phenylsulfonyl, benzyl, ethylaminocarbonyl, isopropylaminocarbonyl, morpholine-4-carbonyl, phenylaminocarbonyl, pyridin-3-ylcarbonyl, and pyridin-3-ylaminocarbonyl.

25 For certain embodiments, R_1 is selected from the group consisting of $-\text{R}_4$, $-\text{Y-R}_4$, $-\text{X-R}_5$, $-\text{X-N(R}_6\text{)-Y-R}_4$, $-\text{X-C(R}_7\text{)-N(R}_6\text{)-R}_4$, and $-\text{X-O-R}_4$; R_1' is selected from the group consisting of hydrogen and alkyl; and R_2 is selected from the group consisting of hydrogen, alkyl, alkenyl, aryl, heteroaryl, heterocyclyl, -alkylene-Z-alkyl, -alkylene-Z-aryl, -alkylene-Z-alkenyl, and -alkyl or alkenyl substituted by one or more 30 substituents selected from the group consisting of $-\text{OH}$, -halogen, $-\text{N(R}_6\text{)}_2$, $-\text{C(R}_7\text{)-N(R}_6\text{)}_2$, $-\text{S(O)}_2\text{-N(R}_6\text{)}_2$, $-\text{N(R}_6\text{)-C(R}_7\text{)-C}_{1-10}$ alkyl, $-\text{N(R}_6\text{)-S(O)}_2\text{-C}_{1-10}$ alkyl, $-\text{C(O)-C}_{1-10}$ alkyl, $-\text{C(O)-O-C}_{1-10}$ alkyl, $-\text{N}_3$, -aryl, -heteroaryl, -heterocyclyl, $-\text{C(O)-aryl}$, and

-C(O)-heteroaryl.

For certain embodiments, R_2 is selected from the group consisting of hydrogen, alkyl, alkenyl, aryl, heteroaryl, heterocyclyl, -alkylene-Z-alkyl, -alkylene-Z-aryl, -alkylene-Z-alkenyl, and alkyl or alkenyl substituted by one or more substituents selected from the group consisting of -OH, halogen, $-N(R_6)_2$, $-C(R_7)-N(R_6)_2$, $-S(O)_2-N(R_6)_2$, $-N(R_6)-C(R_7)-C_{1-10}$ alkyl, $-N(R_6)-C(R_7)-aryl$, $-N(R_6)-S(O)_2-C_{1-10}$ alkyl, $-N(R_6)-S(O)_2-aryl$, $-C(O)-C_{1-10}$ alkyl, $-O-C(R_7)-C_{1-10}$ alkyl, $-O-C(R_7)-aryl$, $-O-C(R_7)-N(R_6)-C_{1-10}$ alkyl, $-O-C(R_7)-N(R_6)-aryl$; $-C(O)-O-C_{1-10}$ alkyl, $-N_3$, aryl, heteroaryl, heterocyclyl, $-C(O)-aryl$, and $-C(O)-heteroaryl$.

For certain embodiments, R_2 is selected from the group consisting of hydrogen, alkyl, alkenyl, aryl, heteroaryl, heterocyclyl, -alkylene-Z-alkyl, -alkylene-Z-aryl, -alkylene-Z-alkenyl, and alkyl or alkenyl substituted by one or more substituents selected from the group consisting of -OH, halogen, $-N(R_6)_2$, $-C(R_7)-N(R_6)_2$, $-S(O)_2-N(R_6)_2$, $-N(R_6)-C(R_7)-C_{1-10}$ alkyl, $-N(R_6)-S(O)_2-C_{1-10}$ alkyl, $-C(O)-C_{1-10}$ alkyl, $-C(O)-O-C_{1-10}$ alkyl, $-N_3$, aryl, heteroaryl, heterocyclyl, $-C(O)-aryl$, and $-C(O)-heteroaryl$.

For certain embodiments, R_2 is selected from the group consisting of hydrogen, alkyl, and alkoxyalkylenyl. For certain embodiments, R_2 is selected from the group consisting of hydrogen, butyl, 2-methoxyethyl, and ethoxymethyl. For certain embodiments, R_2 is selected from the group consisting of hydrogen, methyl, propyl, butyl, 2-methoxyethyl, and ethoxymethyl.

For certain embodiments, R_2 is selected from the group consisting of hydrogen, alkyl, alkoxyalkylenyl, and hydroxyalkylenyl. For certain embodiments, R_2 is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, n-butyl, methoxymethyl, 2-methoxyethyl, ethoxymethyl, hydroxymethyl, 2-hydroxyethyl, and 3-hydroxypropyl.

For certain embodiments, R_3 is selected from the group consisting of $-Z'-R_4'$, $-Z'-X'-R_4'$, $-Z'-X'-Y'-R_4'$, and $-Z'-X'-R_5'$. For certain embodiments, R_3 is selected from the group consisting of $-Z'-R_4'$, $-Z'-X'-R_4'$, and $-Z'-X'-Y'-R_4'$.

For certain embodiments, R_3 is selected from the group consisting of hydroxyl, 2-(pyridin-3-yl)ethyl, pyridinyl, (hydroxymethyl)pyridinyl, phenyl, (hydroxymethyl)phenyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 3-(methanesulfonylamino)phenyl, 2-(methanesulfonylamino)ethoxy, 3-(methanesulfonylamino)propoxy, and benzyloxy. For certain embodiments, R_3 is selected from the group consisting of 2-(pyridin-3-yl)ethyl,

pyridinyl, (hydroxymethyl)pyridinyl, phenyl, (hydroxymethyl)phenyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 3-(methanesulfonylamino)phenyl, 2-(methanesulfonylamino)ethoxy, 3-(methanesulfonylamino)propoxy, and benzyloxy. For certain embodiments, R₃ is selected from the group consisting of 2-(pyridin-3-yl)ethyl, pyridinyl, (hydroxymethyl)pyridinyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 2-(methanesulfonylamino)ethoxy, and benzyloxy.

For certain embodiments, R''' is R or R₃ when n is 1, R or one R and one R₃ when n is 2, R or two R's and one R₃ when n is 3, or R when n is 4. For certain embodiments, R''' is R or R₃ when n is 1, R or one R and one R₃ when n is 2, or R when n is 3 or 4.

For certain embodiments, R''' is R₃. For certain of these embodiments, R₃ is selected from the group consisting of -Z'-R₄', -Z'-X'-R₄', and -Z'-X'-Y'-R₄'.

For certain embodiments, R''' is selected from the group consisting of hydroxyl, 2-(pyridin-3-yl)ethyl, pyridinyl, (hydroxymethyl)pyridinyl, phenyl, (hydroxymethyl)phenyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 3-(methanesulfonylamino)phenyl, 2-(methanesulfonylamino)ethoxy, 3-(methanesulfonylamino)propoxy, and benzyloxy.

For certain embodiments, R''' is selected from the group consisting of 2-(pyridin-3-yl)ethyl, pyridinyl, (hydroxymethyl)pyridinyl, phenyl, (hydroxymethyl)phenyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 3-(methanesulfonylamino)phenyl, 2-(methanesulfonylamino)ethoxy, 3-(methanesulfonylamino)propoxy, and benzyloxy.

For certain embodiments, R''' is selected from the group consisting of 2-(pyridin-3-yl)ethyl, pyridinyl, (hydroxymethyl)pyridinyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 2-(methanesulfonylamino)ethoxy, and benzyloxy.


For certain embodiments, R₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, carboxy, formyl, aryl, aryloxy, arylalkoxy, heteroaryl, heteroaryloxy, heteroarylalkoxy, heterocyclyl, heterocyclylalkylenyl, amino, alkylamino, (arylalkylenyl)amino, dialkylamino, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo, with the proviso that when R₄ is a substituted alkyl group and the substituent contains a hetero atom which bonds directly to

the alkyl group then the alkyl group contains at least two carbons between the substituent and the nitrogen atom to which R₁ is bonded.


For certain embodiments, R₄ is selected from the group consisting of C₁₋₆ alkyl, arylalkylenyl, and heterocyclyl. For certain embodiments, R₄ is C₂₋₆ alkyl.


For certain embodiments, R₄' is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, hydroxyalkyl, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, aryl, aryloxy, arylalkyleneoxy, heteroaryl, heteroaryloxy, heteroarylalkyleneoxy, heterocyclyl, amino, alkylamino, dialkylamino, (dialkylamino)alkyleneoxy, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo. For certain embodiments, R₄' is hydrogen, alkyl, aryl, arylalkylenyl, heteroaryl, or heteroarylalkylenyl wherein the aryl and heteroaryl groups are unsubstituted or substituted by one or more of hydroxyalkyl or alkoxy.

For certain embodiments, R₅ is selected from the group consisting of:



20 For certain embodiments, R_5' is selected from the group consisting of:




For certain embodiments, R₅' is . In certain of these embodiments, V is -C(O)-. In certain of these embodiments, A' is -O- and c and d are each the integer 2.

For certain embodiments, R₆ is selected from the group consisting of hydrogen,
25 alkyl, and arylalkylenyl. For certain embodiments, R₆ is hydrogen.


For certain embodiments, R_7 is selected from the group consisting of $=O$ and $=S$.
 For certain embodiments, R_7 is $=O$.

For certain embodiments, R_8 is C_{2-7} alkylene.

For certain embodiments, R_9 is selected from the group consisting of hydrogen,
 5 alkyl, and arylalkylenyl; or R_9 and R_4 together with the nitrogen atom to which R_9 is
 bonded can join to form the group



For certain embodiments, R_9 and R_4 together with the nitrogen atom to which R_9 is
 bonded join to form the group



For certain embodiments, R_{10} is C_{3-8} alkylene.

For certain embodiments, R_{11} is selected from the group consisting of hydrogen,
 C_{1-10} alkyl, C_{2-10} alkenyl, C_{1-10} alkoxy C_{2-10} alkylenyl, and aryl C_{1-10} alkylenyl. For certain
 15 embodiments, R_{11} is hydrogen.

For certain embodiments, R_{12} is selected from the group consisting of hydrogen
 and alkyl.

For certain embodiments, A is selected from the group consisting of $-\text{CH}(R_6)-$,
 $-O-$, $-\text{N}(R_6)-$, $-\text{N}(Y-R_4)-$, and $-\text{N}(X-\text{N}(R_6)-Y-R_4)-$. For certain embodiments, A is $-O-$.

For certain embodiments, A' is selected from the group consisting of $-\text{CH}_2-$, $-O-$,
 20 $-\text{C}(O)-$, $-\text{S}(O)_{0-2}-$, and $-\text{N}(R_4')-$. For certain embodiments, A' is $-O-$.

For certain embodiments, Q is selected from the group consisting of a bond,
 $-\text{C}(R_7)-$, $-\text{C}(R_7)-\text{C}(R_7)-$, $-\text{S}(O)_2-$, $-\text{C}(R_7)-\text{N}(R_{11})-\text{W}-$, $-\text{S}(O)_2-\text{N}(R_{11})-$, $-\text{C}(R_7)-O-$, and
 $-\text{C}(R_7)-\text{N}(OR_{12})-$. For certain embodiments, Q is $-\text{S}(O)_2-$.

For certain embodiments, V is selected from the group consisting of $-\text{C}(R_7)-$,
 25 $-O-\text{C}(R_7)-$, $-\text{N}(R_{11})-\text{C}(R_7)-$, and $-\text{S}(O)_2-$. For certain embodiments, V is $-\text{C}(R_7)-$.

For certain embodiments, W is selected from the group consisting of a bond,
 $-\text{C}(O)-$, and $-\text{S}(O)_2-$.

For certain embodiments, a and b are independently integers from 1 to 4 with the proviso that when A is -O-, -N(R₆)-, -N(Y-R₄)-, or -N(X-N(R₆)-Y-R₄)- then a and b are independently integers from 2 to 4. For certain embodiments, a and b are each the integer 2.

5 For certain embodiments, c and d are independently integers from 1 to 6 with the proviso that $c + d \leq 7$, and when A' is -O- or -N(R₄')- then c and d are independently integers from 2 to 4. For certain embodiments, c and d are each the integer 2.

For certain embodiments, n is an integer from 0 to 4. For certain embodiments, n is 0. For certain embodiments, n is 1.

10 For certain embodiments, n is 1, and R_B is halogen or hydroxy.

For certain embodiments, m is 0 or 1; with the proviso that when m is 1, then n is 0, 1, or 2. For certain embodiments, m is 0 or 1, and n is 0 or 1. For certain m is 1 and n is 0. For certain embodiments, m is 0, and n is 0.

15 For certain embodiments, m is 0 and n is 1. For certain of these embodiments, R is halogen or hydroxy.

As used herein, the terms "alkyl," "alkenyl," "alkynyl" and the prefix "alk-" are inclusive of both straight chain and branched chain groups and of cyclic groups, i.e. cycloalkyl and cycloalkenyl. Unless otherwise specified, these groups contain from 1 to 20 carbon atoms, with alkenyl groups containing from 2 to 20 carbon atoms, and alkynyl groups containing from 2 to 20 carbon atoms. In some embodiments, these groups have a total of up to 10 carbon atoms, up to 8 carbon atoms, up to 6 carbon atoms, or up to 4 carbon atoms. Cyclic groups can be monocyclic or polycyclic and preferably have from 3 to 10 ring carbon atoms. Exemplary cyclic groups include cyclopropyl, cyclopropylmethyl, cyclopentyl, cyclohexyl, adamantyl, and substituted and unsubstituted bornyl, norbornyl, and norbornenyl.

25 Unless otherwise specified, "alkylene," "-alkylene-", "alkenylene", "-alkenylene-", "alkynylene", and "-alkynylene-" are the divalent forms of the "alkyl", "alkenyl", and "alkynyl" groups defined above. The terms "alkylenyl", "alkenylenyl", and "alkynylenyl" are used when "alkylene", "alkenylene", and "alkynylene", respectively, are substituted. For example, an arylalkylenyl group comprises an "alkylene" moiety to which an aryl group is attached.

30

The term "haloalkyl" is inclusive of alkyl groups that are substituted by one or more halogen atoms, including perfluorinated groups. This is also true of other groups that include the prefix "halo-". Examples of suitable haloalkyl groups are chloromethyl, trifluoromethyl, and the like.

5 The term "aryl" as used herein includes carbocyclic aromatic rings or ring systems. Examples of aryl groups include phenyl, naphthyl, biphenyl, fluorenyl and indenyl.

The term "heteroatom" refers to the atoms O, S, or N.

10 The term "heteroaryl" includes aromatic rings or ring systems that contain at least one ring heteroatom (e.g., O, S, N). Suitable heteroaryl groups include furyl, thienyl, pyridyl, quinoliny, isoquinoliny, indolyl, isoindolyl, triazolyl, pyrrolyl, tetrazolyl, imidazolyl, pyrazolyl, oxazolyl, thiazolyl, benzofuranyl, benzothiophenyl, carbazolyl, benzoxazolyl, pyrimidinyl, benzimidazolyl, quinoxaliny, benzothiazolyl, naphthyridinyl, isoxazolyl, isothiazolyl, purinyl, quinazolinyl, pyrazinyl, 1-oxidopyridyl, pyridazinyl, triazinyl, tetrazinyl, oxadiazolyl, thiadiazolyl, and so on.

15 The term "heterocyclyl" includes non-aromatic rings or ring systems that contain at least one ring heteroatom (e.g., O, S, N) and includes all of the fully saturated and partially unsaturated derivatives of the above mentioned heteroaryl groups. Exemplary heterocyclic groups include pyrrolidinyl, tetrahydrofuranyl, morpholinyl, thiomorpholinyl, piperidinyl, piperazinyl, thiazolidinyl, imidazolidinyl, isothiazolidinyl, tetrahydropyranyl, quinuclidinyl, homopiperidinyl, homopiperazinyl, and the like.

20 The terms "arylene", "heteroarylene", and "heterocyclylene" are the divalent forms of the "aryl", "heteroaryl", and "heterocyclyl" groups defined above. The terms "arylenyl," "heteroarylenyl," and "heterocyclenyl" are used when "arylene", "heteroarylene", and "heterocyclylene", respectively, are substituted. For example, an alkylarylenyl group comprises an arylene moiety to which an alkyl group is attached.

25 The term "acyl" includes alkylcarbonyl and arylcarbonyl. In certain embodiments, "acyl" is alkanoyl or benzoyl.

30 When a group (or substituent or variable) is present more than once in any Formula described herein, each group (or substituent or variable) is independently selected, whether explicitly stated or not. For example, for the formula $-N(R_6)_2$ each R_6 group is independently selected. In another example, when an R_1 and an R_2 group both contain an R_7 group, each R_7 group is independently selected. In a further example, when more than

one $\begin{array}{c} \text{—N—C(R}_7\text{)} \\ | \quad / \\ \text{R}_8 \end{array}$ group is present (i.e., R₅ and R_{5'} both contain a $\begin{array}{c} \text{—N—C(R}_7\text{)} \\ | \quad / \\ \text{R}_8 \end{array}$ group) each R₈ group is independently selected and each R₇ group is independently selected.

The invention is inclusive of the compounds described herein and salts thereof in any of their pharmaceutically acceptable forms, including isomers such as diastereomers and enantiomers, solvates, polymorphs, and the like. In particular, if a compound is optically active, the invention specifically includes each of the compound's enantiomers as well as racemic mixtures of the enantiomers.

Preparation of the Compounds

Compounds of the invention can be prepared according to Reaction Scheme I wherein R, R_{1a}, R_{2a}, and n are as defined above.

In step (1) of Reaction Scheme I, a 4-chloro-3-nitroquinoline of Formula III is reacted with *tert*-butyl carbazate or an alternate carbazate to provide a carbazate compound of Formula IV. The reaction can be carried out by adding *tert*-butyl carbazate to a solution of a compound of Formula III in a suitable solvent such as anhydrous dichloromethane in the presence of a base such as triethylamine. The reaction can be run at ambient temperature. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods. Many compounds of Formula III are known or can be prepared using known synthetic methods, see for example, U.S. Patent Nos. 4,689,338; 5,175,296; 5,367,076; and 5,389,640; and the documents cited therein. *Tertiary*-butyl carbazate is commercially available (for example, from Aldrich, Milwaukee, WI). Many alternate carbazate reagents (for example, benzyl carbazate) may be prepared using known synthetic methods.

In step (2) of Reaction Scheme I a carbazate compound of Formula IV is reduced to provide a compound of Formula V. The reduction can be carried out using a conventional heterogeneous hydrogenation catalyst such as platinum on carbon or palladium on carbon. For some compounds of Formula IV, for example, compounds in which R is halogen, a platinum catalyst is preferred. The reaction can be conveniently carried out on a Parr apparatus in a suitable solvent such as toluene and/or isopropanol. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

Other reduction processes may be used for the reduction in step (2). For example, an aqueous solution of sodium dithionite can be added to a solution or suspension of the compound of Formula IV in a suitable solvent such as ethanol or isopropanol. The reaction can be carried out at an elevated temperature, for example at reflux, or at ambient temperature.

In step (3) of Reaction Scheme I a compound of Formula V is (i) reacted with an acyl halide of Formula $R_{2a}C(O)Cl$ or $R_{2a}C(O)Br$ and then (ii) cyclized to provide a 1*H*-imidazo compound of Formula VI. In part (i) the acyl halide is added to a solution of a compound of Formula V in a suitable solvent such as anhydrous dichloromethane in the presence of a base such as triethylamine. The reaction can be run at a reduced temperature, for example, 0° C, or at ambient temperature. In part (ii) the product of part (i) is heated in an alcoholic solvent in the presence of a base. For example, the product of part (i) is refluxed in ethanol in the presence of excess triethylamine or is heated with methanolic ammonia.

Alternatively, step (3) can be carried out by reacting a compound of Formula V with a carboxylic acid or an equivalent thereof. Suitable equivalents to carboxylic acid include orthoesters and 1,1-dialkoxyalkyl alkanoates. The carboxylic acid or equivalent is selected such that it will provide the desired R_{2a} substituent in a compound of Formula VI. For example, triethyl orthoformate will provide a compound where R_{2a} is hydrogen, and triethyl orthovalerate will provide a compound where R_{2a} is butyl. The reaction can be run in the absence of solvent or in an inert solvent such as anhydrous toluene. The reaction is run with sufficient heating to drive off any alcohol or water formed as a byproduct of the reaction. Optionally a catalyst such as pyridine hydrochloride can be included. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (4) of Reaction Scheme I, the *tert*-butoxycarbonyl or alternate oxycarbonyl group is removed from a 1*H*-imidazo compound of Formula VI by hydrolysis under acidic conditions to provide a 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula VIIa or a salt (for example, hydrochloride salt) thereof. For example, a compound of Formula VI is dissolved in 1.5M HCl in ethanol and heated to reflux. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (5a) of Reaction Scheme I, a 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula VIIa or a salt thereof is treated with a ketone, aldehyde, or corresponding ketal or acetal thereof, under acidic conditions to provide a compound of Formula VIII. For example, a ketone is added to a solution of the hydrochloride salt of a compound of Formula VIIa in a suitable solvent such as isopropanol in the presence of an acid or acid resin, for example, DOWEX W50-X1 acid resin. The ketone, aldehyde, or corresponding ketal or acetal thereof, is selected with R_i and R_{ii} groups that will provide the desired R_{1a} substituent in a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXa. For example, acetone will provide a compound where R_{1a} is isopropyl, and benzaldehyde will provide a compound where R_{1a} is benzyl. The reaction is run with sufficient heating to drive off the water formed as a byproduct of the reaction. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (6) of Reaction Scheme I, a compound of Formula VIII is reduced to provide a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXa. The reaction can be carried out by adding sodium borohydride to a solution of a compound of Formula VIII in a suitable solvent, for example, methanol. The reaction can be run at ambient temperature. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.


Alternatively, in step (5b) of Reaction Scheme I, a 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula VIIa can be treated with a ketone and a borohydride under acidic conditions to provide a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXa. For example, the hydrochloride salt of a 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula VIIa, dissolved in a suitable solvent such as 1,2-dichloroethane, can be treated with a ketone and sodium triacetoxyborohydride at room temperature. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (7) of Reaction Scheme I, a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXa is oxidized to provide an *N*-oxide of Formula Xa using a conventional oxidizing agent that is capable of forming *N*-oxides. The reaction is carried out by treating a solution of a compound of Formula IXa in a suitable solvent such as chloroform or dichloromethane with 3-chloroperoxybenzoic acid at ambient temperature. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (8) of Reaction Scheme I, an N-oxide of Formula Xa is aminated to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula Ia, which is a subgenus of compounds of the Formula I-1. The reaction is carried out in two parts. In part (i) a compound of Formula Xa is reacted with an acylating agent. Suitable acylating agents include alkyl- or arylsulfonyl chlorides (e.g., benzenesulfonyl chloride, methanesulfonyl chloride, and *p*-toluenesulfonyl chloride). In part (ii) the product of part (i) is reacted with an excess of an aminating agent. Suitable aminating agents include ammonia (e.g. in the form of ammonium hydroxide) and ammonium salts (e.g., ammonium carbonate, ammonium bicarbonate, ammonium phosphate). The reaction can be carried out by dissolving a compound of Formula Xa in a suitable solvent such as dichloromethane, adding ammonium hydroxide to the solution, and then adding *p*-toluenesulfonyl chloride. The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Alternatively, the oxidation of step (7) and the amination of step (8) can be carried out sequentially without isolating the product of the oxidation to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula Ia. In step (7), after the 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXa is consumed by reaction with 3-chloroperoxybenzoic acid as described in step (7), the aminating and acylating agents are added to the reaction mixture as in step (8). The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Reaction Scheme I



5 Compounds of the invention can be prepared according to Reaction Scheme II wherein R, R₁, R_{2a} and n are as defined above.

10 In step (1) of Reaction Scheme II, a 1H-imidazo compound of Formula VI is oxidized to provide an N-oxide of Formula XI using the method of step (7) in Reaction Scheme I. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

 In step (2) of Reaction Scheme II, an N-oxide of Formula XI is aminated using the method of step (8) in Reaction Scheme I to provide a 4-amino compound of the Formula XIIa. The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

In step (3) of Reaction Scheme II, the *tert*-butoxycarbonyl or alternate oxycarbonyl group is removed from a 4-amino compound of the Formula XIIa using the method of step (4) in Reaction Scheme I to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula XIIIa or a salt (for example, hydrochloride salt) thereof. The product or a


5 pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (4a) of Reaction Scheme II, a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula XIIIa is treated with a ketone, aldehyde, or corresponding ketal or acetal thereof, using the method of step (5a) in Reaction Scheme I to provide a compound of Formula XIVa. The ketone, aldehyde, or corresponding ketal or acetal thereof, is selected with R_i and R_{ii} groups that will provide the desired R₁ substituent in a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine compound of Formula Ib. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (5) of Reaction Scheme II, a compound of Formula XIVa is reduced to provide a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula Ib using the method of step (6) in Reaction Scheme I. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

Alternatively, in step (4b) of Reaction Scheme II, a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula XIIIa can be treated with a ketone and a borohydride using the method of step (5b) of Reaction Scheme I to provide a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula Ib, which is a subgenus of compounds of the Formula I-1. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

Reaction Scheme II




Compounds of the invention can be prepared according to Reaction Scheme III wherein R, R₁', R_{1a}, R_{2a}, and n are as defined above.

5 In step (1) of Reaction Scheme III, a 4-chloro-3-nitroquinoline of Formula III is reacted with a hydrazino compound of Formula XVa to provide a compound of Formula XVI. The reaction can be carried out by adding the hydrazino compound of Formula XVa to a solution of a compound of Formula III in a suitable solvent such as anhydrous dichloromethane in the presence of a base such as triethylamine. The reaction can be run
10 at ambient temperature. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods. Many hydrazino compounds of Formula XVa are commercially available; others can be readily prepared using known synthetic methods.

In step (2) of Reaction Scheme III, a compound of Formula XVI is reduced to provide a compound of Formula XVII using the methods of step (2) in Reaction Scheme I.
15 The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (3) of Reaction Scheme III, a compound of formula XVII is cyclized using the methods of step (3) in Reaction Scheme I to provide a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXb. The product of step (i) (described in step (3) of Reaction Scheme I) can be isolated to provide a compound of the following formula:




In part (ii) the product of part (i) can be refluxed in suitable solvent such as toluene in the presence of pyridine hydrochloride. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (4) of Reaction Scheme III, a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXb is oxidized to provide an *N*-oxide of Formula X using the method of step (7) in Reaction Scheme I. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (5) of Reaction Scheme III, an *N*-oxide of Formula X is aminated using the method of step (8) in Reaction Scheme I to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula Ic, which is a subgenus of compounds of the Formula I-1. The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Alternatively, the oxidation of step (4) and the amination of step (5) can be carried out sequentially without isolating the product of the oxidation to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula Ic. In step (4), after the 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXb is consumed by reaction with 3-chloroperoxybenzoic acid as described in step (4), the aminating and acylating agents are added to the reaction mixture as in step (5). The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Reaction Scheme III



5 Compounds of the invention can be prepared according to Reaction Scheme IV wherein R , R_1 , R_2 and n are as defined above.

10 In step (1) of Reaction Scheme IV, a 2,4-dichloro-3-nitroquinoline of Formula XVIII is reacted with *tert*-butyl carbazate or an alternate carbazate to provide a carbazate compound of Formula XIX. The reaction can be carried out by adding *tert*-butyl carbazate or an alternate carbazate to a solution of a 2,4-dichloro-3-nitroquinoline of Formula XVIII in a suitable solvent such as anhydrous dichloromethane in the presence of a base such as triethylamine. The reaction can be run at ambient temperature. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods. Many quinolines of Formula XVIII are known or can be prepared using known synthetic methods (see for example, Andre et al., U.S. Patent No. 4,988,815 and references cited therein).

15 In step (2) of Reaction Scheme IV, a carbazate compound of Formula XIX is reduced to provide a 2-chloroquinolin-3-amine of Formula XX using the method of step (2) in Reaction Scheme I. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

20 In step (3) of Reaction Scheme IV, a 2-chloroquinolin-3-amine of Formula XX is reacted with an acyl halide of formula $R_2C(O)Cl$ or $R_2C(O)Br$, or a carboxylic acid or

equivalent thereof, using the methods of step (3) in Reaction Scheme I to provide a 4-chloro-1*H*-imidazo[4,5-*c*]quinoline of Formula XXI. The carboxylic acid or equivalent is selected such that it provides the desired R₂ substituent in compounds of Formula XXI. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (4) of Reaction Scheme IV, the *tert*-butoxycarbonyl or alternate oxycarbonyl group is removed from a 4-chloro-1*H*-imidazo[4,5-*c*]quinoline of Formula XXI using the method of step (4) of Reaction Scheme I to provide a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula XXII or a salt thereof. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (5a) of Reaction Scheme IV, a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula XXII or a salt thereof is treated with a ketone, aldehyde, or corresponding ketal or acetal using the method of step (5a) of Reaction Scheme I to provide a compound of Formula XXIII. The ketone, aldehyde, or corresponding ketal or acetal thereof, is selected with R_i and R_{ii} groups that will provide the desired R₁ substituent in a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula XXIVa. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.


In step (6) of Reaction Scheme IV, a compound of Formula XXIII is reduced using the method of step (6) in Reaction Scheme I to provide a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula XXIVa. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

Alternatively, in step (5b) of Reaction Scheme IV, a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula XXII can be treated with a ketone and a borohydride using the method of step (5b) in Reaction Scheme I to provide a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula XXIVa. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (7) of Reaction Scheme IV, a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula XXIVa is aminated to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Id, which is a subgenus of compounds of the Formula I-1. The reaction is carried out by heating (e.g., 125-175°C) a compound of Formula XXIVa under pressure in a

sealed reactor in the presence of a solution of ammonia in an alkanol. The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Reaction Scheme IV



5

Compounds of the invention can be prepared according to Reaction Scheme V wherein R, R₁, R₂ and n are as defined above.

In step (1) of Reaction Scheme V, a 4-chloro-1H-imidazo[4,5-c]quinoline of Formula XXI is aminated, using the method of step (7) in Reaction Scheme IV, to provide a 4-amino compound of the Formula XII. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (2) of Reaction Scheme V, the *tert*-butoxycarbonyl or alternate oxycarbonyl group is removed from a 4-amino compound of the Formula XII using the method of step (4) of Reaction Scheme I to provide a 1H-imidazo[4,5-c]quinoline-1,4-diamine of

15

Formula XIII or a salt thereof. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.


5 In step (3a) of Reaction Scheme V, a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula XIII or a salt thereof is treated with a ketone, aldehyde, or corresponding ketal or acetal using the method of step (5a) of Reaction Scheme I to provide a compound of Formula XIV. The ketone, aldehyde, or corresponding ketal or acetal thereof, is selected with R_i and R_{ii} groups that will provide the desired R₁ substituent in a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine compound of Formula Id. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

10 In step (4) of Reaction Scheme V, a compound of Formula XIV is reduced using the method of step (6) in Reaction Scheme I to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine compound of Formula Id, which is a subgenus of compounds of the Formula I-1. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

15 Alternatively, in step (3b) of Reaction Scheme V, a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula XIII or a salt thereof can be treated with a ketone and a borohydride using the method of step (5b) in Reaction Scheme I to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine compound of Formula Id. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

20

Reaction Scheme V




5 Compounds of the invention can also be prepared according to Reaction Scheme VI wherein R, R₁', R₁, R₂ and n are as defined above.

In step (1) of Reaction Scheme VI, a 2,4-dichloro-3-nitroquinoline of Formula XVIII is reacted with a hydrazino compound of Formula XV, using the method of step (1) in Reaction Scheme III, to provide a compound of Formula XXV. The product or a
10 pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (2) of Reaction Scheme VI, a compound of Formula XXV is reduced using the method of step (2) in Reaction Scheme I to provide a compound of Formula XXVI. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

15 In step (3) of Reaction Scheme VI, a compound of Formula XXVI is reacted with an acyl halide of formula R₂C(O)Cl or R₂C(O)Br, or a carboxylic acid or equivalent thereof using the methods of step (3) in Reaction Scheme I to provide a 4-chloro-1H-imidazo[4,5-c]quinolin-1-amine compound of Formula XXIV. The carboxylic acid or equivalent is selected such that it provides the desired R₂ substituent in a compound of
20 Formula XXIV. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

In step (4) of Reaction Scheme VI, a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula XXIV is aminated using the method of step (7) in Reaction Scheme IV to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Ie, which is a subgenus of compounds of the Formula I-1. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.



Compounds of the invention can be prepared according to Reaction Scheme VII wherein R, R₁', R_{2a}, R₄, n, and Y are as defined above, and X_a is C₁₋₂₀ alkylene.

In step (1) of Reaction Scheme VII, a 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula VIIa or a salt thereof is treated with a ketal or acetal, containing a protected amino group, using the method of step (5a) of Reaction Scheme I to provide a compound of Formula XXVII. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

The amino ketal or acetal is selected with R₁' and X groups that will provide the desired R₁' and X groups in a 1*H*-imidazo[4,5-*c*]quinolin-1,4-diamine of Formula XXX, XXXI, or XXXII, which are subgenera of compounds of the Formula I-1. For example, *tert*-butyl (3,3-diethoxypropyl)carbamate will provide a compound where R₁' is hydrogen and X is ethylene. The amino group of an amino ketal or acetal can be protected with a

tert-butoxycarbonyl or an alternate oxycarbonyl group. For example, 1-amino-3,3-diethoxypropane can be reacted with di-*tert*-butyl dicarbonate in a suitable solvent such as tetrahydrofuran (THF) in the presence of triethylamine to provide *tert*-butyl (3,3-diethoxypropyl)carbamate.

5 In step (2) of Reaction Scheme VII, a compound of Formula XXVII is reduced using the method of step (6) in Reaction Scheme I to provide a compound of Formula XXVIII, which is a subgenus of compounds of the Formula IX. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

10 In step (3) of Reaction Scheme VII, a compound of Formula XXVIII is oxidized to provide an *N*-oxide of Formula XXIX using the method of step (7) in Reaction Scheme I. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

15 In step (4) of Reaction Scheme VII, an *N*-oxide of Formula XXIX is aminated using the method of step (8) in Reaction Scheme I to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXX, which is a subgenus of compounds of the Formula I-1. The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.


20 In step (5) of Reaction Scheme VII, a the *tert*-butoxycarbonyl or alternate oxycarbonyl group is removed from a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXX using the method of step (4) of Reaction Scheme I to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXXI, which is a subgenus of compounds of the Formula I-1. The product or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.

25 In step (6) of Reaction Scheme VII, a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXXI is converted to a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula XXXII using conventional methods. For example, a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXXI can react with an acid chloride of Formula $R_4C(O)Cl$ to provide a compound of Formula XXXII in which Y is $-C(O)-$. In addition, a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXXI can react with sulfonyl
30 chloride of Formula $R_4S(O)_2Cl$ or a sulfonic anhydride of Formula $(R_4S(O)_2)_2O$ to provide a compound of Formula XXXII in which Y is $-S(O)_2-$. Numerous acid chlorides of Formula $R_4C(O)Cl$, sulfonyl chlorides of Formula $R_4S(O)_2Cl$, and sulfonic anhydrides of

Formula $(R_4S(O)_2)_2O$ are commercially available; others can be readily prepared using known synthetic methods. The reaction can be conveniently carried out by adding the acid chloride of Formula $R_4C(O)Cl$, sulfonyl chloride of Formula $R_4S(O)_2Cl$, or sulfonic anhydride of Formula $(R_4S(O)_2)_2O$ to a cooled solution of a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXXI and a base such as triethylamine in a suitable solvent such as chloroform, dichloromethane, or acetonitrile. The reaction can be carried out at ambient temperature or at a sub-ambient temperature such as 0 °C. The product or pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Ureas of Formula XXXII, where Y is $-C(R_7)-N(R_9)-$, in which R_7 is =O, and R_9 is as defined above, can be prepared by reacting a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXXI with isocyanates of Formula $R_4N=C=O$. Numerous isocyanates of Formula $R_4N=C=O$ are commercially available; others can be readily prepared using known synthetic methods. The reaction can be conveniently carried out by adding the isocyanate of Formula $R_4N=C=O$ to a cooled solution of a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of the Formula XXXI in a suitable solvent such as dichloromethane or chloroform. The reaction can be carried out at ambient temperature or at a sub-ambient temperature such as 0 °C. Alternatively, a compound of Formula XXXI can be treated with a thioisocyanate of Formula $R_4N=C=S$, or a carbamoyl chloride of Formula $R_4N(R_9)-C(O)Cl$ to provide a compound of Formula XXXII, where Y is $-C(S)-N(R_9)-$, in which R_9 is as defined above. The product or pharmaceutically acceptable salt thereof can be isolated using conventional methods.


Reaction Scheme VII



Compounds of the invention can be prepared according to Reaction Scheme VIII where n is as defined above; each R_C is independently selected from the group consisting of hydroxy, alkyl, and alkoxy; and R_{1b} and R_{2b} are a subset of R_1 and R_2 , respectively, as defined above, which do not include those groups that one skilled in the art would recognize as being susceptible to reduction under the acidic hydrogenation conditions in step (1). These susceptible groups include, for example, alkenyl, alkynyl, and aryl groups, and groups bearing nitro substituents.

In step (1) of Reaction Scheme VIII, a 1*H*-imidazo[4,5-*c*]quinolin-4-amine of Formula If is reduced to provide a 6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-4-amine of Formula IIa, which is a subgenus of compounds of the Formula II-1. The reaction can be conveniently carried out by suspending or dissolving a compound of Formula If in trifluoroacetic acid, adding platinum(IV) oxide, and hydrogenating under an atmosphere of hydrogen. The reaction can be carried out in a Parr apparatus. The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Reaction Scheme VIII



Compounds of the invention may be prepared according to Reaction Scheme IX where R_A , R_1 , $R_{1'}$, R_2 , and n is as defined above; and each R_a is independently alkyl. Steps (1) through (4) may be carried out as described in U.S. Patent No. 5,352,784 and documents cited therein. In step (1) the amino group of a compound of Formula XXXIII may be acylated to provide a compound of Formula XXXIV. The reaction may be conveniently carried out by reacting a compound of Formula XXXIII with an alkyl malonyl chloride in the presence of a base such as triethylamine in a suitable solvent such as methylene chloride. The product or a pharmaceutically acceptable salt thereof may be isolated using conventional methods. Certain compounds of Formula XXXIII are commercially available and others can be prepared as described in U.S. Patent No. 5,352,784 and documents cited therein. Alkyl malonyl chlorides are known, some of which are commercially available, and others can be made by known methods.

In step (2) of Reaction Scheme IX, a compound of Formula XXXIV may be cyclized to provide a compound of Formula XXXV. The reaction may be conveniently carried out by adding a solution of a compound of Formula XXXIV in a suitable solvent such as THF to a suspension of sodium hydride (or other base capable of removing a malonyl methylene proton) in a suitable solvent such as THF. The reaction may be run at

an elevated temperature, for example the reflux temperature. The product or a pharmaceutically acceptable salt thereof may be isolated using conventional methods.

5 In step (3) of Reaction Scheme IX, a compound of Formula XXXV may be hydrolyzed and decarboxylated to provide a compound of Formula XXXVI. The reaction may be carried out by conventional methods, for example, by combining a compound of Formula XXXV with an acid, such as hydrochloric acid, with heating. The product may be isolated using conventional methods.

10 In step (4) of Reaction Scheme IX, a compound of Formula XXXVI may be nitrated to provide a compound of Formula XXXVII. The reaction may be carried out under conventional nitration conditions, such as by heating a compound of Formula XXXVI in the presence of nitric acid, preferably in a solvent such as acetic acid. The product or a pharmaceutically acceptable salt thereof may be isolated using conventional methods.

15 In step (5) of Reaction Scheme IX, a compound of Formula XXXVII may be chlorinated to provide a 2,4-dichloro-3-nitro-5,6,7,8-tetrahydroquinoline of Formula XXXVIII. The reaction may be carried out by combining a compound of Formula XXXVII with a conventional chlorinating agent (e.g., phosphorus oxychloride, thionyl chloride, phosgene, oxalyl chloride, or phosphorus pentachloride), optionally in solvent such as *N,N*-dimethylformamide (DMF) or methylene chloride, with heating (e.g., at the
20 reflux temperature). The product or a pharmaceutically acceptable salt thereof may be isolated from the reaction mixture using conventional methods.

In step (6) of Reaction Scheme IX, a 2,4-dichloro-3-nitro-5,6,7,8-tetrahydroquinoline of Formula XXXVIII may be reacted with a hydrazino compound of Formula XV ($\text{H}_2\text{N}-\text{N}(\text{R}_1')(\text{R}_1)$), using the method of step (1) in Reaction Scheme III, to
25 provide a compound of Formula XXXIX. The product or a pharmaceutically acceptable salt thereof may be isolated by conventional methods.


In step (7) of Reaction Scheme IX, a compound of Formula XXXIX may be reduced using the method of step (2) in Reaction Scheme I to provide a compound of Formula XL. The product or a pharmaceutically acceptable salt thereof may be isolated
30 by conventional methods.

In step (8) of Reaction Scheme IX, a compound of Formula XL may be reacted with an acyl halide of formula $\text{R}_2\text{C}(\text{O})\text{Cl}$ or $\text{R}_2\text{C}(\text{O})\text{Br}$, or a carboxylic acid or equivalent

thereof using the methods of step (3) in Reaction Scheme I to provide a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula XLI. The carboxylic acid or equivalent may be selected such that it provides the desired R₂ substituent in a compound of Formula II-1. The product or a pharmaceutically acceptable salt thereof may be isolated by conventional methods.

In step (9) of Reaction Scheme IX, a 4-chloro-1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula XLI may be aminated using the method of step (7) in Reaction Scheme IV to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula II-1. The product or a pharmaceutically acceptable salt thereof may be isolated by conventional methods.

Reaction Scheme IX



5 For some embodiments, compounds of the invention are prepared according to Reaction Scheme X, wherein R , R_{1a} , R_{2a} , and l are as defined above; Hal is chloro, bromo, or iodo; R_{3a} is $-Z'-R_4'$, $-Z'-X'-R_4'$, $-Z'-X'-Y'-R_4'$, or $-Z'-X'-R_5'$; wherein R_4' , Y' , X' , and R_5' are as defined above; and Z' is a bond.

10 In step (1) of Reaction Scheme X, a 4-chloro-3-nitroquinoline of Formula XLIV is converted to a carbazate of Formula XLV according to the method described in step (1) of

Reaction Scheme I. Compounds of Formula XLIV can be readily prepared using known synthetic routes; see for example, U.S. Patent Nos. 4,689,338 (Gerster), 5,367,076 (Gerster), 6,331,539 (Crooks et al.), 6,451,810 (Coleman et al.), 6,541,485 (Crooks et al.) and the documents cited therein.

5 In steps (2) and (3) of Reaction Scheme X, a nitro-substituted quinoline of Formula XLV is first reduced to an amino-substituted quinoline of Formula XLVI, which is then cyclized to a 1*H*-imidazoquinoline of Formula XLVII. Steps (2) and (3) of Reaction Scheme X can be carried out as described for steps (2) and (3) of Reaction Scheme I.

10 In step (4) of Reaction Scheme X, the *tert*-butoxycarbonyl group of a 1*H*-imidazoquinoline of Formula XLVII is hydrolyzed under acidic conditions to provide a 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula VIIb or a pharmaceutically acceptable salt thereof. The reaction is conveniently carried out as described in step (4) of Reaction Scheme I.

15 The 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula VIIb is then converted to a 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula IXc using either a two-step procedure as shown in steps (5a) and (6) of Reaction Scheme X or a one-step procedure as shown in step (5b). The two-step procedure, in which a compound of Formula VIIIb is isolated, can be carried out as described in steps (5a) and (6) of Reaction Scheme I. In step (5a), the ketone, aldehyde, or corresponding ketal or acetal thereof, is selected with R_i and R_{ii} groups that will provide the desired R_{1a} substituent in a 1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula IXc. Step (5b) of Reaction Scheme X can be carried out as described for step (5b) of Reaction Scheme I.

20 In steps (7) and (8) of Reaction Scheme X, a 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula IXc is first oxidized to an *N*-oxide of Formula Xb, which is then aminated to provide a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Ig, which is a subgenus of the compounds of the Formula I-1. Steps (7) and (8) of Reaction Scheme X can be carried out according to the procedures described in steps (7) and (8) of Reaction Scheme I.

30 Step (9) of Reaction Scheme X can be carried out using known palladium-catalyzed coupling reactions such as Suzuki coupling, Stille coupling, Sonogashira coupling, and the Heck reaction. For example, a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Ig undergoes Suzuki coupling with a boronic acid of Formula R_{3a}-B(OH)₂, an anhydride thereof, or a boronic acid ester of Formula R_{3a}-B(O-alkyl)₂ to provide an 1*H*-

imidazo[4,5-*c*]quinoline-1,4-diamine of Formula I-1b, a subgenus of Formula I-1, wherein R_{3a} is -Z'-R₄', -Z'-X'-R₄', -Z'-X'-Y'-R₄', or -Z'-X'-R₅'; -Z' is a bond; -X'- is alkenylene, arylene, heteroarylene, or alkenylene optionally terminated by arylene or heteroarylene; and R₄', Y', and R₅' are as defined above. The coupling is carried out by combining a
5 compound of Formula Ig with a boronic acid or an ester or anhydride thereof in the presence of palladium (II) acetate, triphenylphosphine, and a base such as sodium carbonate in a suitable solvent such as *n*-propanol. The reaction can be carried out at an elevated temperature (e.g., 80-100°C). Numerous boronic acids of Formula R_{3a} -B(OH)₂, anhydrides thereof, and boronic acid esters of Formula R_{3a} -B(O-alkyl)₂ are commercially
10 available; others can be readily prepared using known synthetic methods. See, for example, Li, W. et al, *J. Org. Chem.*, 67, 5394-5397 (2002). The product of Formula I-1b or a pharmaceutically acceptable salt thereof can be isolated by conventional methods.


The Heck reaction can also be used in step (9) of Reaction Scheme X to provide compounds of Formula I-1b, wherein R_{3a} is -Z'-X'-R₄' or -Z'-X'-Y'-R₄'; -Z' is a bond; -X'-
15 is alkenylene optionally terminated by arylene or heteroarylene; and R₄' and Y' are as defined above. The Heck reaction is carried out by coupling a 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Ig with a vinyl-substituted arylene or heteroarylene compound. Several vinyl-substituted arylene or heteroarylene compounds, such as 2-
vinylpyridine, 3-vinylpyridine, and 4-vinylpyridine, are commercially available; others
20 can be prepared by known methods. The reaction is conveniently carried out by combining the 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Ig and the vinyl-substituted compound in the presence of palladium (II) acetate, triphenylphosphine or tri-
ortho-tolylphosphine, and a base such as triethylamine in a suitable solvent such as acetonitrile or toluene. The reaction can be carried out at an elevated temperature such as
25 100-120 °C under an inert atmosphere. The compound or pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Compounds of Formula I-1b, wherein R_{3a} is -Z'-X'-R₄' or -Z'-X'-Y'-R₄', -Z' is a bond and -X'- is alkenylene optionally terminated by arylene or heteroarylene, may be
30 reduced to provide compounds wherein -X'- is alkylene optionally terminated by arylene or heteroarylene. For example, compounds wherein R_{3a} is a 2-(pyridin-3-yl)ethyl group may be prepared in this manner. The reduction can be carried out by hydrogenation using a conventional heterogeneous hydrogenation catalyst such as palladium on carbon. The

reaction can conveniently be carried out on a Parr apparatus in a suitable solvent such as ethanol, methanol, or mixtures thereof. The compound or pharmaceutically acceptable salt thereof can be isolated using conventional methods.

5

Reaction Scheme X



For some embodiments, compounds of the invention can be prepared according to Reaction Scheme XI where R, R_{1a}, R_{2a}, and l are as defined above; Boc is *tert*-

butoxycarbonyl; R_{3b} is $-Z'-R_4'$, $-Z'-X'-R_4'$, $-Z'-X'-Y'-R_4'$, or $-Z'-X'-R_5'$; X' , Y' , and R_4' are as defined above; and Z' is $-O-$.

5 In step (1) of Reaction Scheme XI, a benzyloxyaniline of Formula XLVIII is treated with the condensation product generated from 2,2-dimethyl-1,3-dioxane-4,6-dione (Meldrum's acid) and triethyl orthoformate to provide an imine of Formula XLIX. The reaction is conveniently carried out by adding a solution of a benzyloxyaniline of Formula XLVIII to a heated mixture of Meldrum's acid and triethyl orthoformate and heating the reaction at an elevated temperature such as 45 °C. The product can be isolated using conventional methods.

10 In step (2) of Reaction Scheme XI, an imine of Formula XLIX undergoes thermolysis and cyclization to provide a benzyloxyquinolin-4-ol of Formula L. The reaction is conveniently carried out in a heat transfer fluid such as DOWTHERM A heat transfer fluid at a temperature between 200 and 250 °C. The product can be isolated using conventional methods.

15 In step (3) of Reaction Scheme XI, a benzyloxyquinolin-4-ol of Formula L is nitrated under conventional nitration conditions to provide a benzyloxy-3-nitroquinolin-4-ol of Formula LI. The reaction is conveniently carried out by adding nitric acid to the benzyloxyquinolin-4-ol of Formula L in a suitable solvent such as propionic acid and heating the mixture at an elevated temperature such as 125 °C. The product can be
20 isolated using conventional methods.

In step (4) of Reaction Scheme XI, a benzyloxy-3-nitroquinolin-4-ol of Formula LI is chlorinated using conventional chlorination chemistry to provide a benzyloxy-4-chloro-3-nitroquinoline of Formula LII. The reaction is conveniently carried out by treating the benzyloxy-3-nitroquinolin-4-ol of Formula LI with phosphorous oxychloride in a suitable
25 solvent such as DMF. The reaction can be carried out at ambient temperature or at an elevated temperature such as 100 °C, and the product can be isolated using conventional methods.

In step (5) of Reaction Scheme XI, a benzyloxy-4-chloro-3-nitroquinoline of Formula LII is converted to a carbazate of Formula LIII. The reaction is conveniently
30 carried out as described in step (1) of Reaction Scheme I.

In steps (6) and (7) of Reaction Scheme XI, a nitro-substituted quinoline of Formula LIII is first reduced to an amino-substituted quinoline of Formula LIV, which is

then cyclized to a benzyloxy-1*H*-imidazo[4,5-*c*]quinoline of Formula LV. Steps (6) and (7) of Reaction Scheme XI can be carried out as described for steps (2) and (3) of Reaction Scheme I.

5 In step (8) of Reaction Scheme XI, the Boc group of a benzyloxy-1*H*-imidazo[4,5-*c*]quinoline of Formula LV is hydrolyzed under acidic conditions to provide a benzyloxy-1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula XLIIa or a pharmaceutically acceptable salt thereof. The reaction is conveniently carried out as described in step (4) of Reaction Scheme I.

10 The benzyloxy-1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula XLIIa is then converted to a benzyloxy-1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula XLIIIa using either a two-step procedure as shown in steps (9a) and (10) of Reaction Scheme XI or a one-step procedure as shown in step (9b). The two-step procedure, in which a compound of Formula LVI is isolated, can be carried out as described in steps (5a) and (6) of Reaction Scheme I. In step (9a), the ketone, aldehyde, or corresponding ketal or acetal
15 thereof, is selected with R_i and R_{ii} groups that will provide the desired R_{1a} substituent in a benzyloxy-1*H*-imidazo[4,5-*c*]quinolin-1-amine compound of Formula XLIIIa. Step (9b) of Reaction Scheme XI can be carried out as described for step (5b) of Reaction Scheme I.

In steps (11) and (12) of Reaction Scheme XI, a benzyloxy-1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula XLIIIa is first oxidized to an *N*-oxide of Formula LVII, which is then aminated to provide a benzyloxy-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula LVIII, which is a subgenus of the compounds of the Formula I-1. Steps (11) and (12) of Reaction Scheme XI can be carried out according to the procedures described in steps (7) and (8) of Reaction Scheme I.


25 In step (13) of Reaction Scheme XI, the benzyl group of a benzyloxy-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula LVIII is cleaved to provide a hydroxy-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Ih. The cleavage is conveniently carried out on a Parr apparatus under hydrogenolysis conditions using a suitable heterogeneous catalyst such as palladium on carbon in a solvent such as ethanol. The product or pharmaceutically acceptable salt thereof can be isolated using conventional
30 methods.

In step (14) of Reaction Scheme XI a hydroxy-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Ih is converted to an ether-substituted 1*H*-imidazo[4,5-*c*]quinoline-

1,4-diamine of Formula I-1c (a subgenus of compounds of Formula I-1) using a Williamson-type ether synthesis. The reaction is effected by treating a compound of Formula Ih with an alkyl halide of Formula Halide-R₄', Halide-X'-Y'-R₄', Halide-X'-R₄', or Halide-X'-R₅' in the presence of a base. The reaction is conveniently carried out by combining the alkyl halide with a compound of Formula Ih in a solvent such as DMF in the presence of a suitable base such as cesium carbonate. The reaction can be carried out at ambient temperature or at an elevated temperature, for example 65 °C or 85 °C. Alternatively, the reaction can be carried out by treating a solution of a compound of Formula Ih in a solvent such as DMF with sodium hydride and then adding the alkyl halide. The product or pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Numerous reagents of Formulas Halide-R₄', Halide-X'-R₄', and Halide-X'-Y'-R₄' are commercially available, for example, bromo-substituted ketones, esters, and heterocycles. Other reagents of Formulas Halide-R₄', Halide-X'-Y'-R₄', or Halide-X'-R₅' can be prepared using conventional synthetic methods; for example, a bromo-substituted acid halide of Formula ClC(O)-X'-Br can be treated with a secondary amine in a suitable solvent such as dichloromethane to provide a variety of bromo-substituted amides of Formula

Br-X'-C(O)-N(R₁₁)-R₄' or



The reaction can be run at a sub-ambient temperature such as -25 °C, and the product or pharmaceutically acceptable salt thereof can be isolated using conventional methods.


Reagents of Formula I-X'-NH-C(O)-O-C(CH₃)₃ can be prepared in two steps from amino alcohols of Formula HO-X'-NH₂, many of which are commercially available or readily prepared by known synthetic methods. An amino alcohol of Formula HO-X'-NH₂ is first protected with a *tert*-butoxy carbonyl group by treating the amino alcohol with di-*tert*-butyl dicarbonate in the presence of a base such as aqueous sodium hydroxide in a suitable solvent such as tetrahydrofuran. The resulting hydroxyalkylcarbamate of Formula HO-X'-NH-C(O)-O-C(CH₃)₃ is then treated with a solution of iodine, triphenylphosphine,

and imidazole at ambient temperature in a suitable solvent such as dichloromethane. The product of Formula I-X'-NH-C(O)-O-C(CH₃)₃ can be isolated using conventional methods.

Step (14) of Reaction Scheme XI can alternatively be carried out by treating a hydroxy-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula Ih with an alcohol of
5 Formula HO-X'-Y'-R₄', HO-X'-R₅', HO-X'-R₄', or HO-R₄' under Mitsunobu reaction conditions. Numerous alcohols of these formulas are commercially available, and others can be prepared using conventional synthetic methods. The reaction is conveniently carried out by adding triphenylphosphine and an alcohol of Formula HO-X'-Y'-R₄', HO-X'-R₅', HO-X'-R₄', or HO-R₄' to a solution of a compound of Formula Ih in a suitable
10 solvent such as tetrahydrofuran and then slowly adding diisopropyl azodicarboxylate or diethyl azodicarboxylate. The reaction can be carried out at ambient temperature or at a sub-ambient temperature, such as 0 °C. The product or pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Compounds of Formula I-1c, wherein R_{3b} is -O-X'-NH-C(O)-O-C(CH₃)₃, can be
15 prepared by treating compounds of Formula Ih with alcohols such as *tert*-butyl *N*-(4-hydroxybutyl)carbamate and *tert*-butyl *N*-(5-hydroxypentyl)carbamate under Mitsunobu conditions or with alkyl halides of Formula I-X'-NH-C(O)-O-C(CH₃)₃ in a Williamson-type ether synthesis. These compounds of Formula I-1c, wherein R_{3b} is -O-X'-NH-C(O)-O-C(CH₃)₃, are then readily converted to other compounds of Formula
20 I-1c using conventional synthetic methods. For example, compounds in which R_{3b} is -O-X'-NH-C(O)-O-C(CH₃)₃ can be deprotected and treated according to the methods described in steps (5) and (6) of Reaction Scheme VII, Parts F and G of Example 14, and Examples 15 and 23 to provide compounds of Formula I-1c wherein R_{3b} is -Z'-X'-Y'-R₄'; Z' is -O-; Y' is -NH-Q-; Q is -C(R₇)-, -S(O)₂-, or -C(R₇)-N(R₁₁)-; and X', R₄', R₇, and R₁₁
25 are as defined above. Compounds in which R_{3b} is a 2-methanesulfonylaminoethoxy group or a 3-methanesulfonylaminoethoxy group are available using these methods.


Reaction Scheme XI



For some embodiments, compounds of Formula I-1c can be prepared according to Reaction Scheme XII, in which R, R_{1a}, R_{2a}, R_{3b}, and l are as defined above. In step (1) of Reaction Scheme XII, the benzyl group of a benzyloxy-1H-imidazo[4,5-c]quinolin-1-amine of Formula XLIIa is cleaved to provide a hydroxy-1H-imidazo[4,5-c]quinolin-1-amine of Formula IXd. In step (2) of Reaction Scheme XII a hydroxy-1H-imidazo[4,5-c]quinolin-1-amine of Formula IXd is converted to an ether-substituted 1H-imidazo[4,5-c]quinolin-1-amine of Formula LIX. In steps (3) and (4) of Reaction Scheme XII, an

ether-substituted 1*H*-imidazo[4,5-*c*]quinolin-1-amine of Formula LIX is first oxidized to an *N*-oxide of Formula LX, which is then aminated to provide an ether-substituted 1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine of Formula I-1c, which is a subgenus of the compounds of Formula I-1. Steps (1), (2), (3), and (4) of Reaction Scheme XII can be carried out as described in steps (13), (14), (11), and (12), respectively, of Reaction Scheme XI.

Reaction Scheme XII



Pharmaceutical Compositions and Biological Activity

Pharmaceutical compositions of the invention contain a therapeutically effective amount of a compound of the invention as described above in combination with a pharmaceutically acceptable carrier.

The term "a therapeutically effective amount" or "effective amount" means an amount of the compound sufficient to induce a therapeutic or prophylactic effect, such as cytokine induction, cytokine inhibition, immunomodulation, antitumor activity, and/or antiviral activity. Although the exact amount of active compound used in a pharmaceutical composition of the invention will vary according to factors known to those of skill in the art, such as the physical and chemical nature of the compound, the nature of the carrier, and the intended dosing regimen, it is anticipated that the compositions of the invention will contain sufficient active ingredient to provide a dose of about 100 nanograms per kilogram (ng/kg) to about 50 milligrams per kilogram (mg/kg), preferably

about 10 micrograms per kilogram ($\mu\text{g/kg}$) to about 5 mg/kg, of the compound to the subject. A variety of dosage forms may be used, such as tablets, lozenges, capsules, parenteral formulations, syrups, creams, ointments, aerosol formulations, transdermal patches, transmucosal patches and the like.

5 The compounds of the invention can be administered as the single therapeutic agent in the treatment regimen, or the compounds of the invention may be administered in combination with one another or with other active agents, including additional immune response modifiers, antivirals, antibiotics, antibodies, proteins, peptides, oligonucleotides, etc.

10 Compounds of the invention have been shown to induce, and certain compounds of the invention may inhibit, the production of certain cytokines in experiments performed according to the tests set forth below. These results indicate that the compounds are useful as immune response modifiers that can modulate the immune response in a number of different ways, rendering them useful in the treatment of a variety of disorders.

15 Cytokines whose production may be induced by the administration of compounds according to the invention generally include interferon- α (IFN- α) and/or tumor necrosis factor- α (TNF- α) as well as certain interleukins (IL). Cytokines whose biosynthesis may be induced by compounds of the invention include IFN- α , TNF- α , IL-1, IL-6, IL-10 and IL-12, and a variety of other cytokines. Among other effects, these and other cytokines
20 can inhibit virus production and tumor cell growth, making the compounds useful in the treatment of viral diseases and neoplastic diseases. Accordingly, the invention provides a method of inducing cytokine biosynthesis in an animal comprising administering an effective amount of a compound or composition of the invention to the animal. The animal to which the compound or composition is administered for induction of cytokine
25 biosynthesis may have a disease as described *infra*, for example a viral disease or a neoplastic disease, and administration of the compound may provide therapeutic treatment. Alternatively, the compound may be administered to the animal prior to the animal acquiring the disease so that administration of the compound may provide a prophylactic treatment.

30 In addition to the ability to induce the production of cytokines, compounds of the invention may affect other aspects of the innate immune response. For example, natural killer cell activity may be stimulated, an effect that may be due to cytokine induction. The

compounds may also activate macrophages, which in turn stimulate secretion of nitric oxide and the production of additional cytokines. Further, the compounds may cause proliferation and differentiation of B-lymphocytes.

5 Compounds of the invention can also have an effect on the acquired immune response. For example, the production of the T helper type 1 (T_H1) cytokine IFN- γ may be induced indirectly and the production of the T helper type 2 (T_H2) cytokines IL-4, IL-5 and IL-13 may be inhibited upon administration of the compounds.

10 Other cytokines whose production may be inhibited by the administration of certain compounds according to the invention include tumor necrosis factor- α (TNF- α). Among other effects, inhibition of TNF- α production can provide prophylaxis or therapeutic treatment of diseases in animals in which TNF is mediated, making the compounds useful in the treatment of, for example, autoimmune diseases. Accordingly, the invention provides a method of inhibiting TNF- α biosynthesis in an animal comprising administering an effective amount of a compound or composition of the invention to the
15 animal. The animal to which the compound or composition is administered for inhibition of TNF- α biosynthesis may have a disease as described *infra*, for example an autoimmune disease, and administration of the compound may provide therapeutic treatment. Alternatively, the compound may be administered to the animal prior to the animal acquiring the disease so that administration of the compound may provide a prophylactic
20 treatment.

Whether for prophylaxis or therapeutic treatment of a disease, and whether for effecting innate or acquired immunity, the compound or composition may be administered alone or in combination with one or more active components as in, for example, a vaccine adjuvant. When administered with other components, the compound and other component
25 or components may be administered separately; together but independently such as in a solution; or together and associated with one another such as (a) covalently linked or (b) non-covalently associated, e.g., in a colloidal suspension.

Conditions for which IRMs identified herein may be used as treatments include, but are not limited to:

30 (a) viral diseases such as, for example, diseases resulting from infection by an adenovirus, a herpesvirus (e.g., HSV-I, HSV-II, CMV, or VZV), a poxvirus (e.g., an orthopoxvirus such as variola or vaccinia, or molluscum contagiosum), a picornavirus

(e.g., rhinovirus or enterovirus), an orthomyxovirus (e.g., influenzavirus), a paramyxovirus (e.g., parainfluenzavirus, mumps virus, measles virus, and respiratory syncytial virus (RSV), a coronavirus (e.g., SARS), a papovavirus, (e.g., papillomaviruses, such as those that cause genital warts, common warts, or plantar warts), a hepadnavirus (e.g., hepatitis B virus), a flavivirus (e.g., hepatitis C virus or Dengue virus), or a retrovirus (e.g., a lentivirus such as HIV);

(b) bacterial diseases, such as, for example, diseases resulting from infection by bacteria of, for example, the genus *Escherichia*, *Enterobacter*, *Salmonella*, *Staphylococci*, *Shigella*, *Listeria*, *Aerobacter*, *Helicobacter*, *Klebsiella*, *Proteus*, *Pseudomonas*, *Streptococcus*, *Chlamydia*, *Mycoplasma*, *Pneumococcus*, *Neisseria*, *Clostridium*, *Bacillus*, *Corynebacterium*, *Mycobacterium*, *Campylobacter*, *Vibrio*, *Serratia*, *Providencia*, *Chromobacterium*, *Brucella*, *Yersinia*, *Haemophilus*, or *Bordetella*;

(c) other infectious diseases, such as chlamydia, fungal diseases, such as, for example, candidiasis, aspergillosis, histoplasmosis, cryptococcal meningitis, or parasitic diseases, such as, for example, malaria, pneumocystis carinii pneumonia, leishmaniasis, cryptosporidiosis, toxoplasmosis, and trypanosome infection;

(d) neoplastic diseases, such as intraepithelial neoplasias, cervical dysplasia, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, aenal cell leukemia, Karposi's sarcoma, melanoma, renal cell carcinoma, leukemias, such as, for example, myelogenous leukemia, chronic lymphocytic leukemia, and multiple myeloma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, B-cell lymphoma, hairy cell leukemia, and other cancers; and

(e) T_H2 -mediated, atopic, and autoimmune diseases, such as atopic dermatitis or eczema, eosinophilia, asthma, allergy, allergic rhinitis, systemic lupus erythematosus, essential thrombocythaemia, multiple sclerosis, Ommen's syndrome, discoid lupus, alopecia areata, inhibition of keloid formation and other types of scarring, and enhancing wound healing, including chronic wounds.

IRMs identified herein also may be useful as a vaccine adjuvant for use in conjunction with any material that raises either humoral and/or cell mediated immune response, such as, for example, live viral, bacterial, or parasitic immunogens; inactivated viral, tumor-derived, protozoal, organism-derived, fungal, or bacterial immunogens, toxoids, toxins; self-antigens; polysaccharides; proteins; glycoproteins; peptides; cellular

vaccines; DNA vaccines; recombinant proteins; and the like, for use in connection with, for example, BCG, cholera, plague, typhoid, hepatitis A, hepatitis B, and hepatitis C, influenza A and influenza B, parainfluenza, polio, rabies, measles, mumps, rubella, yellow fever, tetanus, diphtheria, hemophilus influenza b, tuberculosis, meningococcal and
5 pneumococcal vaccines, adenovirus, HIV, chicken pox, cytomegalovirus, dengue, feline leukemia, fowl plague, HSV-1 and HSV-2, hog cholera, Japanese encephalitis, respiratory syncytial virus, rotavirus, papilloma virus, yellow fever, and Alzheimer's Disease.

IRMs identified herein may also be particularly helpful in individuals having compromised immune function. For example, IRM compounds may be used for treating
10 the opportunistic infections and tumors that occur after suppression of cell mediated immunity in, for example, transplant patients, cancer patients and HIV patients.

Thus, one or more of the above diseases or types of diseases, for example, a viral disease or a neoplastic disease may be treated in an animal in need thereof (having the disease) by administering a therapeutically effective amount of a compound or salt of
15 Formula I-1, II-1, any of the embodiments described herein, or a combination thereof to the animal. An animal may also be vaccinated by administering an effective amount of a compound or salt of Formula I-1, II-1, any of the embodiments described herein, or a combination thereof to the animal as a vaccine adjuvant.

An amount of a compound effective to induce cytokine biosynthesis is an amount
20 sufficient to cause one or more cell types, such as monocytes, macrophages, dendritic cells and B-cells to produce an amount of one or more cytokines such as, for example, IFN- α , TNF- α , IL-1, IL-6, IL-10 and IL-12 that is increased over the background level of such cytokines. The precise amount will vary according to factors known in the art but is expected to be a dose of about 100 ng/kg to about 50 mg/kg, preferably about 10 μ g/kg to
25 about 5 mg/kg. The invention also provides a method of treating a viral infection in an animal and a method of treating a neoplastic disease in an animal comprising administering an effective amount of a compound or composition of the invention to the animal.

An amount effective to treat or inhibit a viral infection is an amount that will cause
30 a reduction in one or more of the manifestations of viral infection, such as viral lesions, viral load, rate of virus production, and mortality as compared to untreated control animals. The precise amount that is effective for such treatment will vary according to

factors known in the art but is expected to be a dose of about 100 ng/kg to about 50 mg/kg, preferably about 10 µg/kg to about 5 mg/kg. An amount of a compound effective to treat a neoplastic condition is an amount that will cause a reduction in tumor size or in the number of tumor foci. Again, the precise amount will vary according to factors known in the art but is expected to be a dose of about 100 ng/kg to about 50 mg/kg, preferably about 10 µg/kg to about 5 mg/kg.


In certain embodiments, there is provided a method of inducing cytokine biosynthesis in an animal comprising administering an effective amount of a compound or salt described herein to the animal. In another embodiment, there is provided a method of treating a viral disease in an animal comprising administering a therapeutically effective amount of a compound or salt described herein to the animal. In another embodiment, there is provided a method of treating a neoplastic disease in an animal comprising administering a therapeutically effective amount of a compound or salt described herein to the animal. In another embodiment, there is provided a method of vaccinating an animal comprising administering an effective amount of a compound or salt described herein to the animal as a vaccine adjuvant.

EXAMPLES

Objects and advantages of this invention are further illustrated by the following examples, but the particular materials and amounts thereof recited in these examples, as well as other conditions and details, should not be construed to unduly limit this invention.

Example 1

2-Butyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine



Part A

A solution of 4-chloro-3-nitroquinoline (5.00 g, 24.0 mmol) in 120 mL of anhydrous CH₂Cl₂ was treated with triethylamine (6.7 mL, 48.2 mmol) and *tert*-butyl

carbazate (3.20 g, 24.2 mmol). After stirring under nitrogen for 2.5 hour (h), an additional portion of *tert*-butyl carbazate (3.2 g, 24.2 mmol) was added. After stirring overnight, the deep red solution was washed with H₂O (2X) and brine. The organic portion was dried over Na₂SO₄ and concentrated to give a red foam. The material was passed through a
5 SiO₂ column eluting with 2.5% methanol/CH₂Cl₂. The resulting red powder was treated with 5:1 hexanes/CH₂Cl₂ and filtered. The solid was washed several times with hexanes and was dried under vacuum to give *tert*-butyl *N'*-(3-nitroquinolin-4-yl)hydrazinecarboxylate (4.97 g) as an orange powder.

10 Part B

A suspension of *tert*-butyl *N'*-(3-nitroquinolin-4-yl)hydrazinecarboxylate (2.50 g, 8.22 mmol) in 150 mL of isopropanol was treated with 1.0 g of 10% palladium on carbon and the mixture was shaken under an atmosphere of hydrogen (3.8×10^5 Pa) for 2 h. The reaction mixture was then filtered through a pad of CELITE filter agent and rinsed with
15 isopropanol, and the filtrate was concentrated under reduced pressure to give *tert*-butyl *N'*-(3-aminoquinolin-4-yl)hydrazinecarboxylate (2.18 g) as a yellow solid.

Part C

A solution of *tert*-butyl *N'*-(3-aminoquinolin-4-yl)hydrazinecarboxylate (2.18 g, 7.96 mmol) in 80 mL of anhydrous CH₂Cl₂ was cooled to 0 °C and treated with
20 triethylamine (1.12 mL, 8.00 mmol) and valeryl chloride (0.95 mL, 8.00 mmol) under an atmosphere of nitrogen. After stirring for 3 h, the reaction mixture was concentrated under reduced pressure and the residue was treated with Et₂O and filtered. The filtrate was concentrated and the resulting black tar was dissolved in 80 mL of ethanol and treated
25 with 3 mL of triethylamine and the mixture was refluxed overnight. The reaction mixture was concentrated under reduced pressure. Chromatography (SiO₂, 1-5% methanol (MeOH)/CHCl₃) gave *tert*-butyl *N*-(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (1.41 g) as a mauve foam.

30 Part D

tert-Butyl *N*-(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (830 mg, 2.44 mmol) was dissolved in 20 mL of 1.5 M HCl in ethanol (EtOH) and the reaction mixture

was heated to reflux for 1.5 h. The reaction mixture was cooled and concentrated under reduced pressure to give a brown solid. The material was dissolved in 50 mL of hot isopropanol and the solution was allowed to cool overnight. The resulting crystals were isolated by filtration. A second crop was obtained from the filtrate by crystallization from isopropanol/Et₂O. The total yield of 2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine hydrochloride was 570 mg. mp > 250 °C. ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.68 (s, 1H), 9.35 (d, *J* = 8.3 Hz, 1H), 8.47 (d, *J* = 8.0 Hz, 1H), 8.03 (t, *J* = 7.1 Hz, 1H), 7.98 (t, *J* = 7.1 Hz, 1H), 6.85 (s, 2H), 3.13 (t, *J* = 7.6 Hz, 2H), 1.89, (m, 2H), 1.49 (m, 2H), 0.98 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 163.5, 139.4, 136.1, 134.0, 131.8, 130.4, 128.9, 122.6, 120.2, 115.6, 28.2, 25.7, 22.1, 13.3; Anal. Calcd for C₁₄H₁₆N₄·HCl: C, 60.76; H, 6.19; N, 20.24; Cl, 12.81. Found: C, 60.78; H, 6.19; N, 20.21; Cl, 12.78.

Part E

A solution of 2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine hydrochloride (520 mg, 2.17 mmol) in 10 mL of isopropanol was treated with 2 mL of acetone and 200 mg of DOWEX W50-X1 acid resin. The reaction mixture was heated to 55 °C overnight. The reaction mixture was treated with an additional 10 mL of isopropanol and 5 mL of acetone and heated to 70 °C for 2 h. The reaction mixture was filtered and the filtrate was treated with 0.5 mL of triethylamine and concentrated under reduced pressure. Chromatography (SiO₂, 3% MeOH/CHCl₃) gave *N*-(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylideneamine (421 mg) as a brown oil.

Part F

A solution of *N*-(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylideneamine (406 mg, 1.45 mmol) in 15 mL of MeOH was treated with NaBH₄ (500 mg, 13.2 mmol). After stirring for 2 days (d), the reaction was quenched with saturated NaHCO₃ solution and extracted into ethyl acetate (EtOAc). The organic portion was washed with H₂O and brine and dried over Na₂SO₄. Chromatography (SiO₂, EtOAc) gave *N*-(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (372 mg) as a mauve solid.


Part G

A solution of *N*-(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (334 mg, 1.18 mmol) in 10 mL of CH₂Cl₂ was treated with 3-chloroperoxybenzoic acid (MCPBA) (77% max., 334 mg, 1.45 mmol). After stirring for 3 h, the reaction was quenched with saturated NaHCO₃ solution and extracted into CH₂Cl₂. The organic portion was washed with saturated NaHCO₃ solution, H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated to give *N*-(2-butyl-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (338 mg) as a light brown solid.

Part H

A solution of *N*-(2-butyl-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (332 mg, 1.11 mmol) in 15 mL of 1,2-dichloroethane was placed in a pressure vessel and heated to 70 °C. The rapidly stirred solution was then treated with 3 mL of concentrated NH₄OH solution and *p*-toluenesulfonyl chloride (233 mg, 1.22 mmol), the reaction vessel was capped, and heating was continued for 2 h. The reaction mixture was then cooled to ambient temperature and treated with 50 mL of CH₂Cl₂. The reaction mixture was washed with H₂O, 1% Na₂CO₃ solution (3X), H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated. Chromatography (SiO₂, 5-10% MeOH/CHCl₃) gave 320 mg of a light brown solid. Crystallization from CH₂Cl₂/hexanes gave 2-butyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (230 mg) as colorless crystals. mp 157.1-158.7 °C. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.40 (m, 1H), 7.80 (m, 1H), 7.50 (m, 1H), 7.31 (m, 1H), 5.41 (s, 2H), 4.95 (s, 1H), 3.68 (m, 1H), 2.96 (t, *J* = 7.6 Hz, 2H), 1.93-1.82 (m, 2H), 1.48 (m, 2H), 1.16 d, *J* = 6.4 Hz, 6H), 1.00 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 155.1, 151.8, 144.7, 133.1, 127.3, 126.6, 124.7, 122.0, 120.4, 115.3, 52.1, 30.3, 26.8, 23.0, 20.8, 14.2; MS *m/z* 298 (*M* + *H*)⁺; Anal. Calcd for C₁₇H₂₃N₅: C, 68.66; H, 7.80; N, 23.55. Found: C, 68.30; H, 7.68; N, 23.33.

Example 2

*N*¹-Benzyl-2-butyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

Part A

5 A solution of 2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine hydrochloride (503 mg, 1.82 mmol) in 10 mL of isopropanol was treated with benzaldehyde (220 μ L, 2.17 mmol) and 200 mg of DOWEX W50-X1 acid resin. The reaction mixture was heated to reflux overnight. The reaction mixture was filtered, and the filtrate was treated with 0.5 mL of triethylamine and concentrated under reduced pressure. The resulting oil was dissolved in
10 75 mL of CH₂Cl₂ and washed with saturated NaHCO₃ solution, H₂O and brine. The organic was dried over Na₂SO₄, filtered and concentrated to give *N*-benzylidene(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (575 mg) as a light yellow solid.

Part B

15 A solution of *N*-benzylidene(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (575 mg, 1.75 mmol) in 40 mL of MeOH was treated with NaBH₄ (250 mg, 6.58 mmol). After stirring for 4 h, the reaction was quenched with saturated NaHCO₃ solution and extracted into CHCl₃. The organic portion was washed with H₂O and brine and dried over Na₂SO₄. Chromatography (SiO₂, 50-67% EtOAc/hexanes) gave *N*-benzyl(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (427 mg) as a yellow solid.
20

Part C

 A solution of *N*-benzyl(2-butyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (427 mg, 1.29 mmol) in 20 mL of CH₂Cl₂ was treated with MCPBA (77% max., 325 mg, 1.41
25 mmol). After stirring for 3 h, the reaction was quenched with saturated NaHCO₃ solution and extracted into CH₂Cl₂. The organic portion was washed with saturated NaHCO₃ solution, H₂O and brine. The organic was dried over Na₂SO₄, filtered and concentrated to

give *N*-benzyl(2-butyl-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (393 mg) as a light brown foam.


Part D

5 A solution of *N*-benzyl(2-butyl-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (393 mg, 1.14 mmol) in 20 mL of 1,2-dichloroethane was placed in a pressure vessel and heated to 70 °C. The rapidly stirred solution was then treated with 5 mL of concentrated NH₄OH solution and *p*-toluenesulfonyl chloride (239 mg, 1.25 mmol), the reaction vessel was capped, and heating was continued for 2 h. The reaction mixture was then cooled to
10 ambient temperature and treated with 50 mL of CH₂Cl₂. The reaction mixture was washed with H₂O, 1% Na₂CO₃ solution (3X), H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated. Chromatography (SiO₂, 5% MeOH/CHCl₃) followed by crystallization from propyl acetate/hexanes gave *N*¹-benzyl-2-butyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (237 mg) as light-yellow crystals. mp 159.3-160.5 °C. ¹H NMR
15 (300 MHz, DMSO-*d*₆) δ 8.31 (d, *J* = 8.2 Hz, 1H), 7.83 (d, *J* = 8.4 Hz, 1H), 7.54 (m, 1H), 7.42-7.31 (m, 6H), 5.44 (s, 2H), 5.26 (t, *J* = 5.6 Hz, 1H), 4.37 (d, *J* = 5.6 Hz, 2H), 2.71 (t, *J* = 8.4 Hz, 2H), 1.74 (m, 2H), 1.42 (m, 2H), 0.95 (t, *J* = 7.3 Hz, 3H); MS *m/z* 346 (*M* + H)⁺; Anal. Calcd for C₂₁H₂₃N₅: C, 73.02; H, 6.71; N, 20.27. Found: C, 72.75; H, 6.55; N, 20.46.

20

Example 3

*N*¹-Isopropyl-2-methyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine



Part A

25 A solution of *tert*-butyl *N*'-(3-aminoquinolin-4-yl)hydrazinecarboxylate (11.67 g, 42.5 mmol) in 400 mL of anhydrous toluene was treated with trimethyl orthoacetate (5.96 mL, 46.8 mmol) and pyridine hydrochloride (100 mg) under an atmosphere of N₂ and heated to reflux. After stirring for 3 h, the reaction mixture was concentrated under reduced pressure to give a red solid. Chromatography (SiO₂, 0-10% MeOH/EtOAc) gave

tert-butyl *N*-(2-methyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (10.7 g) as a yellow foam.

Part B

5 *tert*-Butyl *N*-(2-methyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (5.00 g, 16.8 mmol) was dissolved in 40 mL of 1.65 M HCl in EtOH, and the reaction mixture was heated to reflux for 2 h. The reaction mixture was cooled and concentrated under reduced pressure to give a brown solid. The brown solid was crystallized from ethanol/H₂O to give 3.13 g of 2-methyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine hydrochloride.

10

Part C

 A suspension of 2-methyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine hydrochloride (1.79 g, 7.62 mmol) in 30 mL of 2,2-dimethoxypropane was treated with 90 mg of *p*-toluenesulfonic acid. The reaction mixture was heated to 100 °C overnight. The reaction mixture was then treated with 10 mL of H₂O and 10 mL of MeOH, and heating was continued for 24 h. The reaction mixture was cooled and concentrated under reduced pressure. The resulting oil was dissolved in 50 mL of CHCl₃ and washed with 2% Na₂CO₃ solution, H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated to give *N*-isopropylidene(2-methyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.82 g) as a yellow solid.

20

Part D

 A solution of *N*-isopropylidene(2-methyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.82 g, 7.64 mmol) dissolved in 40 mL of MeOH was treated with NaBH₄ (1.16 g, 30.6 mmol). After stirring for 18 h, the reaction was quenched with saturated NH₄Cl solution and partitioned between CH₂Cl₂ and 2% Na₂CO₃ solution. The organic portion was washed with 2% Na₂CO₃ solution, H₂O and brine and dried over Na₂SO₄. The resulting organic portion was filtered and concentrated under reduced pressure to give *N*-isopropyl(2-methyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.84 g) as a yellow foam.

30

Part E


A solution of *N*-isopropyl(2-methyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.84 g, 7.66 mmol) dissolved in 50 mL of 1,2-dichloroethane was treated with MCPBA (77% max., 2.36 g, 9.58 mmol). After stirring for 3 h, the reaction mixture was treated with 2% Na₂CO₃ solution and extracted into CH₂Cl₂. The organic portion was washed with saturated 2% Na₂CO₃ solution, H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated to give *N*-isopropyl(2-methyl-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.95 g) as a light orange solid.

10 Part F

A solution of *N*-isopropyl(2-methyl-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.95 g, 7.61 mmol) in 75 mL of CH₂Cl₂ was treated with 35 mL of concentrated NH₄OH solution. To the rapidly stirred solution was added *p*-toluenesulfonyl chloride (1.52 g, 7.99 mmol). After stirring for 30 min, the reaction mixture was treated with CHCl₃ (25 mL) and H₂O (35 mL). The layers were separated and the organic portion was washed with 2% Na₂CO₃ solution (2X), H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated to give a light-yellow solid. Crystallization from propyl acetate gave *N*¹-isopropyl-2-methyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (747 mg) as off-white crystals. mp 227–229 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.19 (dd, *J* = 8.2, 1.1 Hz, 1 H), 7.79 (dd, *J* = 8.4, 0.7 Hz, 1 H), 7.53–7.45 (m, 1 H), 7.33–7.26 (m, 1 H), 5.42 (s, 2 H), 4.91 (d, *J* = 1.4 Hz, 1 H), 3.73–3.62 (m, 1 H), 2.64 (s, 3 H), 1.15 (d, *J* = 6.2 Hz, 6 H); ¹³C NMR (75 MHz, CDCl₃) δ 151.4, 151.3, 144.9, 133.3, 127.6, 127.3, 124.6, 122.4, 120.2, 115.4, 52.3, 20.9, 13.8; MS *m/z* 256 (*M* + H)⁺; Anal. Calcd for C₁₄H₁₇N₅: C, 65.86; H, 6.71; N, 27.43; Found: C, 65.59; H, 6.56; N, 27.09.

25

Example 4

*N*¹-Benzyl-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

Part A

5 A solution of *tert*-butyl *N*'-(3-aminoquinolin-4-yl)hydrazinecarboxylate (12.15 g, 44.3 mmol) in 200 mL of anhydrous CH₂Cl₂ was cooled to 0 °C and treated with triethylamine (7.72 mL, 55.4 mmol) and 2-ethoxyacetyl chloride (5.70 g, 46.5 mmol) under an atmosphere of N₂. After 3 h, an additional 1 mL of 2-ethoxyacetyl chloride was added. After stirring for 2 h, the reaction mixture was concentrated under reduced
10 pressure to give a brown solid. This was dissolved in 150 mL of EtOH and treated with 18.5 mL of triethylamine, and the mixture was refluxed overnight. The reaction mixture was concentrated under reduced pressure to give a dark-red oil. The red oil was dissolved in 200 mL of CH₂Cl₂ and washed with H₂O (2 X 75 mL) and brine (75 mL). The organic portion was dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give a
15 red solid. The solid was treated with a minimum amount of hot Et₂O and filtered to remove insoluble material. The filtrate was concentrated to give *tert*-butyl *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (14.3 g) as a tan solid.

Part B

20 *tert*-Butyl *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (14.3 g, 41.8 mmol) was dissolved in 150 mL of 2 M HCl in EtOH, and the reaction mixture was heated to reflux for 3 h. The reaction mixture was cooled and concentrated under reduced pressure to give a brown solid. The brown solid was dissolved in 100 mL of H₂O and treated with 100 mL of concentrated NH₄OH solution. The basic, aqueous solution was
25 then extracted with CH₂Cl₂ (4X). The combined organic layers were then washed with brine and dried over Na₂SO₄. The solution was filtered and concentrated under reduced pressure to give a brown foam. The foam was triturated with Et₂O (150 mL) and filtered.

The filtrate was concentrated to give 2-ethoxymethy-1*H*-imidazo[4,5-*c*]quinolin-1-amine (5.77 g) as a tan solid.

Part C

5 A solution of 2-ethoxymethy-1*H*-imidazo[4,5-*c*]quinolin-1-amine (1.50 g, 6.19 mmol) in 50 mL of isopropanol was treated with benzaldehyde (0.66 mL, 6.50 mmol) and 10 mg of *p*-toluenesulfonic acid. The reaction mixture was heated to 120 °C for 3 d. The reaction mixture was cooled, and a precipitate started to form. The reaction mixture was treated with Et₂O and then filtered to give *N*-benzylidene-(2-ethoxymethy-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.21 g) as a gray solid.

Part D

15 A solution of *N*-benzylidene-(2-ethoxymethy-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.00 g, 3.03 mmol) in 50 mL of MeOH was treated with NaBH₄ (458 mg, 12.1 mmol). After stirring for 1.5 h, the reaction mixture was concentrated, then treated with saturated NaHCO₃ solution, and extracted into CHCl₃. The organic portion was washed with H₂O and brine and dried over Na₂SO₄. The resulting solution was filtered and concentrated to give *N*-benzyl-(2-ethoxymethy-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.01 g) as a tan solid.

20

Part E

25 A solution of *N*-benzyl-(2-ethoxymethy-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (1.01 g, 3.04 mmol) in 50 mL of CH₂Cl₂ was treated with MCPBA (77% max., 1.02 g, 4.56 mmol). After stirring for 3 h, the reaction mixture was quenched with 2% Na₂CO₃ solution and extracted into CH₂Cl₂. The organic portion was washed with H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated to give *N*-benzyl-(2-ethoxymethy-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (0.99 g) as a light-yellow solid.

30


Part F

 A solution of *N*-benzyl-(2-ethoxymethy-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (0.99 g, 2.84 mmol) in 50 mL of CH₂Cl₂ was treated with 25 mL of concentrated

NH₄OH solution. To the rapidly stirred solution was added *p*-toluenesulfonyl chloride (569 mg, 2.98 mmol). After stirring for 30 min, the reaction was treated with CH₂Cl₂ (50 mL) and H₂O (25 mL). The layers were separated and the organic portion was washed 2% Na₂CO₃ solution, H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated to give a tan solid. Chromatography (SiO₂, 2% MeOH/CHCl₃ containing 0.5% concentrated NH₄OH) followed by crystallization from propyl acetate gave *N*¹-benzyl-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (148 mg) as white needles. mp 152–155 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.61 (dd, *J* = 8.2, 1.2 Hz, 1 H), 7.85–7.77 (m, 1 H), 7.59–7.52 (m, 1 H), 7.42–7.34 (m, 4 H), 7.33–7.24 (m, 2 H), 6.02 (t, *J* = 6.6 Hz, 1 H), 5.39 (s, 2 H), 4.43 (s, 2 H), 4.40 (d, *J* = 6.7 Hz, 2 H), 3.55 (q, *J* = 7.0 Hz, 2 H), 1.22 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 151.1, 147.9, 144.9, 135.7, 129.2, 129.1, 128.6, 127.8, 126.7, 122.4, 120.7, 66.7, 65.3, 56.7, 15.0; MS *m/z* 348 (*M* + H)⁺; Anal. Calcd for C₂₀H₂₁N₅O·0.36H₂O: C, 68.90; H, 6.11; N, 20.09; Found: C, 68.50; H, 6.07; N, 20.11.

Example 5'

2-Ethoxymethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine



Part A

A solution of 2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (2.50 g, 10.3 mmol) in 250 mL of 1,2-dichloroethane was treated with acetone (0.83 mL, 11.3 mmol), acetic acid (0.65 mL, 11.3 mmol) and sodium triacetoxymethylborohydride (2.39 g, 11.3 mL). After stirring overnight, additional acetone (5 mL), acetic acid (0.65 mL, 11.3 mmol) and sodium triacetoxymethylborohydride (2.39 g, 11.3 mL) were added. After 2 d, the reaction was carefully quenched by addition of saturated NaHCO₃ solution. The layers were separated and the aqueous portion was extracted with additional CH₂Cl₂. The combined organic layers were washed with H₂O and brine, dried over Na₂SO₄, and concentrated under

reduced pressure to give a brown oil. Some isopropylidene intermediate was still present, so the material was dissolved in 50 mL of MeOH and treated with NaBH₄ (1.0 g). After 2 h, the reaction was quenched by the addition of H₂O and the reaction mixture was concentrated under reduced pressure. The residue was partitioned between saturated NaHCO₃ solution and CH₂Cl₂. The layers were separated and the organic portion was washed with saturated NaHCO₃, H₂O and brine. The organic portion was dried over Na₂SO₄, filtered, and concentrated under reduced pressure. Chromatography (SiO₂, 4% MeOH/CHCl₃) gave *N*-(2-ethoxymethy-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (0.98 g) as a brown oil.

Part B

A solution of *N*-(2-ethoxymethy-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (0.98 g, 3.45 mmol) in 35 mL of CH₂Cl₂ was treated with MCPBA (77% max., 1.10 g, 4.48 mmol). After stirring for 3 h, the reaction was quenched with 2% Na₂CO₃ solution and extracted into CH₂Cl₂. The organic portion was washed with H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated to give *N*-(2-ethoxymethy-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (0.93 g) as a light-orange solid.

Part C


A solution of *N*-(2-ethoxymethy-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (0.93 g, 3.10 mmol) in 25 mL of CH₂Cl₂ was treated with 15 mL of concentrated NH₄OH solution. To the rapidly stirred solution was added *p*-toluenesulfonyl chloride (620 mg, 3.25 mmol). After stirring for 30 min, the reaction was treated with CH₂Cl₂ (20 mL) and H₂O (15 mL). The layers were separated and the organic portion was washed with 2% Na₂CO₃ solution, H₂O and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated to give a tan solid. Chromatography (SiO₂, 5% MeOH/CHCl₃) gave 2-ethoxymethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (368 mg) as a tan solid. mp 162–164 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.60 (dd, *J* = 8.2, 1.1 Hz, 1 H), 7.77 (dd, *J* = 8.4, 0.7 Hz, 1 H), 7.54-7.47 (m, 1 H), 7.33-7.24 (m, 1 H), 5.55 (d, *J* = 3.2 Hz, 1 H), 5.41 (s, 2 H), 4.89 (s, 2 H), 3.73-3.60 (m, 3 H), 1.26 (t, *J* = 7.0 Hz, 3H); 1.15 (d, *J* = 6.2 Hz, 6 H); ¹³C NMR (75 MHz, CDCl₃) δ 151.1, 148.7, 145.0,

127.7, 126.6, 123.9, 121.9, 121.3, 115.4, 66.8, 65.7, 52.5, 20.6, 15.1; MS m/z 300 ($M + H$)⁺; Anal. Calcd for $C_{16}H_{21}N_5O \cdot 0.48 H_2O$: C, 62.39; H, 7.19; N, 22.74; Found: C, 62.38; H, 6.90; N, 22.79.

5

Example 6

*N*¹-Cyclohexyl-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine



Part A

2-(Ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-amine (0.900 g, 3.71 mmol) was placed in a 50 mL round bottom flask, dissolved in 1,2-dichloromethane, and placed under N₂. Cyclohexanone (1.19 mL, 11.5 mmol), acetic acid (0.45 mL, 7.79 mmol) and sodium triacetoxyborohydride (1.65 g, 7.79 mmol) were added and the reaction was stirred under N₂ at room temperature for 5 days. The reaction was quenched by slow addition of saturated NaHCO₃ solution (25 mL) and dichloromethane (25 mL). The mixture was transferred to a separatory funnel and the phases separated. The aqueous portion was extracted with dichloromethane (25 mL). The combined organic portions were washed sequentially with water (25 mL) and brine (25 mL), dried (Na₂SO₄), filtered and then concentrated to yield a thick brown oil. Analysis by liquid chromatography/mass spectroscopy (LC/MS) of the crude product showed it to be a mixture of the hydrazone and hydrazine. The oil was dissolved in methanol (25 mL), chilled in an ice water bath and then treated with sodium borohydride (1.25 g). The reaction was quenched with water (25 mL) and the mixture concentrated. The residue was partitioned between dichloromethane 50 mL) and water (15 mL), transferred to a separatory funnel, and the phases were separated. The organic portion was washed sequentially with saturated NaHCO₃ solution (20 mL), water (20 mL) and brine (20 mL), dried (Na₂SO₄), filtered and then concentrated to yield a thick brown oil. The material was purified by column chromatography (35 g SiO₂, 97:3 chloroform:methanol) to yield 0.51 g of *N*-cyclohexyl-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-amine as a light brown oil / solid.

Part B

N-Cyclohexyl-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-amine (0.51 g, 1.57 mmol) was placed in a 200 mL round bottom flask, purged with N₂ and dissolved in dichloromethane (25 mL). MCPBA (0.484 g, 1.96 mmol, 77% max) was added over a 5 min period. The reaction was stirred at room temperature under N₂. After 2 h, analysis by thin layer chromatography (TLC) (SiO₂, 95:5 chloroform:methanol) showed complete conversion. The solution was diluted with dichloromethane (15 mL) and 2% sodium carbonate solution (15 mL). The mixture was transferred to a separatory funnel, and the phases were separated. The organic portion was washed sequentially with 2% sodium carbonate solution (15 mL), water (15 mL) and brine (15 mL), dried (Na₂SO₄), filtered and then concentrated to yield 0.431 g of *N*-cyclohexyl-2-(ethoxymethyl)-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-amine as a tan foam.


15 Part C

N-Cyclohexyl-2-(ethoxymethyl)-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-amine (0.425 g, 1.25 mmol) was placed in a 100 mL round bottom flask and dissolved in dichloromethane (20 mL). Ammonium hydroxide solution (10 mL) was added and the mixture was stirred vigorously. The stirred mixture was chilled in an ice water bath. *Para*-toluenesulfonyl chloride (0.250 g, 1.31 mmol) was added over 5 min. After 30 min of stirring at 0 °C TLC (SiO₂, 95:5 chloroform:methanol) showed complete conversion. The mixture was warmed to room temperature and then diluted with dichloromethane (25 mL) and water (10 mL). The mixture was transferred to a separatory funnel and the phases separated. The organic portion was washed sequentially with 2% sodium carbonate solution (15 mL), water (15 mL) and brine (15 mL), dried over Na₂SO₄, filtered and then concentrated to yield an orange/tan foamy solid. The material was purified by column chromatography (40 g SiO₂, 95:5 chloroform:methanol) to yield the product as an off white solid. The off-white solid was dissolved in 3 mL of a 9:1 chloroform:methanol mixture. A small spatula tip full of activated carbon (DARCO G 60-100 mesh) was added and the mixture was stirred at room temperature for 3 h. The mixture was filtered through a short column of SiO₂ (5 g) eluting with 9:1 chloroform:methanol. The filtrate was concentrated to yield a glassy solid. The glassy solid was triturated in 15 mL diethyl ether

for 2 h to provide a white solid. The solid was collected by vacuum filtration and rinsed with diethyl ether. The solid was dried in a vacuum oven (70 °C) to yield 0.062 g of *N*¹-cyclohexyl-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine. mp 143–145 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.61 (dd, *J* = 8.1, 1.1 Hz, 1 H), 7.58 (dd, *J* = 8.3, 0.9 Hz, 1 H), 7.46–7.38 (m, 1 H), 7.28–7.21 (m, 1 H), 6.99 (d, *J* = 1.9 Hz, 1 H), 6.69 (s, 2 H), 4.77 (s, 2 H), 3.63 (q, *J* = 7.0 Hz, 2 H), 3.32–3.23 (m, 1 H), 1.71–1.52 (m, 5 H), 1.30–1.05 (m, 8 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ ; MS *m/z* 152.1, 150.3, 145.0, 133.4, 127.4, 125.8, 123.9, 121.6, 121.1, 115.0, 65.8, 63.1, 59.8, 30.9, 25.8, 24.3, 15.4; MS *m/z* 340 (M + H)⁺; Anal. Calcd for C₁₉H₂₅N₅O: C, 67.23; H, 7.42; N, 20.63; Found: C, 67.32; H, 7.37; N, 20.55.

Example 7

*N*¹,*N*¹-Dimethyl-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine



Part A

A solution of 4-chloro-3-nitroquinoline (5.00 g, 24.0 mmol) in 100 mL CH₂Cl₂ was cooled to 0 °C and treated with triethylamine (8.40 mL, 60.0 mmol) and *N,N*-dimethylhydrazine (5.65 mL, 74.4 mmol) under an atmosphere of nitrogen. After 18 h, the mixture was diluted with 2% Na₂CO₃ solution and CHCl₃ and separated. The organic portion was washed with water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 4-(2,2-dimethylhydrazino)-3-nitroquinoline (5.33 g) as a yellow/orange crystalline solid.

Part B

A suspension of 4-(2,2-dimethylhydrazino)-3-nitroquinoline (5.33 g, 23.0 mmol) in 125 mL of acetonitrile was treated with 5% platinum on carbon (0.45 g, 0.11 mmol) and the mixture was shaken under an atmosphere of hydrogen (3.8 x 10⁵ Pa). After 5 h, the reaction mixture was filtered through a pad of CELITE filter agent and rinsed with 80:20 acetonitrile:MeOH. The filtrate was concentrated under reduced pressure. The resulting

oil was dissolved in CH₂Cl₂, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 4-(2,2-dimethylhydrazino)quinolin-3-amine (4.64 g) as a red foam.

Part C


5 A solution of 4-(2,2-dimethylhydrazino)quinolin-3-amine (4.64 g, 23.0 mmol) in 75 mL of CH₂Cl₂ was cooled to 0 °C under an atmosphere of nitrogen. The reaction mixture was treated with triethylamine (6.72 mL, 48.2 mmol) followed by dropwise addition of ethoxyacetyl chloride (2.95 g, 24.1 mmol). After 1.5 h, the reaction mixture was concentrated under reduced pressure. The resulting oil was dissolved in 75 mL of ethanol, treated with triethylamine (9.60 mL, 68.9 mmol) and heated to reflux. After 5 d, the reaction mixture was concentrated under reduced pressure. The resulting oil was dissolved in CH₂Cl₂, washed with 2% Na₂CO₃ solution, water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a brown oil. Chromatography (SiO₂, 5-10% MeOH/CHCl₃) gave *N,N*-dimethyl-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-amine (0.89 g) as a brown oil.

Part D

A solution of *N,N*-dimethyl-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-amine (0.89 g, 3.3 mmol) in 25 mL of CH₂Cl₂ was treated with MCPBA (1.01 g, 4.10 mmol, 77% max). After 1.5 h, the reaction mixture was treated with 7 mL of concentrated NH₄OH solution and *p*-toluenesulfonyl chloride (0.69 g, 3.6 mmol). After 30 min, the reaction was diluted with CH₂Cl₂ and water and the phases were separated. The organic portion was washed with 2% Na₂CO₃ solution (2X), water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield an orange solid.

25 Recrystallization twice from acetonitrile gave *N,N'*-dimethyl-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.208 g) as gold, needle-like crystals. mp 213–215 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.57 (dd, *J* = 8.3, 1.4 Hz, 1 H), 7.79 (dd, *J* = 8.4, 0.7 Hz, 1 H), 7.56–7.48 (m, 1 H), 7.38–7.29 (m, 1 H), 5.45 (s, 2 H), 4.48 (s, 2 H), 3.69 (q, *J* = 7.0 Hz, 2 H), 3.20 (s, 6 H), 1.29 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 151.2, 149.3, 145.1, 133.5, 127.7, 126.7, 123.8, 122.1, 115.3, 66.4, 65.6, 45.3, 15.1; MS (APCI) *m/z* 286 (M + H)⁺; Anal. Calcd for C₁₅H₁₉N₅O: C, 63.14; H, 6.71; N, 24.54; Found: C, 63.02; H, 6.91; N, 24.57.

Example 8

2-Ethoxymethyl-*N*¹-(furan-2-ylmethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

5 Part A

A solution of 2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (1.50 g, 6.19 mmol) in 20 mL of isopropanol was treated with 2-furaldehyde (1.08 mL, 13.0 mmol) and 2 drops of concentrated HCl and heated to reflux under an atmosphere of nitrogen. After 48 h, the reaction was concentrated under reduced pressure to yield a brown oil. The oil was dissolved in 30 mL of CHCl₃ and washed with 5% Na₂CO₃ solution, water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(furan-2-ylmethylene)amine (1.86 g) as a light brown solid.

15 Part B

A solution of *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(furan-2-ylmethylene)amine (1.86 g, 5.81 mmol) in 20 mL of methanol was treated with NaBH₄ (0.659 g, 17.4 mmol) and stirred under an atmosphere of nitrogen. After 18 h the reaction was quenched by addition of 20 mL of water. The reaction mixture was concentrated under reduced pressure and dissolved in CHCl₃. The organic portion was washed with 2% Na₂CO₃ solution, water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(furan-2-ylmethyl)amine (1.70 g) as a thick orange syrup.


25 Part C

A solution of *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(furan-2-ylmethyl)amine (1.70 g, 5.27 mmol) in 45 mL of CH₂Cl₂ was treated with MCPBA (1.48 g, 6.59 mmol, 77% max). After 1.5 h the reaction mixture was treated with 15 mL of

concentrated NH_4OH solution and *p*-toluenesulfonyl chloride (1.06 g, 5.54 mmol). After 45 min the reaction mixture was diluted with water and CHCl_3 and separated. The organic portion was washed with 3% Na_2CO_3 solution, water and brine, dried over Na_2SO_4 , and concentrated under reduced pressure to yield a yellow foam. Chromatography (SiO_2 , 95:5 CHCl_3 :MeOH) gave an off white foam. The foam was triturated with diethyl ether and filtered to give 2-ethoxymethyl-*N*¹-(furan-2-ylmethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.03 g) as an off white powder. mp dec. > 200 °C; ^1H NMR (300 MHz, CDCl_3) δ 8.57 (dd, J = 8.1, 1.1 Hz, 1 H), 7.80 (dd, J = 8.4, 0.8 Hz, 1 H), 7.57-7.51 (m, 1 H), 7.45 (d, J = 1.8 Hz, 1 H), 7.39-7.33 (m, 1 H), 6.34-6.32 (m, 1 H), 6.24 (t, J = 5.3 Hz, 1 H), 6.07 (d, J = 3.1 Hz, 1 H), 5.43 (s, 2 H), 4.40-4.38 (m, 4 H), 3.57 (q, J = 7.0 Hz, 2 H), 1.25 (t, J = 7.0 Hz, 3 H); ^{13}C NMR (75 MHz, CDCl_3) δ 151.1, 149.5, 147.8, 144.8, 143.0, 132.6, 127.8, 126.6, 124.1, 122.5, 120.7, 115.1, 111.1, 110.1, 66.8, 64.9, 48.5, 15.0; MS (APCI) m/z 338 ($\text{M} + \text{H}$)⁺; Anal. Calcd for $\text{C}_{18}\text{H}_{19}\text{N}_5\text{O}_2$: C, 64.08; H, 5.68; N, 20.76; Found: C, 63.89; H, 5.75; N, 20.48.

Example 9

2-Ethoxymethyl-*N*¹-(1-ethylpropyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine



Part A

A solution of 2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (1.50 g, 6.19 mmol) in 20 mL of toluene and 5 mL of isopropanol was treated with 3-pentanone (5.00 mL, 47.2 mmol) and pyridinium *p*-toluenesulfonate (0.015 g, 0.062 mmol) and the reaction mixture was heated to reflux under an atmosphere of nitrogen. After 7 d, the reaction mixture was concentrated under reduced pressure, dissolved in CHCl_3 , washed with water (2X) and brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield a light brown oil. Chromatography (SiO_2 , 95:5 CHCl_3 :MeOH) gave *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(1-ethylpropylidene)amine (1.78 g) as a yellow/green syrup.


Part B

A solution of *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(1-ethylpropylidene)amine (1.78 g, 5.73 mmol) in 20 mL of methanol was treated with NaBH₄ (0.867 g, 22.9 mmol) and CeCl₃·7H₂O (15 mg, catalytic) and stirred under an atmosphere of nitrogen. After 24 h, the reaction was concentrated under reduced pressure, dissolved CHCl₃, washed with water (2X) and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a yellow/green syrup. Chromatography (SiO₂, 93:7 CHCl₃:MeOH) gave *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(1-ethylpropyl)amine (1.01 g) as a yellow/green oil.

Part C

A solution of *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(1-ethylpropyl)amine (1.01 g, 3.23 mmol) in 30 mL of CH₂Cl₂ was treated with MCPBA (1.04 g, 4.20 mmol, 77% max). After 1.5 h the reaction mixture was treated with 15 mL of concentrated NH₄OH solution and *p*-toluenesulfonyl chloride (0.65 g, 3.39 mmol). After 30 min, the reaction mixture was diluted with CH₂Cl₂ and water and the phases were separated. The organic portion was washed with 2 % Na₂CO₃ solution and water. The combined aqueous washes were back extracted with CHCl₃ (2X). The combined organic portions were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a light yellow foam. Chromatography (SiO₂, 97:3 CHCl₃:MeOH) gave a white foam. The foam was triturated with CH₂Cl₂/hexanes and filtered to give 2-ethoxymethyl-*N*¹-(1-ethylpropyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.652 g) as a white solid. mp 125–128 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.66 (dd, *J* = 8.3, 1.1 Hz, 1 H), 7.77 (dd, *J* = 7.6, 0.8 Hz, 1 H), 7.55-7.48 (m, 1 H), 7.33-7.26 (m, 1 H), 5.66, (d, *J* = 3.0 Hz, 1 H), 5.41 (s, 2 H), 4.87 (s, 2 H), 3.64 (q, *J* = 7.0 Hz, 2 H), 3.32-3.23 (m, 1 H), 1.70-1.56 (m, 2 H), 1.55-1.41 (m, 2 H), 1.27 (t, *J* = 7.1 Hz, 3 H), 0.94 (t, *J* = 7.5 Hz, 6 H); ¹³C NMR (75 MHz, CDCl₃) δ 151.5, 149.1, 145.4, 135.0, 132.4, 128.1, 126.9, 124.1, 122.2, 122.0, 115.9, 67.2, 66.2, 64.0, 24.5, 15.5, 10.2; MS (APCI) *m/z* 328 (M + H)⁺; Anal. Calcd for C₁₈H₂₅N₅O: C, 66.03; H, 7.70; N, 21.39; Found: C, 65.64; H, 7.89; N, 21.02.

Example 10

2-Ethoxymethyl-*N*¹-isobutyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

5 Part A

A solution of 2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (0.940 g, 3.88 mmol) in 20 mL of toluene and 5 mL of isopropanol was treated with isobutyraldehyde (0.800 mL, 8.81 mmol) and pyridinium *p*-toluenesulfonate (0.098 g, 0.39 mmol) and the reaction mixture was heated to reflux under an atmosphere of nitrogen. After 48 h, the reaction mixture was concentrated under reduced pressure and dissolved in CHCl₃. The organic portion was washed with water (2X) and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a light brown oil which solidified under vacuum to yield *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isobutylideneamine (1.15 g) as a tan solid.

15

Part B

A solution of *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isobutylideneamine (1.15 g, 3.88 mmol) in 15 mL of methanol was treated with NaBH₄ (0.44 g, 11.6 mmol) and stirred under an atmosphere of nitrogen. After 18 h, the reaction was concentrated under reduced pressure. The residue was partitioned between CHCl₃ and water, and the phases were separated. The organic portion was washed with water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield an orange oil. Chromatography (SiO₂, 97:3 CHCl₃:MeOH), gave *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isobutylamine (0.69 g) as clear, colorless crystals.

25


Part C

A solution of *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isobutylamine (1.16 g, 3.89 mmol) in 30 mL of CH₂Cl₂ was treated with MCPBA (1.25 g, 5.05 mmol,

77% max). After 1.5 h, the reaction mixture was treated with 15 mL of concentrated NH_4OH solution and *p*-toluenesulfonyl chloride (0.78 g, 4.08 mmol). After 30 min the reaction mixture was diluted with CH_2Cl_2 and water, and the phases were separated. The organic portion was washed with 2% Na_2CO_3 solution and water. The combined aqueous washes were back extracted with CHCl_3 (2X). The combined organic portions were washed with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield a brown foam. Chromatography (SiO_2 , 97:3 CHCl_3 :MeOH) yielded 2-ethoxymethyl-*N*¹-isobutyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.049 g) as an off white solid. mp 137–140 °C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$, 350 K) δ 8.47 (dd, $J = 8.1$, 0.9 Hz, 1 H), 7.60 (d, $J = 8.3$ Hz, 1 H), 7.45–7.36 (m, 1 H), 7.28–7.19 (m, 1 H), 6.67, (t, $J = 6.2$ Hz, 1 H), 6.22 (s, 2 H), 4.76 (s, 2 H), 3.64 (q, $J = 7.0$ Hz, 2 H), 3.02 (t, $J = 6.4$ Hz, 2 H), 1.97 (s, $J = 6.7$ Hz, 1 H), 1.19 (t, $J = 7.0$ Hz, 3 H), 1.05 (d $J = 6.7$ Hz, 6 H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 151.9, 148.9, 144.8, 131.9, 126.9, 125.7, 123.8, 120.8, 114.2, 65.4, 62.8, 59.6, 26.7, 20.5, 14.9; MS (APCI) m/z 314 ($\text{M} + \text{H}$)⁺; Anal. Calcd for $\text{C}_{17}\text{H}_{23}\text{N}_5\text{O}$: C, 65.15; H, 7.40; N, 22.35; Found: C, 64.88; H, 7.39; N, 22.38.

Example 11

2-Ethoxymethyl-*N*¹-isopropyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine




20 Part A

A solution of 2-ethoxymethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.700 g, 2.34 mmol) in 25 mL of trifluoroacetic acid was treated with platinum(IV) oxide (0.27 g, 1.2 mmol) and the mixture was shaken under an atmosphere of hydrogen (3.8×10^5 Pa). After 15 h, the reaction mixture was filtered through a pad of CELITE filter agent, rinsed with 9:1:0.5 CHCl_3 :MeOH:trifluoroacetic acid (TFA) and concentrated under reduced pressure to yield a creamy white solid. The solid was triturated with concentrated NH_4OH solution for 2 h and then extracted with CHCl_3 (3X). The organic portion was washed with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield a white foam. The foam was triturated with diethyl ether,

filtered and dried under reduced pressure to yield 2-ethoxymethyl-*N*¹-isopropyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.376 g) as a fine white solid. mp 144–146 °C; ¹H NMR (300 MHz, CDCl₃) δ 5.08 (d, *J* = 2.7 Hz, 1 H), 4.92 (s, 2 H), 4.78 (s, 2 H), 3.61 (q, *J* = 7.0 Hz, 2 H), 3.53–3.43 (m, 1 H), 3.07–3.03 (m, 2 H), 2.85–2.81 (m, 2 H), 1.92–1.79 (m, 4 H), 1.25 (t, *J* = 7.0 Hz, 3 H), 1.08 (d, *J* = 6.3 Hz, 6 H); ¹³C NMR (75 MHz, CDCl₃) δ 149.4, 148.9, 148.1, 138.8, 122.9, 107.4, 66.6, 65.4, 53.0, 32.5, 23.7, 23.2, 22.8, 20.5, 15.1; MS (APCI) *m/z* 304 (*M* + *H*)⁺; Anal. Calcd for C₁₆H₂₅N₅O: C, 63.34; H, 8.31; N, 23.08; Found: C, 63.32; H, 8.31; N, 22.97.

10

Example 12

2-Ethoxymethyl-*N*¹-(3-methylbutyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

Part A

15

A solution of 2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (1.00 g, 4.13 mmol) in 20 mL of toluene and 5 mL of isopropanol was treated with isovaleraldehyde (0.94 mL, 8.76 mmol) and pyridinium *p*-toluenesulfonate (0.052 g, 0.21 mmol) and the reaction mixture was heated to reflux under an atmosphere of nitrogen. After 15 h, the reaction mixture was concentrated under reduced pressure to yield a brown oil. The oil was dissolved in CHCl₃ and washed with water (2X) and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(3-methylbutylidene)amine (1.28 g) as a dark orange oil.

20

Part B

25

A solution of *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(3-methylbutylidene)amine (1.28 g, 4.13 mmol) in 25 mL of methanol was treated with NaBH₄ (0.47 g, 12.39 mmol). After 1 h, the reaction was quenched with saturated NH₄Cl solution and the mixture was concentrated under reduced pressure. The residue was

partitioned between CHCl_3 and saturated NaHCO_3 solution and the phases were separated. The organic portion was washed with water and brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(3-methylbutyl)amine (1.24 g) as a dark orange oil.


5

Part C

A solution of *N*-(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)(3-methylbutyl)amine (1.24 g, 3.97 mmol) in 45 mL of CH_2Cl_2 was treated with MCPBA (1.87 g, 7.04 mmol, 77% max). After 1.5 h, the reaction mixture was treated with 15 mL of concentrated NH_4OH solution and *p*-toluenesulfonyl chloride (0.795 g, 4.17 mmol). After 30 min, the reaction mixture was diluted with CHCl_3 and water and the phases were separated. The organic portion was washed with 5% Na_2CO_3 solution, water and brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield a sticky orange foam. Chromatography (SiO_2 , 97:3 CHCl_3 :MeOH) gave an off white foam. The foam was triturated with diethyl ether and hexanes and filtered to give 2-ethoxymethyl-*N*¹-(3-methylbutyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.435 g) as a cream colored solid. mp 129–132 °C; ^1H NMR (300 MHz, CDCl_3) δ 8.48 (dd, $J = 8.1, 1.1$ Hz, 1 H), 7.78 (d, $J = 8.3$ Hz, 1 H), 7.56–7.50 (m, 1 H), 7.36–7.30 (m, 1 H), 5.59 (t, $J = 6.7$ Hz, 1 H), 5.42 (s, 2 H), 4.87 (s, 2 H), 3.64 (q, $J = 7.0$ Hz, 2 H), 3.29 (q, $J = 7.0$ Hz, 2 H), 1.76 (s, $J = 6.7$ Hz, 1 H), 1.60 (q, $J = 6.9$ Hz, 2 H), 1.27 (t, $J = 7.0$ Hz, 3 H), 0.97 (d, $J = 6.6$ Hz, 6 H); ^{13}C NMR (75 MHz, CDCl_3) δ 151.2, 147.8, 144.9, 133.1, 127.8, 126.6, 124.0, 122.3, 120.7, 115.2, 66.8, 65.3, 51.1, 36.7, 26.0, 22.6, 15.1; MS (APCI) m/z 328 ($\text{M} + \text{H}$)⁺; Anal. Calcd for $\text{C}_{18}\text{H}_{25}\text{N}_5\text{O} \cdot 0.06\text{H}_2\text{O}$: C, 65.81; H, 7.71; N, 21.32; Found: C, 65.42; H, 7.75; N, 21.11. Karl Fischer analysis 0.32% water.

25

Example 13

2-Ethoxymethyl-1-(morpholin-4-yl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine

Part A

A solution of 4-chloro-3-nitroquinoline (5.00 g, 24.0 mmol) in 100 mL of CH₂Cl₂ was treated with triethylamine (6.37 mL, 48.0 mmol) and 4-aminomorpholine (3.47 mL, 36.0 mL) under an atmosphere of nitrogen. After 15 h, the reaction mixture was diluted with 5% Na₂CO₃ solution and CHCl₃, and the phases were separated. The organic portion was washed with another portion of 5% Na₂CO₃ solution, water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a bright yellow solid. Recrystallization from acetonitrile gave *N*-(morpholin-4-yl)(3-nitroquinolin-4-yl)amine (4.54 g) as bright yellow needle-like crystals.

Part B

A solution of *N*-(morpholin-4-yl)(3-nitroquinolin-4-yl)amine (4.54 g, 16.6 mmol) in 150 mL of toluene was treated with 5% platinum on carbon (0.65 g, 0.17 mmol) and the mixture was shaken under an atmosphere of hydrogen (3.8 x 10⁵ Pa). After 15 h, the reaction mixture was filtered through a pad of CELITE filter agent and rinsed with 4:1 toluene:MeOH. The filtrate was concentrated under reduced pressure to yield *N*⁴-(morpholin-4-yl)quinoline-3,4-diamine (4.06 g) as a red foam.

20 Part C

A solution of *N*⁴-(morpholin-4-yl)quinoline-3,4-diamine (4.06 g, 16.6 mmol) in 50 mL of CH₂Cl₂ was treated with triethylamine (4.40 mL, 33.2 mmol) and cooled to 0 °C. The solution was treated dropwise with ethoxyacetyl chloride (2.40 g, 17.4 mmol) and stirred under an atmosphere of nitrogen. The reaction mixture was allowed to slowly come to room temperature. After 2 d, the reaction mixture was concentrated under reduced pressure to yield a red semi-solid. The material was dissolved in CHCl₃ and washed with water, 5% Na₂CO₃ solution and brine, dried over Na₂SO₄, filtered and dried to yield 2-ethoxy-*N*-{4-[(morpholin-4-yl)amino]quinolin-3-yl}acetamide (5.35 g) as a red/orange foam.

30

Part D


A suspension of 2-ethoxy-*N*-{4-[(morpholin-4-yl)amino]quinolin-3-yl}acetamide (5.35 g, 16.2 mmol) in 65 mL of toluene was treated with pyridine hydrochloride (0.94 g, 0.081 mmol). The reaction flask was equipped with a Dean-Stark trap and the reaction mixture was heated to reflux under an atmosphere of nitrogen. After 2.5 d, the reaction mixture was concentrated under reduced pressure to yield a brown oil. The oil was dissolved in CHCl₃ and was washed with 5% Na₂CO₃ solution, water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a brown foam. Chromatography (SiO₂, 95:5 CHCl₃:MeOH) gave 2-ethoxymethyl-1-(morpholin-4-yl)-1*H*-imidazo[4,5-*c*]quinoline (1.61 g) as a light brown solid.

Part E

A solution of 2-ethoxymethyl-1-(morpholin-4-yl)-1*H*-imidazo[4,5-*c*]quinoline (1.61 g, 5.51 mmol) in 40 mL of CH₂Cl₂ was treated with MCPBA (1.78 g, 6.70 mmol, 77% max). After 30 min, the reaction mixture was treated with 20 mL of concentrated NH₄OH solution and *p*-toluenesulfonyl chloride (1.03 g, 5.41 mmol). After 15 min, the reaction mixture was diluted with CH₂Cl₂ and water and the phases were separated. The organic portion was washed with 5% Na₂CO₃ solution, water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan foam. Chromatography (SiO₂, 97:3 CHCl₃:MeOH) gave a light yellow foam. The foam was triturated with diethyl ether and filtered to give 2-ethoxymethyl-1-(morpholin-4-yl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (0.794 g) as a light cream colored solid. mp 223–224 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.77 (d, *J* = 8.1 Hz, 1 H), 7.79 (d, *J* = 8.4 Hz, 1 H), 7.54 (t, *J* = 8.2 Hz, 1 H), 7.34 (t, *J* = 8.1 Hz, 1 H), 5.48 (s, 2 H), 4.85 (s, 2 H), 4.06–4.03 (m, 4 H), 3.74–3.66 (m, 4 H), 3.42–3.38 (m, 2 H), 1.29 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 151.2, 149.0, 145.3, 133.5, 127.9, 126.9, 123.7, 122.2, 121.3, 115.3, 67.5, 66.5, 65.9, 53.5, 15.1; MS (APCI) *m/z* 328 (M + H)⁺; Anal. Calcd for C₁₇H₂₁N₅O₂: C, 62.37; H, 6.47; N, 21.39; Found: C, 62.14; H, 6.19; N, 21.34.

Example 14

N-{3-[(4-Amino-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}
methanesulfonamide



5 Part A

A solution of 1-amino-3,3-diethoxypropane (5.00 mL, 30.9 mmol) in 5 mL of tetrahydrofuran (THF) was treated with triethylamine (4.51 mL, 34.0 mmol) under an atmosphere of nitrogen and cooled to 0 °C. The reaction mixture was then treated dropwise with a solution of di-*tert*-butyl dicarbonate (7.42 g, 34.0 mmol) in 25 mL of THF. The reaction mixture was stirred for 2 h at 0 °C and then allowed to come to room temperature. After 15 h, the reaction mixture was concentrated under reduced pressure, dissolved in ethyl acetate, washed with water (2X) and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield *tert*-butyl (3,3-diethoxypropyl)carbamate (8.40 g) as a clear, faintly yellow oil.

15

Part B

A solution of 2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (1.00 g, 4.13 mmol) in 20 mL of acetonitrile and 5 mL of glacial acetic acid was treated with *tert*-butyl (3,3-diethoxypropyl)carbamate (2.55 g, 10.3 mmol) and heated to reflux under an atmosphere of nitrogen. After 15 h, the reaction mixture was concentrated under reduced pressure to yield a brown oil. The oil was partitioned between CHCl₃ and saturated NaHCO₃ solution and the phases were separated. The organic portion was washed with water (2X) and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield *tert*-butyl {3-[(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)imino]propyl}carbamate (1.64 g) as a dark red/orange oil.

25

Part C

A solution of *tert*-butyl {3-[(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)imino]propyl} carbamate (1.64 g, 4.13 mmol) in 20 mL of methanol was treated with NaBH₄ (0.78 g, 20.6 mmol) under an atmosphere of nitrogen. After 1.5 h, the reaction mixture was quenched with saturated NH₄Cl solution and concentrated under reduced pressure. The residue was partitioned between saturated NaHCO₃ solution and CHCl₃ and the phases were separated. The organic portion was washed with water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a light brown solid. Chromatography [SiO₂, 95:5 CHCl₃:(80:18:2 CHCl₃:MeOH:NH₄OH)] yielded *tert*-butyl {3-[(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl} carbamate (1.34 g) as a tan foam.

Part D

A solution of *tert*-butyl {3-[(2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl} carbamate (1.34 g, 3.35 mmol) in 30 mL of CHCl₃ was treated with MCPBA (1.45 g, 5.03 mmol, 77% max). After 3 h, the reaction mixture was diluted with 10% Na₂CO₃ solution and CHCl₃ and the phases were separated. The organic portion was washed with water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield *tert*-butyl {3-[(2-ethoxymethyl-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl} carbamate (1.39 g) as an orange foam.

Part E

A solution of *tert*-butyl {3-[(2-ethoxymethyl-5-oxido-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl} carbamate (1.39 g, 3.35 mmol) in 35 mL of CHCl₃ was treated with 15 mL of concentrated NH₄OH solution and *p*-toluenesulfonyl chloride (0.67 g, 3.51 mmol). After 15 min, the reaction mixture was diluted with water and CHCl₃ and the phases were separated. The organic portion was washed with 10% Na₂CO₃ solution and water. The combined aqueous washes were back-extracted with CHCl₃. The combined organic extracts were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield *tert*-butyl {3-[(4-amino-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl} carbamate (1.30 g) as an orange foam.

Part F

A solution of *tert*-butyl {3-[(4-amino-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl} carbamate (1.30 g, 3.14 mmol) in 10 mL of ethanol was treated with a solution of 3 M hydrogen chloride in ethanol (5.0 mL, 15 mmol) and heated to 100 °C.


5 After 30 min, the solvent was concentrated under reduced pressure to yield a brown sludge. The material was triturated with diethyl ether and filtered to give a tan solid. The solid was dissolved in water and treated with 10% NaOH solution until pH 13 was reached. The aqueous solution was extracted with CH₂Cl₂ (4X). The combined organic extracts were washed with brine, dried over Na₂SO₄, filtered and concentrated under
10 reduced pressure to yield *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.77 g) as a gold colored foam.

Part G

A solution of *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-
15 1,4-diamine (0.250 g, 0.795 mmol) in 10 mL of CH₂Cl₂ was treated with triethylamine (0.221 mL, 1.67 mmol) under an atmosphere of nitrogen and cooled to 0 °C. The reaction mixture was treated dropwise with methanesulfonyl chloride (0.065 mL, 0.835 mmol). After 16 h, the reaction mixture was quenched by 10% Na₂CO₃ solution, diluted with CHCl₃ and the phases were separated. The organic portion was washed with water and
20 brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a light yellow solid. Chromatography (SiO₂, 95:5 CHCl₃:MeOH) gave an off-white foam. The foam was triturated with diethyl ether and filtered to give *N*-{3-[(4-amino-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}methanesulfonamide (0.164 g) as an off white solid. mp 148–150 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.46 (d, *J* =
25 7.8 Hz, 1 H), 7.58 (d, *J* = 8.2 Hz, 1 H), 7.44 (t, *J* = 7.1 Hz, 1 H), 7.25 (t, *J* = 7.4 Hz, 1 H), 7.05-6.95 (m, 2 H), 6.61 (s, 2 H), 4.76 (s, 2 H), 3.62 (q, *J* = 7.0 Hz, 2 H), 3.22 (q, *J* = 6.8 Hz, 2 H), 3.07 (q, *J* = 6.2 Hz, 2 H), 2.88 (s, 3 H), 1.78 (p, *J* = 6.3 Hz, 2 H), 1.18 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (125 MHz, DMSO-*d*₆) δ 152.3, 149.5, 145.3, 132.5, 127.4, 126.1, 124.2, 121.3, 121.3, 114.7, 65.9, 63.1, 49.9, 39.6, 28.1, 15.4; MS (APCI) *m/z* 393 (*M* +
30 H)⁺; Anal. Calcd for C₁₇H₂₄N₆O₃: C, 52.03; H, 6.16; N, 21.41; Found: C, 51.84; H, 6.28; N, 21.18.

Example 15


1-{3-[(4-Amino-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}-3-phenylurea



5 Part A

A solution of *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.250 g, 0.795 mmol) in 10 mL of CH₂Cl₂ was cooled to 0 °C under an atmosphere of nitrogen. The reaction mixture was treated dropwise with phenyl isocyanate (0.091 mL, 0.835 mmol). After 16 h, the reaction mixture was quenched by 10% Na₂CO₃ solution, diluted with CHCl₃ and the phases were separated. The organic portion was washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield an off-white solid. Chromatography (SiO₂, 95:5 CHCl₃:MeOH) gave an off-white foam. The foam was triturated with diethyl ether and filtered to give 1-{3-[(4-amino-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}-3-phenylurea (0.115 g) as an off-white solid. mp 177–179 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.46 (dd, *J* = 8.1, 1.0 Hz, 1 H), 8.39 (s, 1 H), 7.58 (dd, *J* = 8.4, 0.9 Hz, 1 H), 7.44–7.35 (m, 3 H), 7.25–7.18 (m, 3 H), 6.99 (t, *J* = 5.6 Hz, 1 H), 6.90–6.85 (m, 1 H), 6.60 (s, 2 H), 6.16 (t, *J* = 5.6 Hz, 1 H), 4.76 (s, 2 H), 3.60 (q, *J* = 7.0 Hz, 2 H), 3.26–3.18 (m, 4 H), 1.76 (t, *J* = 7.0 Hz, 2 H), 1.15 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (125 MHz, DMSO-*d*₆) δ 155.2, 151.8, 149.0, 144.8, 140.4, 132.0, 128.5, 126.9, 125.7, 123.7, 120.9, 120.8, 120.8, 117.6, 114.3, 65.4, 62.7, 49.7, 37.0, 28.1, 14.9; MS (APCI) *m/z* 434 (M + H)⁺; Anal. Calcd for C₂₃H₂₇N₇O₂: C, 63.72; H, 6.28; N, 22.62; Found: C, 63.45; H, 6.04; N, 22.28.

Example 16

*N*¹-Isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

Part A

5 A suspension of *tert*-butyl *N*'-(3-aminoquinolin-4-yl)hydrazinecarboxylate (6.50 g, 23.7 mmol) in 100 mL of toluene was treated with trimethyl orthobutyrate (4.18 mL, 26.1 mmol) and pyridine hydrochloride (0.14 g, 1.2 mmol) and heated to 130 °C under an atmosphere of nitrogen. After 18 h, the reaction mixture was concentrated under reduced pressure to yield a brown oil. The oil was dissolved in 150 mL CHCl₃, washed with water
10 (2 X 50 mL), brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 7.23 g of *tert*-butyl (2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate as an orange foam.

Part B

15 A solution of *tert*-butyl (2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (7.23 g, 22.2 mmol) in 40 mL of ethanol was treated with HCl (37 mL, 111 mmol, 3 M in ethanol) and heated to reflux. After 1 h, the reaction mixture was cooled to ambient temperature, diluted with 80 mL of diethyl ether, and cooled in an ice water bath. The HCl salt of the product was collected by vacuum filtration and rinsed with diethyl ether
20 until the filtrate ran clear. The dried HCl salt was dissolved in 75 mL of water and treated with 50% NaOH solution until the pH of the water was 12-13. The free base of the product precipitated out and was triturated in the basic water for 30 min while being cooled in an ice water bath. The solid was collected by vacuum filtration and dried under vacuum to give 4.64 g of 2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine as a tan granular
25 solid.

Part C

 A solution of 2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (4.64 g, 20.5 mmol) in 60 mL of acetonitrile and 15 mL of glacial acetic acid was treated with 2,2-

dimethoxypropane (12.6 mL, 103 mmol) and heated to 100 °C under an atmosphere of nitrogen. After 6 d, the reaction mixture was concentrated under reduced pressure to yield a brown oil. The oil was dissolved in 100 mL of CHCl₃ and washed with 10% Na₂CO₃ (2 X 25 mL), water (25 mL), brine (25 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 4.30 g of *N*-isopropylidene-(2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine as a brown oil.

Part D

A solution of *N*-isopropylidene-(2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (4.30 g, 16.1 mmol) in 100 mL of methanol was cooled in an ice water bath. The solution was treated with sodium borohydride (3.05 g, 80.7 mmol) over 5 min. The reaction mixture was allowed to warm to ambient temperature. After 2.5, the reaction was quenched by addition of 15 mL of saturated NH₄Cl solution. The mixture was concentrated under reduced pressure to yield a light brown solid. The solid was partitioned between 100 mL CHCl₃ and 25 mL of saturated NaHCO₃ solution and then separated. The organic portion was washed with water (25 mL), brine (25 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a light brown solid. The solid was purified by chromatography (SiO₂, 97:2.5:0.5 CHCl₃:MeOH:NH₄OH) to give 2.48 g of *N*-isopropyl-(2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine as a tan solid.

Part E


A solution of *N*-isopropyl-(2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amine (2.48 g, 9.24 mmol) in 75 mL of chloroform was cooled in a cold water bath. The solution was treated with MCPBA (3.32 g, 11.6 mmol) over 6 min. The reaction was allowed to come to ambient temperature. After 1.5 h, TLC showed complete conversion to the 5-*N*-oxide intermediate. The reaction mixture was again cooled in a cold water bath and then treated with concentrated ammonium hydroxide solution (30 mL, 30%) and stirred rapidly. The reaction mixture was treated with *p*-toluenesulfonyl chloride (1.85 g, 9.70 mmol) over 5 min. The reaction was allowed to come to ambient temperature. After 30 min, the reaction mixture was diluted with 50 mL of chloroform and 30 mL of water and the phases were separated. The organic portion was washed with 5% Na₂CO₃ solution (30 mL), water (30 mL) and brine (30 mL). The organic portion was dried over Na₂SO₄, filtered

and concentrated under reduced pressure to yield a light brown foam. The material was purified by chromatography (SiO₂, 97:3 CHCl₃:MeOH) and recrystallized from EtOAc to yield 1.39 g of *N*¹-isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine as amber crystals.

5 mp 181–184 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.44 (d, *J* = 8.1 Hz, 1 H), 7.57 (d, *J* = 8.3 Hz, 1 H), 7.41–7.35 (m, 1 H), 7.23–7.18 (m, 1 H), 6.95 (d, *J* = 1.6 Hz, 1 H), 6.48 (s, 2 H), 3.52–3.45 (m, 1 H), 2.98–2.85 (m, 2 H), 1.91–1.79 (m, 2 H), 1.03–0.98 (m, 9 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 154.5, 152.0, 144.9, 132.6, 126.8, 126.1, 124.2, 121.2, 120.9, 115.0, 51.2, 28.2, 21.1, 20.6, 14.3; MS (APCI) *m/z* 284 (M + H)⁺; Anal. Calcd for C₁₆H₂₁N₅: C, 67.82; H, 7.47; N, 24.71; Found: C, 67.66; H, 7.39; N, 24.66.

Example 17

*N*¹-Isopropyl-2-propyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine




15 Part A

A solution of *N*¹-isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.59 g, 2.1 mmol) in 15 mL of trifluoroacetic acid was treated with platinum(IV) oxide (0.55 g, 2.4 mmol) and shaken under an atmosphere of hydrogen (3.8 x 10⁵ Pa). After 6 days, the reaction mixture was filtered through a pad of CELITE filter agent and rinsed with a mixture of 85:15:0.1 CHCl₃:MeOH:TFA until the filtrate ran clear. The filtrate was concentrated under reduced pressure to yield a white foam. The material was suspended in water and treated with 50 % NaOH solution until the pH reached 13. A white solid precipitated and was triturated in the basic mixture for 1 h. The white solid was collected by vacuum filtration. The solid was purified by chromatography (SiO₂, 95:5:0.1 CHCl₃:MeOH:NH₄OH) to yield 0.23 g of *N*¹-isopropyl-2-propyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine as a white solid.

mp 162–164 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 6.34 (s, 1 H), 5.64 (s, 2 H), 3.38–3.23 (m, 2 H), 2.85–2.79 (m, 3 H), 2.78–2.71 (m, 2 H), 1.84–1.71 (m, 6 H), 0.99–0.86 (m, 9 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 154.4, 149.3, 146.1, 137.9, 122.8, 105.7, 52.4, 32.5,

28.4, 23.3, 23.1, 22.9, 21.0, 20.7, 14.3; MS (APCI) m/z 288 ($M + H$)⁺; Anal. Calcd for C₁₆H₂₅N₅: C, 66.87; H, 8.77; N, 24.37; Found: C, 66.65; H, 8.90; N, 24.08.

Example 18

5 *N*¹-Isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

Part A

A suspension of *tert*-butyl *N*-(3-aminoquinolin-4-yl)hydrazinecarboxylate (6.50 g, 23.7 mmol) in 100 mL of toluene was treated with triethyl orthoformate (8.68 mL, 52.2 mmol) and pyridine hydrochloride (0.14 g, 1.2 mmol) and heated to 130 °C under an atmosphere of nitrogen. After 23 h, the reaction mixture was concentrated under reduced pressure to yield a red/brown oil. The oil was dissolved in CHCl₃ (150 mL) and washed with water (2 X 50 mL), brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 6.74 of *tert*-butyl *N*-(1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate as a red/orange oil.

Part B

A solution of *tert*-butyl *N*-(1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (6.74 g, 23.7 mmol) in 40 mL of ethanol was treated with 40 mL of HCl (40 mL, 119 mmol, 3 M in ethanol) and heated to reflux. After 1 h, the reaction mixture was cooled to ambient temperature, diluted with 80 mL of diethyl ether, and cooled in an ice water bath which precipitated a tan solid. The HCl salt of the product was collected by vacuum filtration and rinsed with diethyl ether until the filtrate ran clear. The dried HCl salt was dissolved in 75 mL of water and made basic by addition of 50% NaOH solution until the pH of the water was 12-13. The free base of the product precipitated out and was triturated in the basic water for 30 min while being cooled in an ice water bath. The solid was collected by vacuum filtration and dried under vacuum to give 2.86 g of 1*H*-imidazo[4,5-*c*]quinolin-1-amine as a tan granular solid.

Part C

A solution of 1*H*-imidazo[4,5-*c*]quinolin-1-amine (2.86 g, 15.5 mmol) in 60 mL of acetonitrile and 15 mL of glacial acetic acid was treated with 2,2-dimethoxypropane (9.53 mL, 77.5 mmol) and heated to 100 °C under an atmosphere of nitrogen. After 18 h, the reaction mixture was concentrated under reduced pressure to give a brown oil. The oil was dissolved in 100 mL of CHCl₃ and washed with 5% Na₂CO₃ solution (2 X 30 mL), water (30 mL) and brine (30 mL). The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 3.48 g of *N*-(1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylideneamine as a brown oil.

Part D

A solution of *N*-(1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylideneamine (3.48 g, 15.5 mmol) in 75 mL of methanol was cooled in an ice water bath. The solution was treated over 5 min with sodium borohydride (2.94 g, 77.6 mmol). After 1 h, the reaction mixture was quenched with 20 mL of saturated NH₄Cl solution and then concentrated under reduced pressure to yield a brown solid. The solid was partitioned between 80 mL CHCl₃ and 20 mL saturated NaHCO₃ solution and the phases were separated. The organic portion was washed with water (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a brown solid. The solid was purified by chromatography (SiO₂, 95:5:0.5 CHCl₃:MeOH:NH₄OH) to give 1.28 g of *N*-(1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine as a tan foam.

Part E


A solution of *N*-(1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (1.36 g, 5.66 mmol) in 50 mL of chloroform was cooled in a cold water bath. The solution was treated with MCPBA (2.03 g, 7.07 mmol) over 5 min and then allowed to warm to ambient temperature. After 1 h, TLC showed complete conversion to the intermediate 5-*N*-oxide. The reaction mixture was again cooled with a cold water bath. The solution was treated with concentrated ammonium hydroxide solution (25 mL, 30%) and stirred rapidly to homogenize. The reaction mixture was treated with *p*-toluenesulfonyl chloride (1.13 g, 5.94 g) over 5 min and allowed to warm to ambient temperature. After 30 min, the reaction mixture was diluted with 50 mL of CHCl₃ and 25 mL of water. An undissolved

solid between the phases was filtered off, saved, and the phases were separated. The organic portion was washed with saturated NaHCO₃ solution (30 mL), water (30 mL) and brine (30 mL). The organic portion was then dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan/orange solid. A high-performance liquid chromatography (HPLC) analysis of the filtered solid matched that of the solid from the concentrated organic extracts. The combined solid was recrystallized twice from MeOH to give 1.18 g of *N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine as an off-white solid.

mp dec. > 250 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.61 (dd, *J* = 8.1, 1.1 Hz, 1 H), 8.23 (s, 1 H), 7.56 (d, *J* = 7.6 Hz, 1 H), 7.43-7.37 (m, 1 H), 7.23-7.18 (m, 1 H), 7.04 (d, *J* = 3.4 Hz, 1 H), 6.58 (s, 2 H), 3.57-3.47 (m, 1 H), 1.03 (d, *J* = 6.2 Hz, 6 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 152.4, 145.3, 132.3, 127.3, 126.0, 125.1, 121.5, 121.0, 115.1, 52.6, 20.6; MS (APCI) *m/z* 242 (M + H)⁺; Anal. Calcd for C₁₃H₁₅N₅: C, 64.71; H, 6.27; N, 29.02; Found: C, 63.11; H, 6.30; N, 27.96.

Example 19

*N*¹-Isopropyl-2-propyl-7-(pyridin-3-yl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine



Part A

A suspension of 7-bromo-4-chloro-3-nitroquinoline (75.00 g, 260.9 mmol) in 350 mL of dichloromethane was cooled to 0 °C under an atmosphere of nitrogen. The suspension was treated with triethylamine (43.25 mL, 326.1 mmol), which dissolved most of the material. A solution of *tert*-butyl carbazate (37.93 g, 287.0 mmol) in 250 mL of dichloromethane was added to the reaction mixture over 20 min. The reaction was allowed to slowly come to ambient temperature. After 15 h, the reaction mixture was washed with 5% Na₂CO₃ solution (2 X 100 mL) and water (100 mL). The combined aqueous washes were back-extracted with CHCl₃ (50 mL). The combined organic portions were washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated

under reduced pressure to yield 99.98 g of *tert*-butyl *N'*-(7-bromo-3-nitroquinolin-4-yl)hydrazinecarboxylate as a dark red solid.

Part B

5 A suspension of *tert*-butyl *N'*-(7-bromo-3-nitroquinolin-4-yl)hydrazinecarboxylate (50.0 g, 131 mmol) in 320 mL of acetonitrile (MeCN) and 80 mL of methanol was treated with platinum on carbon (5.0 g, 1.3 mmol, 5% w/w) and shaken under an atmosphere of hydrogen (3.8×10^5 Pa). After 4 h, the reaction mixture was filtered through a pad of
10 CELITE filter agent and rinsed with portions of MeCN:MeOH (1:1) until the filtrate ran clear. The filtrate was concentrated under reduced pressure to yield 37.1 g of *tert*-butyl *N'*-(3-amino-7-bromoquinolin-4-yl)hydrazinecarboxylate as a tan solid.

Part C

15 A solution of *tert*-butyl *N'*-(3-amino-7-bromoquinolin-4-yl)hydrazinecarboxylate (37.1 g, 105 mmol) in 315 mL of toluene was treated with trimethyl orthobutyrate (16.7 mL, 105 mmol) and pyridine hydrochloride (0.12 g, 1.05 mmol). The reaction mixture was heated to reflux under an atmosphere of nitrogen. After 4 h, the reaction mixture was cooled to ambient temperature and concentrated under reduced pressure to give a brown oil. The oil was dissolved in 300 mL of CHCl_3 . The solution was washed with 5%
20 Na_2CO_3 (100 mL), water (100 mL) and brine (100 mL). The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield a brown foam. The foam was purified by chromatography (SiO_2 , 100:0 gradient to 95:5 CHCl_3 :MeOH) to yield 30.1 g of *tert*-butyl (7-bromo-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate as a light brown solid.

25

Part D

30 A suspension of *tert*-butyl (7-bromo-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (30.1 g, 74.3 mmol) in 25 mL of ethanol was treated with HCl in ethanol (86.4 mL, 37.1 mmol, 4.3 M) and heated to 100 °C. After 30 min, the reaction mixture was cooled to ambient temperature and concentrated under reduced pressure to yield a brown solid. The solid was suspended in 100 mL of water, stirred vigorously and treated with 50% NaOH solution until the pH of the liquid rose to 12-13. A brown solid collected

around the stir bar. The water was diluted with 200 mL of dichloromethane and the solid was broken apart. The material was triturated in the biphasic mixture overnight. After triturating for 15 h, the mixture was filtered to give the crude free base as a light brown solid. The solid was dried under vacuum to give 17.6 g of 7-bromo-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine as a light brown solid.

Part E

A suspension of 7-bromo-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (17.6 g, 57.7 mmol) in 160 mL of acetonitrile and 40 mL of glacial acetic acid was treated with 2,2-dimethoxypropane (35.5 mL, 288 mmol). The reaction mixture was heated to 100° C under an atmosphere of nitrogen. After 16 h, the reaction was cooled to ambient temperature and concentrated under reduced pressure to yield a brown oil. The oil was dissolved in CHCl₃ (200 mL). The CHCl₃ solution was washed with saturated NaHCO₃ solution (2 X 50 mL), water (50 mL) and brine (50 mL). The organic portion was then dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 18.4 g of *N*-(7-bromo-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylideneamine as a red/brown foam.

Part F

A solution of *N*-(7-bromo-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylideneamine (18.4 g, 53.3 mmol) in 100 mL of methanol was placed under an atmosphere of nitrogen and cooled in an ice water bath. The solution was treated with sodium borohydride (2.32 g, 61.3 mmol) over 30 min. The reaction mixture was allowed to slowly come to ambient temperature. After 1.5 h, the reaction was quenched by the addition of 25 mL of saturated NH₄Cl solution. The reaction mixture was concentrated under reduced pressure to remove the methanol. The residue was partitioned between chloroform (150 mL) and 10% Na₂CO₃ solution (35 mL), and the phases were separated. The organic portion was washed with another portion of 10% Na₂CO₃ solution (35 mL), water (35 mL) and brine (35 mL). The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a brown foam. The foam was purified by chromatography (SiO₂, 97:3 CHCl₃:MeOH gradient to 9:1) to give 16.3 g of *N*-(7-bromo-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine as a dark tan solid.

Part G

A solution of *N*-(7-bromo-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (9.10 g, 26.2 mmol) in 200 mL of chloroform was placed under an atmosphere of nitrogen and cooled in an ice water bath. The solution was treated with MCPBA (8.28 g, 28.8 mmol, 77% max) and allowed to slowly come to ambient temperature. After 2 h, LC/MS and HPLC indicated complete conversion to the 5-*N*-oxide intermediate. The reaction mixture was again cooled in an ice water bath. The reaction mixture was treated with ammonium hydroxide solution (50 mL, 30%) and stirred vigorously. The mixture was treated with *p*-toluenesulfonyl chloride (5.24 g, 27.5 mmol) and allowed to come to ambient temperature. After 30 min, the reaction was diluted with 50 mL of water, and the phases were separated. The organic portion was washed with water (75 mL), brine (75 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a light brown solid. The solid was purified by chromatography (SiO₂, 95:5 CHCl₃:MeOH) and then recrystallized from acetonitrile to give 4.52 g of 7-bromo-*N*¹-isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine as off white crystals. mp 226–228 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.44 (d, *J* = 8.7 Hz, 1 H), 7.71 (d, *J* = 2.1 Hz, 1 H), 7.36 (dd, *J* = 8.7, 2.1 Hz, 1 H), 6.99 (d, *J* = 1.7 Hz, 1 H), 6.73 (s, 2 H), 3.53–3.40 (m, 1 H), 2.90 (s, 2 H), 1.93–1.80 (m, 2 H), 1.05–1.00 (m, 9 H); ¹³C NMR (125 MHz, DMSO-*d*₆) δ 154.9, 152.9, 146.3, 132.5, 127.8, 124.2, 123.5, 123.1, 119.7, 114.0, 79.5, 51.4, 28.2, 21.1, 20.6, 14.3; MS (APCI) *m/z* 362, 364 (*M* + *H*)⁺; Anal. Calcd for C₁₆H₂₀BrN₅·0.25H₂O: C, 52.40; H, 5.63; N, 19.09; Found: C, 52.03; H, 5.42; N, 19.14.

Part H


A suspension of 7-bromo-*N*¹-isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.00 g, 2.76 mmol) in 20 mL of 1-propanol was treated with pyridine-3-boronic acid 1,3-propane diol cyclic ester (0.540 g, 3.31 mmol). The head-space of the reaction flask was purged and back-filled with nitrogen (3X). The reaction mixture was then treated with triphenylphosphine (11 mg, 0.041 mmol), sodium carbonate (1.66 mL, 3.31 mmol, 2 M solution in water), water (2 mL) and palladium(II) acetate (3.1 mg, 0.014 mmol). Again the head-space of the reaction flask was purged and back-filled with nitrogen (3X). The reaction was heated to 100° C. After 17 h, the reaction was cooled to

ambient temperature and concentrated under reduced pressure to yield a brown solid. The solid was dissolved and partitioned between 15 mL of water and 15 mL of chloroform and then separated. The aqueous portion was extracted with chloroform (2 X 15 mL). The combined organic extracts were washed with brine (15 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan solid. The solid was purified by chromatography (SiO₂, 95:5 CHCl₃:MeOH) and recrystallized from acetonitrile to give 0.515 g of *N*¹-isopropyl-2-propyl-7-(pyridin-3-yl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine as white crystals.

mp 218–219 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.99 (d, *J* = 1.7 Hz, 1 H), 8.60–8.57 (m, 2 H), 8.19–8.16 (m, 1 H), 7.88 (d, *J* = 1.9 Hz, 1 H), 7.61 (dd, *J* = 8.5, 1.9 Hz, 1 H), 7.53–7.49 (m, 1 H), 7.04 (s, 1 H), 6.59 (s, 2 H), 3.57–3.49 (m, 1 H), 2.92–2.87 (m, 2 H), 1.94–1.82 (m, 2 H), 1.06–1.01 (m, 9 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 154.8, 152.5, 148.6, 148.1, 145.4, 136.2, 135.4, 134.5, 132.5, 124.5, 124.3, 123.9, 122.2, 119.6, 114.7, 51.3, 28.2, 21.1, 20.6; MS (APCI) *m/z* 361 (M + H)⁺; Anal. Calcd for C₂₁H₂₄N₆: C, 69.97; H, 6.71; N, 23.31; Found: C, 69.78; H, 6.55; N, 23.51.

Example 20

7-Benzyloxy-2-ethoxymethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine



Part A

A mixture of triethyl orthoformate (92 mL, 0.55 mol) and 2,2-dimethyl-1,3-dioxane-4,6-dione (75.3 g, 0.522 mol) (Meldrum's acid) was heated at 55 °C for 90 minutes and then cooled to 45 °C. A solution of 3-benzyloxyaniline (100.2 g, 0.5029 mol) in methanol (200 mL) was slowly added to the reaction over a period 45 minutes while maintaining the reaction temperature below 50 °C. The reaction was then heated at 45 °C for one hour, allowed to cool to room temperature, and stirred overnight. The reaction mixture was cooled to 1 °C, and the product was isolated by filtration and washed with cold ethanol (~400 mL) until the filtrate was colorless.

Benzyloxy)phenylimino]methyl}-2,2-dimethyl-1,3-dioxane-4,6-dione (170.65 g) was isolated as a tan, powdery solid.

¹H NMR (300 MHz, DMSO-*d*₆) δ 11.21 (d, *J* = 14.2 Hz, 1H), 8.61 (d, *J* = 14.2 Hz, 1H), 7.49-7.30 (m, 7H), 7.12 (dd, *J* = 8.1, 1.96 Hz, 1H), 6.91 (dd, *J* = 8.4, 2.1 Hz, 1H), 5.16 (s, 2H), 1.68 (s, 6H).

Part B

A mixture of 5-{[(3-benzyloxy)phenylimino]methyl}-2,2-dimethyl-1,3-dioxane-4,6-dione (170.65 g, 0.483 mol) and DOWTHERM A heat transfer fluid (800 mL) was heated to 100 °C and then slowly added to a flask containing DOWTHERM A heat transfer fluid (1.3 L, heated at 210 °C) over a period of 40 minutes. During the addition, the reaction temperature was not allowed to fall below 207 °C. Following the addition, the reaction was stirred at 210 °C for one hour, and then allowed to cool to ambient temperature. A precipitate formed, which was isolated by filtration, washed with diethyl ether (1.7 L) and acetone (0.5 L), and dried in an oven to provide 76.5 g of 7-benzyloxyquinolin-4-ol as a tan powder.

¹H NMR (300 MHz, DMSO-*d*₆) δ 11.53 (s, 1H), 7.99 (dd, *J* = 7.4, 2.4 Hz, 1H), 7.79 (d, *J* = 7.4 Hz, 1H), 7.50-7.32 (m, 5H), 7.00 (s, 1H), 6.98 (dd, *J* = 7.4, 2.5 Hz, 1H), 5.93 (d, *J* = 7.5 Hz, 1H), 5.20 (s, 2H).

Part C

A mixture of 7-benzyloxyquinolin-4-ol (71.47 g, 0.2844 mol) and propionic acid (700 mL) was heated to 125 °C with vigorous stirring. Nitric acid (23.11 mL of 16 M) was slowly added over a period of 30 minutes while maintaining the reaction temperature between 121 °C and 125 °C. After the addition, the reaction was stirred at 125 °C for 1 hour then allowed to cool to ambient temperature. The resulting solid was isolated by filtration, washed with water, and dried in an oven for 1.5 days to provide 69.13 g of 7-benzyloxy-3-nitroquinolin-4-ol as a grayish powder.

¹H NMR (300 MHz, DMSO-*d*₆) δ 12.77 (s, 1H), 9.12 (s, 1H), 8.17 (dd, *J* = 6.3, 3.3 Hz, 1H), 7.51-7.33 (m, 5H), 7.21-7.17 (m, 2H), 5.25 (s, 2H).

Part D

A suspension of 7-benzyloxy-3-nitroquinolin-4-ol (75.0 g, 253 mmol), which was made in a separate run, in 500 mL of *N,N*-dimethylformamide was placed under an atmosphere of nitrogen. The suspension was treated with phosphorous oxychloride (27.8 mL, 304 mmol) dropwise over 1.5 h. After 18 h, the reaction mixture was cooled to 0 °C and then poured into 1 L of ice water. The mixture was stirred until the ice had melted. A tan/yellow precipitate was collected by vacuum filtration. The solid was dissolved in dichloromethane (500 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 71.7 g of 7-benzyloxy-4-chloro-3-nitro-quinoline as an orange solid.

10

Part E

A solution of *tert*-butyl carbazate (33.1 g, 251 mmol) in 150 mL of dichloromethane was treated with triethylamine (66.5 mL, 502 mmol). The solution was placed under an atmosphere of nitrogen and cooled in a cold-water bath. The solution was treated with a solution of 7-benzyloxy-4-chloro-3-nitroquinoline (71.7 g, 228 mmol) in 350 mL of dichloromethane over 1 h. The reaction was stirred and allowed to warm to ambient temperature. After 15 h, the reaction was diluted with 200 mL of water and 250 mL of CHCl₃ and the phases were separated. The organic portion was washed with water (200 mL), brine (200 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield an orange solid. The solid was recrystallized from dichloromethane to yield 53.5 g of *tert*-butyl *N'*-(7-benzyloxy-3-nitroquinolin-4-yl)hydrazinecarboxylate as yellow crystals.

15

20

Part F

A solution of *tert*-butyl *N'*-(7-benzyloxy-3-nitroquinolin-4-yl)hydrazinecarboxylate (20.00 g, 48.73 mmol) in 200 mL of methanol and 200 mL of acetonitrile was treated with platinum on carbon (2.00 g, 0.51 mmol) and shaken under an atmosphere of hydrogen (3.8 x 10⁵ Pa). After 17 h, the mixture was filtered through a pad of CELITE filter agent and rinsed with MeOH:MeCN (1:1) until the filtrate ran clear. The filtrate was concentrated under reduced pressure to yield 18.21 g of *tert*-butyl *N'*-(3-amino-7-benzyloxyquinolin-4-yl)hydrazinecarboxylate as a red/orange solid.

25

30

Part G

A suspension of *tert*-butyl *N'*-(3-amino-7-benzyloxyquinolin-4-yl)hydrazinecarboxylate (29.6 g, 77.8 mmol) in 250 mL of 1,2-dichloroethane was placed under an atmosphere of nitrogen. The mixture was treated with triethylamine (30.9 mL, 233 mmol). The mixture was then treated dropwise with ethoxyacetyl chloride (10.5 g, 85.6 mmol). After 2 h, the reaction was concentrated under reduced pressure to give a brown oil. The oil was dissolved in 200 mL of 1-butanol and treated with pyridinium *p*-toluenesulfonate (0.25 g, 1.0 mmol). The mixture was heated to 135 °C under an atmosphere of nitrogen. After 20 h, the reaction mixture was cooled to ambient temperature and concentrated under reduced pressure to give a brown oil. The oil was dissolved in 250 mL of CHCl₃ and washed with saturated NaHCO₃ solution (75 mL), water (75 mL) and brine (75 mL). The organic portion was then dried over Na₂SO₄, filtered and concentrated under reduced pressure to give an orange/brown oil. The oil was purified by chromatography (SiO₂, 9:1 CHCl₃:MeOH) to yield 14.4 g of *tert*-butyl (7-benzyloxy-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate as an orange/brown foam.

Part H

A suspension of *tert*-butyl (7-benzyloxy-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)carbamate (14.4 g, 32.1 mmol) in 100 mL of ethanol was treated with HCl in ethanol (38 mL, 160 mmol, 4.3 M). The mixture was heated to 100 °C under an atmosphere of nitrogen. After 2 h, the reaction mixture was cooled to ambient temperature at which point a solid precipitated from solution. The mixture was diluted with 100 mL of diethyl ether and the solid was triturated for 15 min. The solid was collected by vacuum filtration and washed with several portions of diethyl ether. The solid was dried under vacuum for 2 h. The dry solid was suspended in 150 mL of water and treated with 50% NaOH solution until the pH of the liquid was 12. A brown solid precipitated. The mixture was diluted with 200 mL of CH₂Cl₂ and stirred until the solid dissolved. The layers were then separated. The aqueous portion was extracted with CH₂Cl₂ (2 X 100 mL). The combined organic extracts were washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 6.91 g of 7-benzyloxy-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine as a dark tan solid.

Part I

A suspension of 7-benzyloxy-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-amine (6.91 g, 19.8 mmol) in 55 mL of acetonitrile was treated with 2,2-dimethoxypropane (12.2 mL, 99.2 mmol) and 14 mL of glacial acetic acid. The reaction mixture was heated to 100 °C under an atmosphere of nitrogen. After 22 h, the reaction was cooled to ambient temperature and concentrated under reduced pressure to yield a brown oil. The oil was dissolved in 125 mL of CHCl₃ and washed with saturated NaHCO₃ solution (2 X 30 mL) and water (30 mL). The combined aqueous washes were back-extracted with CHCl₃ (25 mL). The combined organic extracts were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 7.69 g of *N*-(7-benzyloxy-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylideneamine as a brown solid.

Part J


A solution of *N*-(7-benzyloxy-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylideneamine (7.69 g, 19.8 mmol) in 50 mL of methanol was cooled to 0 °C. The solution was treated with sodium borohydride (1.12 g, 29.7 mmol) over 10 min. The reaction was allowed to slowly come to ambient temperature. After 2 h, the reaction was quenched with 15 mL of saturated NH₄Cl solution and concentrated under reduced pressure to yield a tan solid residue. The solid was dissolved in 100 mL of CHCl₃ and 25 mL of saturated K₂CO₃ solution then separated. The organic portion was washed with water (25 mL), brine (25 mL), dried over Na₂SO₄, filtered and concentrated to yield a brown oil. The oil was purified by chromatography (SiO₂, 98:2 CHCl₃:MeOH) to yield 6.63 g of *N*-(7-benzyloxy-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine as a tan foam.

Part K

A solution of *N*-(7-benzyloxy-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)isopropylamine (6.63 g, 17.0 mmol) in 90 mL of CHCl₃ was treated with MPCBA (6.29 g, 25.5 mmol, 70%). After 3 h, HPLC and LC/MS indicated complete conversion to the intermediate 5-*N*-oxide. The reaction mixture was then treated with concentrated ammonium hydroxide solution (30 mL, 30%). The biphasic reaction mixture was stirred

vigorously while *p*-toluenesulfonyl chloride (3.40 g, 17.9 mmol) was added. After 45 min, LC/MS indicated complete conversion to the 4-amine. The reaction mixture was diluted with 30 mL of water and 45 mL of CHCl₃ and separated. The organic portion was washed with 10% Na₂CO₃ solution (50 mL) and water (50 mL). The combined aqueous
5 portions were then back-extracted with CHCl₃ (25 mL). The combined organic portions were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated to yield a tan solid. The solid was purified by chromatography (SiO₂, 96:4 CHCl₃:MeOH) to give 5.90 g of 7-benzyloxy-2-ethoxymethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine as a light tan solid.
10 mp 194–196 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.47 (d, *J* = 8.9 Hz, 1 H), 7.50-7.48 (m, 2 H), 7.43-7.38 (m, 2 H), 7.35-7.30 (m, 1 H), 7.09 (d, *J* = 2.6 Hz, 1 H), 6.96 (dd, *J* = 9.0, 2.5 Hz, 1 H), 6.91 (d, *J* = 1.5 Hz, 1 H), 6.57 (s, 2 H), 5.20 (s, 2 H), 4.72 (s, 2 H), 3.64-3.57 (m, 3 H), 1.15 (t, *J* = 7.0 Hz, 3 H), 1.01 (d, *J* = 6.1 Hz, 6 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 157.9, 152.6, 149.4, 147.1, 137.7, 133.7, 128.8, 128.1, 128.0, 122.7, 111.8,
15 109.2, 108.4, 69.5, 65.8, 63.0, 51.6, 20.6, 15.3; MS (APCI) *m/z* 406 (M + H)⁺; Anal. Calcd for C₂₃H₂₇N₅O₂: C, 68.13; H, 6.71; N, 17.27; Found: C, 68.15; H, 6.91; N, 17.24.

Example 21

4-Amino-2-ethoxymethyl-1-isopropylamino-1*H*-imidazo[4,5-*c*]quinolin-7-ol


Part A

- 5 A solution of 7-benzyloxy-2-ethoxymethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.67 g, 4.12 mmol) in 25 mL of toluene and 25 mL of methanol was treated with palladium on carbon (0.44 g, 0.42 mmol, 10% w/w). The mixture was shaken under an atmosphere of hydrogen (3.8×10^5 Pa). After 16 h, the reaction was filtered through a pad of CELITE filter agent and rinsed with solvent until the filtrate ran clear. The filtrate was concentrated under reduced pressure to provide a white solid. Purification by chromatography (SiO₂, 3:1 CHCl₃:(80:18:2 CHCl₃:MeOH:NH₄OH) gradient to 1:1) gave 0.50 g of 4-amino-2-ethoxymethyl-1-isopropylamino-1*H*-imidazo[4,5-*c*]quinolin-7-ol as a white solid. MS (APCI) *m/z* 316 (M + H)⁺.

15

Example 22

tert-Butyl [3-(4-amino-2-ethoxymethyl-1-isopropylamino-1*H*-imidazo[4,5-*c*]quinolin-7-yloxy)propyl]carbamate



Part A

- 20 A solution of di-*tert*-butyl dicarbonate (19.05 g, 87.29 mmol) in tetrahydrofuran (20 mL) was added dropwise to a mixture of 3-amino-1-propanol (6.55 g, 87.2 mmol), tetrahydrofuran (50 mL), and 10% aqueous sodium hydroxide (35 mL). The reaction was stirred for 16 hours. The tetrahydrofuran was removed under reduced pressure, and the residue was adjusted to pH 3 with the slow addition of 15% aqueous potassium hydrogen sulfate. The mixture was extracted with ethyl acetate (3 x), and the combined organic

25

fractions were washed sequentially with water and brine, dried over anhydrous magnesium sulfate, filtered, and concentrated under reduced pressure to provide 16.6 g of *tert*-butyl 3-hydroxypropylcarbamate as a colorless oil containing some residual ethyl acetate.

5 Part B


Iodine (21.1 g, 83.1 mmol) was added in three portions to a solution of triphenylphosphine (19.83 g, 75.6 mmol) and imidazole (5.15 g, 75.6 mmol) in dichloromethane (300 mL). The resulting reddish-brown solution with a white precipitate was stirred until all of the iodine had dissolved. A solution of *tert*-butyl 3-hydroxypropylcarbamate (13.25 g, 75.61 mmol) in dichloromethane (150 mL) was added over a period of 45 minutes, and the reaction was stirred for 16 hours at ambient temperature. The reaction mixture was poured into saturated aqueous sodium thiosulfate and stirred until solution became colorless. The organic layer was separated and washed sequentially with saturated aqueous sodium thiosulfate, water, and brine; dried over anhydrous magnesium sulfate; filtered; and concentrated under reduced pressure to a pale yellow oil. The oil was purified by flash column chromatography (eluting with 80:20 hexanes:ethyl acetate) to a pale yellow oil which slowly crystallizes upon standing to afford 16.2 g of *tert*-butyl 3-iodopropylcarbamate as a yellow solid.

20 Part C

A solution of 4-amino-2-ethoxymethyl-1-isopropylamino-1*H*-imidazo[4,5-*c*]quinolin-7-ol (0.11 g, 0.35 mmol) in 10 mL of *N,N*-dimethylformamide was placed under an atmosphere of nitrogen and was treated with cesium carbonate (0.23 g, 0.70 mmol). After 5 min of stirring the mixture was treated with *tert*-butyl 3-iodopropylcarbamate (0.12 g, 0.35 mmol) and heated to 65 °C. After 60 h, the reaction mixture was cooled to ambient temperature and then poured into 100 mL of ice water which resulted in a cloudy suspension. The mixture was extracted with CHCl₃ (5 X 25 mL). The combined organic extracts were then washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan oil. Chromatography (95:5 CHCl₃:(80:18:2 CHCl₃:MeOH:NH₄OH) gradient to 1:1 gave 0.040 g of *tert*-butyl [3-(4-amino-2-ethoxymethyl-1-isopropylamino-1*H*-imidazo[4,5-*c*]quinolin-7-yloxy)propyl]carbamate as a light tan solid. LC/MS (APCI) *m/z* 473 (M+H)⁺.

Example 23

[3-(4-Amino-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-ylamino)propyl]morpholine-4-carboxamide



5

A solution of *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.500 g, 1.59 mmol) in 10 mL of CH₂Cl₂ was treated with triethylamine (0.443 mL, 3.34 mmol) under an atmosphere of nitrogen and cooled to 0 °C. The reaction mixture was treated dropwise with 4-morpholinecarbonyl chloride (0.065 mL, 0.835 mmol) and allowed to slowly come to ambient temperature. After 60 h, the reaction mixture was quenched with 10% Na₂CO₃ solution, diluted with CHCl₃ and the phases were separated. The organic portion was washed with water and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a light yellow solid. Chromatography (SiO₂, 9:1 CHCl₃:(80:18:2 CHCl₃:MeOH:NH₄OH) gradient to 1:1) gave a glassy solid. The solid was triturated with diethyl ether and filtered to give 0.046 g of [3-(4-amino-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinolin-1-ylamino)propyl]morpholine-4-carboxamide as a white solid.

15


20

mp 158–160 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.44 (d, *J* = 7.9 Hz, 1 H), 7.58 (d, *J* = 8.1 Hz, 1 H), 7.46–7.41 (m, 1 H), 7.26–7.21 (m, 1 H), 6.96 (t, *J* = 5.5 Hz, 1 H), 6.60 (s, 2 H), 6.53 (t, *J* = 5.1 Hz, 1 H), 4.75 (s, 2 H), 3.61 (q, *J* = 7.0 Hz, 2 H), 3.50 (t, *J* = 4.7 Hz, 4 H), 3.22–3.15 (m, 8 H), 1.72 (p, *J* = 6.9 Hz, 2 H), 1.17 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 158.0, 152.3, 149.5, 145.3, 132.4, 127.4, 126.1, 124.2, 121.2, 114.7, 66.3, 65.8, 63.1, 50.2, 44.1, 38.3, 28.5, 15.4; MS (APCI) *m/z* 428 (M + H)⁺; Anal. Calcd for C₂₁H₂₉N₇O₃: C, 59.00; H, 6.84; N, 22.93; Found: C, 58.76; H, 7.04; N, 22.82.

25

Example 24

N-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)cyclopentanecarboxamide




5

A solution of *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.500 g, 1.59 mmol) in 20 mL of CH₂Cl₂ was placed under an atmosphere of nitrogen, treated with triethylamine (0.443 mL, 3.34 mmol) and then cooled to 0 °C with an ice water bath. The reaction mixture was treated with cyclopentanecarbonyl chloride (0.203 mL, 1.67 mmol) over 2 min. The reaction was stirred and allowed to come to ambient temperature over 2 h. After 18 h, the reaction was quenched with 15 mL of 10% Na₂CO₃ solution. The mixture was diluted with 25 mL CHCl₃, mixed and then separated. The organic portion was washed with H₂O (15 mL) and brine (15 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan foam. The foam was purified by chromatography (SiO₂, 96:4 CHCl₃: methanol (MeOH)) to yield a light tan solid. The solid was dissolved in boiling 2-propanol (IPA), cooled to ambient temperature and then water was added until an off-white solid precipitated from solution. The precipitate was triturated in the IPA/H₂O mixture for 2 h and then collected by vacuum filtration to give *N*-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)cyclopentanecarboxamide (0.106 g) as an off-white solid. mp 75–77 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.42 (d, *J* = 7.9 Hz, 1 H), 7.74 (t, *J* = 5.2 Hz, 1 H), 7.58 (d, *J* = 7.9 Hz, 1 H), 7.46-7.41 (m, 1 H), 7.26-7.21 (m, 1 H), 6.97 (t, *J* = 5.4 Hz, 1 H), 6.60 (s, 2 H), 4.75 (s, 2 H), 3.61 (q, *J* = 7.0 Hz, 2 H), 3.19-3.13 (m, 4 H), 2.48-2.44 (m, 1 H), 1.75-1.38 (m, 10 H), 1.17 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 175.5, 152.3, 149.5, 145.3, 132.4, 127.3, 126.1, 124.2, 121.2, 114.7, 65.8, 63.1, 50.2, 44.7, 36.8, 30.3, 28.0, 25.9, 15.4; MS (APCI) *m/z* 411 (M + H)⁺; Anal. Calcd for C₂₂H₃₀N₆O₂•0.25H₂O: C, 63.67; H, 7.41; N, 20.25; Found: C, 36.37; H, 7.64; N, 20.32.

25

Example 25

1-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea




5

A solution of *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.04 g, 3.31 mmol) in 30 mL of CH₂Cl₂ and 5 mL of *N,N*-dimethylformamide was placed under an atmosphere of nitrogen and slowly treated with isopropyl isocyanate (0.320 mL, 3.31 mmol). After 15 h, the reaction mixture was concentrated under reduced pressure to yield a yellow/orange foam. The foam was purified by chromatography (SiO₂, 30% (80:18:2 CHCl₃:MeOH: conc. ammonium hydroxide (NH₄OH)) in CHCl₃) to give an off-white solid. The solid was triturated with diethyl ether (Et₂O) and collected by vacuum filtration to yield 1-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea (0.603 g) as an off-white solid. mp 174–177 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.45 (d, *J* = 8.1 Hz, 1 H), 7.58 (d, *J* = 8.2 Hz, 1 H), 7.46–7.41 (m, 1 H), 7.26–7.22 (m, 1 H), 6.96 (t, *J* = 5.5 Hz, 1 H), 6.60 (s, 2 H), 5.72 (t, *J* = 5.5 Hz, 1 H), 5.59 (d, *J* = 7.7 Hz, 1 H), 4.75 (s, 2 H), 3.70–3.58 (m, 3 H), 3.19–3.08 (m, 4 H), 1.72–1.63 (m, 2 H), 1.17 (t, *J* = 7.0 Hz, 3 H), 0.99 (d, *J* = 6.5 Hz, 6 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 157.8, 152.3, 149.5, 145.3, 132.4, 127.4, 126.1, 124.2, 121.2, 114.7, 65.9, 63.1, 50.2, 41.2, 37.4, 28.9, 23.6, 15.4; MS (APCI) *m/z* 400 (M + H)⁺; Anal. Calcd for C₂₀H₂₉N₇O₂: C, 60.13; H, 7.32; N, 24.54; Found: C, 59.94; H, 7.58; N, 24.45.

20

Example 26

1-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea




5

A solution of 1-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea (0.450 g, 1.13 mmol) in 25 mL of trifluoroacetic acid (TFA) was treated with platinum (IV) oxide (0.256 g, 1.13 mmol) and shaken under an atmosphere of hydrogen (3.8×10^5 Pa) at ambient temperature. After 72 h, the reaction mixture was diluted with 4:1 CHCl_3 :MeOH (25 mL) and filtered through a pad of CELITE filter agent. The filter pad was washed with additional 4:1 CHCl_3 :MeOH (1% TFA). The clear colorless filtrate was concentrated under reduced pressure to yield a white semi-solid. The material was suspended in 15 mL of H_2O and treated with 50% NaOH solution until the pH of the liquid reached 13. The material was triturated with the basic H_2O for 2 h. An off white solid was collected by vacuum filtration and dried. The material was purified by chromatography (SiO_2 , 30-50% (80:18:2 CHCl_3 :MeOH: NH_4OH) in CHCl_3) to yield the product as a white foam. The foam was triturated with Et_2O , filtered and dried to yield 1-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea (0.223 g) as a white powdery solid. mp 185–187 °C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 6.31 (t, $J = 6.0$ Hz, 1 H), 5.83 (s, 2 H), 5.71 (t, $J = 5.7$ Hz, 1 H), 5.60 (d, $J = 7.8$ Hz, 1 H), 4.61 (s, 2 H), 3.70-3.58 (m, 1 H), 3.55 (q, $J = 7.0$ Hz, 2 H), 3.11-2.98 (m, 6 H), 2.66 (t, $J = 5.8$ Hz, 2 H), 1.81-1.67 (m, 4 H), 1.59 (p, $J = 7.1$ Hz, 2 H), 1.14 (t, $J = 7.0$ Hz, 3 H), 1.00 (d, $J = 6.5$ Hz, 6 H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 157.8, 149.9, 149.3, 146.9, 137.6, 122.7, 105.5, 65.7, 63.3, 51.7, 41.2, 37.4, 32.4, 28.9, 23.6, 23.2, 23.0, 22.4, 15.3; MS (APCI) m/z 404 ($\text{M} + \text{H}$) $^+$; Anal. Calcd for $\text{C}_{20}\text{H}_{33}\text{N}_7\text{O}_2$: C, 59.53; H, 8.24; N, 24.30; Found: C, 59.28; H, 8.46; N, 24.04.

25

Example 27

N-{3-[4-Amino-1-(isopropylamino)-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-7-yl]phenyl}methanesulfonamide




5

A suspension of 7-bromo-*N*¹-isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.00 g, 2.76 mmol) in 20 mL of 1-propanol was treated with 3-(methanesulfonylamino)phenylboronic acid (0.653 g, 3.04 mmol). The headspace of the reaction flask was purged and back-filled with nitrogen (3X). The reaction mixture was then treated with triphenylphosphine (11 mg, 0.041 mmol), sodium carbonate (1.66 mL, 3.31 mmol, 2 M solution in H₂O), H₂O (2 mL) and palladium (II) acetate (3.1 mg, 0.014 mmol). Again the headspace of the reaction flask was purged and back-filled with nitrogen (3X). The reaction was heated to 100 °C. After 17 h, the reaction was cooled to ambient temperature and concentrated under reduced pressure to yield a brown solid. The solid was partitioned between 15 mL of H₂O and 15 mL of CHCl₃ and then separated. The aqueous portion was extracted with CHCl₃ (2 X 15 mL). The combined organic extracts were washed with brine (15 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan foam. The material was purified by chromatography (SiO₂, 50% (80:18:2 CHCl₃:MeOH:NH₄OH) in CHCl₃) and then dissolved in boiling CHCl₃ and slowly cooled to yield *N*-{3-[4-amino-1-(isopropylamino)-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-7-yl]phenyl}methanesulfonamide (0.700 g) as a white solid. mp 217–227 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.85 (s, 1 H), 8.54 (d, *J* = 8.5 Hz, 1 H), 7.79 (d, *J* = 1.4 Hz, 1 H), 7.60 (s, 1 H), 7.52-7.42 (m, 3 H), 7.22 (d, *J* = 7.8 Hz, 1 H), 7.01 (s, 1 H), 6.55 (s, 2 H), 3.55-3.51 (m, 1 H), 3.06 (s, 3 H), 2.92 (bs, 2 H), 1.91-1.84 (m, 2 H), 1.05-1.01 (m, 9 H); ¹³C NMR (125 MHz, DMSO-*d*₆) δ 155.4, 152.9, 145.8, 142.4, 139.9, 138.6, 133.1, 130.8, 124.9, 124.2, 123.1, 122.5, 120.2, 119.5, 118.7, 114.9, 51.8, 28.7, 21.6, 21.1, 14.8; MS (APCI) *m/z* 453 (M + H)⁺; Anal. Calcd for C₂₃H₂₈N₆O₂S: C, 61.04; H, 6.24; N, 18.57; Found: C, 60.66; H, 6.40; N, 18.47.

25

Example 28

N-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)morpholine-4-carboxamide




5

A solution of *N*-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)morpholine-4-carboxamide (0.120 g, 0.281 mmol) in 12 mL of TFA was treated with platinum (IV) oxide (0.064 g, 0.28 mmol). The mixture was placed under an atmosphere of hydrogen (3.8×10^5 Pa) and shaken at ambient temperature. After 17 h, the reaction mixture was diluted with 15 mL of 4:1 CHCl_3 :MeOH and then filtered through a pad of CELITE filter agent and rinsed through with portions of 4:1 CHCl_3 :MeOH (1% TFA). The clear colorless filtrate was concentrated under reduced pressure to yield a white solid. The material was suspended in 15 mL of H_2O and treated with 10% NaOH solution until the pH of the liquid was 13. A white solid was collected by vacuum filtration. The solid was purified by chromatography (SiO_2 , 25-30% (80:18:2 CHCl_3 :MeOH: NH_4OH) in CHCl_3) to yield a white foam. The foam was triturated with Et_2O and filtered to yield *N*-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)morpholine-4-carboxamide (0.050 g) as a white solid. mp 152–154 °C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 6.52 (t, $J = 5.4$ Hz, 1 H), 6.31 (t, $J = 5.9$ Hz, 1 H), 5.83 (s, 2 H), 4.61 (s, 2 H), 3.58-3.50 (m, 6 H), 3.24-3.21 (m, 4 H), 3.12 (q, $J = 6.7$ Hz, 2 H), 3.06-3.00 (m, 4 H), 2.67-2.64 (m, 2 H), 1.78-1.72 (m, 4 H), 1.66-1.61 (m, 2 H), 1.14 (t, $J = 7.0$ Hz, 3 H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 157.6, 149.4, 148.9, 146.5, 137.1, 105.0, 65.8, 65.2, 62.9, 51.3, 43.7, 37.8, 32.0, 28.1, 22.8, 22.5, 22.0, 14.9; MS (APCI) m/z 432 ($\text{M} + \text{H}$) $^+$; Anal. Calcd for $\text{C}_{21}\text{H}_{33}\text{N}_7\text{O}_3 \cdot 0.5\text{H}_2\text{O}$: C, 57.25; H, 7.78; N, 22.26; Found: C, 57.30; H, 7.82; N, 22.29.

25

Example 29

*N*¹-Isopropyl-7-[3-(morpholin-4-ylcarbonyl)phenyl]-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine




5

A suspension of 7-bromo-*N*¹-isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.00 g, 2.76 mmol) in 20 mL of 1-propanol was treated with the 3-(morpholine-4-carbonyl)phenylboronic acid (0.715 g, 3.04 mmol). The headspace of the flask was purged and back-filled with nitrogen (3X). The reaction mixture was then treated with triphenylphosphine (11 mg, 0.041 mmol), sodium carbonate (1.66 mL, 3.31 mmol, 2 M solution in H₂O), H₂O (2 mL), and palladium (II) acetate (3.1 mg, 0.014 mmol). Again the headspace of the flask was purged and back filled with nitrogen (3X). The reaction mixture was heated to 100 °C. After 16 h, the reaction mixture was concentrated under reduced pressure to yield a brown solid. The solid was partitioned between CHCl₃ (45 mL) and H₂O (15 mL) and separated. The organic portion was washed again with H₂O (15 mL) and brine (15 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan solid. The solid was purified by chromatography (SiO₂, 95:5 CHCl₃: MeOH) to yield a light yellow foam. The foam was crystallized from 1:1 acetonitrile (MeCN):MeOH to give *N*¹-isopropyl-7-[3-(morpholin-4-ylcarbonyl)phenyl]-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.683 g) as light yellow crystals. mp 274–276 °C; ¹H NMR (300 MHz, DMSO-*d*₆, 350 K) δ 8.56 (d, *J* = 8.5 Hz, 1 H), 7.86–7.81 (m, 2 H), 7.74 (s, 1 H), 7.57–7.52 (m, 2 H), 7.38 (d, *J* = 7.6 Hz, 1 H), 6.81 (s, 1 H), 6.21 (s, 2 H), 3.65–3.62 (m, 4 H), 3.58–3.54 (m, 5 H), 2.93 (t, *J* = 7.5 Hz, 2 H), 1.96–1.83 (m, 2 H), 1.07–1.02 (m, 9 H); ¹³C NMR (75 MHz, DMSO-*d*₆, 350 K) δ 169.5, 154.9, 152.3, 145.4, 141.1, 137.8, 136.9, 132.8, 129.4, 128.1, 126.1, 125.5, 124.6, 124.1, 122.0, 119.9, 114.7, 66.5, 51.4, 45.4, 28.3, 20.9, 20.6, 14.1; MS (APCI) *m/z* 473 (M + H)⁺; Anal. Calcd for C₂₇H₃₂N₆O₂: C, 68.62; H, 6.83; N, 17.78; Found: C, 68.73; H, 6.71; N, 17.85.

25

Example 30

N-(3-{[4-Amino-2-(ethoxymethyl)-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-7-yl]oxy}propyl)methanesulfonamide



5

Part A

A solution of *tert*-butyl [3-(4-amino-2-ethoxymethyl-1-isopropylamino-1*H*-imidazo[4,5-*c*]quinolin-7-yloxy)propyl]carbamate (1.24 g, 2.62 mmol) in 5 mL of ethanol (EtOH) was treated with 5 mL of 4.3 M HCl in EtOH. The reaction was heated to 100 °C. After 1 h, the reaction was cooled to ambient temperature and concentrated under reduced pressure to yield a brown solid. The solid was dissolved in a minimum amount of H₂O and treated with 10% NaOH solution until the pH of the liquid was 13. The basic aqueous mixture was extracted with CH₂Cl₂ (3 X 25 mL). The combined organic extracts were washed with brine (25 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield 7-(3-aminopropoxy)-2-ethoxymethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.84 g) as a tan solid.

15

Part B

A solution of 7-(3-aminopropoxy)-2-ethoxymethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.84 g, 2.26 mmol) in 25 mL of CH₂Cl₂ was placed under an atmosphere of nitrogen. The solution was treated with triethylamine (0.63 mL, 4.75 mmol) and cooled in an ice water bath. The solution was slowly treated with methanesulfonyl chloride (0.184 mL, 2.37). The reaction was stirred and allowed to slowly come to ambient temperature. After 16 h, the reaction mixture was diluted with CHCl₃ (20 mL) and transferred to a separatory funnel. The organic solution was washed with 10% Na₂CO₃ solution (15 mL), H₂O (15 mL) and brine (15 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan foam. The foam was purified by chromatography (SiO₂, 10-50% (80:18:2 CHCl₃:MeOH:NH₄OH) in CHCl₃) to give a light yellow solid. The solid was crystallized from MeCN to yield *N*-(3-{[4-amino-2-(ethoxymethyl)-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-7-

20


25

30

yl]oxy}propyl)methanesulfonamide (0.295 g) as white crystals. mp 181–183 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.46 (d, *J* = 9.0 Hz, 1 H), 7.09 (t, *J* = 6.8 Hz, 1 H), 7.02 (d, *J* = 2.6 Hz, 1 H), 6.90 (d, *J* = 2.1 Hz, 1 H), 6.87 (d, *J* = 2.5 Hz, 1 H), 6.54 (s, 2 H), 4.71 (s, 2 H), 4.11 (t, *J* = 6.1 Hz, 2 H), 3.64–3.58 (m, 3 H), 3.14 (q, *J* = 6.1 Hz, 2 H), 2.91 (s, 3 H), 1.96 (p, *J* = 6.6 Hz, 2 H), 1.15 (t, *J* = 7.0 Hz, 3 H), 1.01 (d, *J* = 6.1 Hz, 6 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 158.1, 152.6, 149.4, 147.2, 133.7, 122.6, 111.5, 109.1, 107.9, 65.8, 65.0, 63.0, 51.6, 29.7, 20.6, 15.3; MS (APCI) *m/z* 451 (M + H)⁺; Anal. Calcd for C₂₀H₃₀N₆O₄S: C, 53.32; H, 6.71; N, 18.65; S, 7.12; Found: C, 53.44; H, 6.68; N, 18.83; S, 7.09.

10

Example 31

*N*¹-Isopropyl-7-phenyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

15 A solution of 7-bromo-*N*¹-isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.00 g, 2.76 mmol) in 20 mL 1-propanol was treated with phenylboronic acid (0.371 g, 3.04 mmol) giving a suspension. The headspace of the flask was purged and back-filled with nitrogen (3X). The reaction mixture was then treated with triphenylphosphine (11 mg, 0.041 mmol), sodium carbonate (1.66 mL, 3.31 mmol, 2 M solution in H₂O), H₂O (2 mL), and palladium (II) acetate (3.1 mg, 0.014 mmol). Again the headspace of the flask was purged and back-filled with nitrogen (3X). The reaction mixture was heated to 100 °C. After 16 h, the reaction mixture was concentrated under reduced pressure to yield a brown solid. The solid was partitioned between CHCl₃ (45 mL) and H₂O (15 mL) and separated. The organic portion was washed again with H₂O (15 mL) and brine (15 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a tan solid. The solid was recrystallized from MeCN to yield *N*¹-isopropyl-7-phenyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.401 g) as light yellow crystals. mp 217–220 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.53 (d, *J* = 8.5 Hz, 1 H), 7.83 (d, *J* = 1.8 Hz, 1 H), 7.78–7.76 (m, 2 H), 7.55 (dd, *J* = 7.5, 1.9 Hz, 1 H), 7.52–7.47


20

25

(m, 2 H), 7.40-7.35 (m, 1 H), 7.01 (s, 1 H), 6.53 (s, 2 H), 3.57-3.49 (m, 1 H), 2.92 (bs, 2 H), 1.94-1.81 (m, 2 H), 1.06-1.01 (m, 9 H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 154.2, 151.9, 144.9, 140.3, 138.1, 132.1, 128.9, 127.2, 126.7, 123.9, 123.2, 121.4, 119.4, 113.8, 50.8, 27.8, 20.7, 20.2, 13.9; MS (ESI) m/z 360 ($\text{M} + \text{H}$) $^+$; Anal. Calcd for $\text{C}_{22}\text{H}_{25}\text{N}_5$: C, 73.51; H, 7.01; N, 19.48; Found: C, 73.39; H, 7.20; N, 19.36.

Example 32

N-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide




A solution of *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.00 g, 3.18 mmol) in 25 mL of CH_2Cl_2 was placed under an atmosphere of nitrogen and then chilled in an ice water bath. The solution was treated with triethylamine (0.89 mL, 6.7 mmol) and then slowly with isobutyryl chloride (0.35 mL, 3.3 mmol). The reaction was stirred and allowed to slowly come to ambient temperature. After 16 h, the reaction was diluted with 15 mL of CHCl_3 and 15 mL of 10% Na_2CO_3 solution and the layers were separated. The organic portion was washed with H_2O (15 mL). The combined aqueous washes were back-extracted with CHCl_3 (10 mL). The combined organic extracts were washed with brine (15 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield a light yellow foam. The foam was purified by chromatography (SiO_2 , 10-50% (80:18:2 CHCl_3 : MeOH : NH_4OH) in CHCl_3) to yield a white foam. The foam was triturated with hexanes and filtered to yield *N*-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide (0.821 g) as a white solid. mp 108–110 °C; ^1H NMR (300 MHz, DMSO- d_6) δ 8.43 (d, J = 7.5 Hz, 1 H), 7.72 (t, J = 5.4 Hz, 1 H), 7.58 (d, J = 7.9 Hz, 1 H), 7.46-7.41 (m, 1 H), 7.26-7.21 (m, 1 H), 6.97 (t, J = 5.5 Hz, 1 H), 6.62 (s, 2 H), 4.75 (s, 2 H), 3.61 (q, J = 7.0 Hz, 2 H), 3.17-3.13 (m, 4 H), 2.29 (h, J = 6.9 Hz, 1 H), 1.71 (p, J = 7.1 Hz, 2 H), 1.17 (t, J = 7.0 Hz, 3 H), 0.95 (d, J = 6.8 Hz, 3 H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 176.4, 152.3, 149.5, 145.3, 132.4, 127.4, 126.1, 124.2, 121.2, 114.7, 65.8, 63.1, 50.2, 36.6, 34.4, 28.0, 19.9,

15.3; MS (APCI) m/z 385 ($M + H$)⁺; Anal. Calcd for C₂₀H₂₈N₆O₂•0.25H₂O: C, 61.76; H, 7.38; N, 21.61; Found: C, 61.67; H, 7.55; N, 21.72.

Example 33


5 *N*-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide



A solution of *N*-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide (0.450 g, 1.17 mmol) in 25 mL of TFA was treated with platinum (IV) oxide (0.266 g, 1.17 mmol). The reaction mixture was shaken under an atmosphere of hydrogen (3.8 x 10⁵ Pa) at ambient temperature. After 17 h, the reaction mixture was filtered through a pad of CELITE filter agent and rinsed with portions of 4:1 CHCl₃:MeOH (1% TFA). The filtrate was concentrated under reduced pressure to yield a white oil. The oil was suspended in a minimum amount of H₂O and treated with 50% NaOH solution until the pH was 13. A white solid precipitated and was triturated in the basic H₂O for 1 h and then filtered to yield a sticky white solid. The solid was then triturated with Et₂O for 2 h, filtered and dried to yield *N*-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide (0.237 g) as a white solid. mp 163–165 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ 7.69 (t, *J* = 5.5 Hz, 1 H), 6.31 (t, *J* = 5.9 Hz, 1 H), 5.80 (s, 2 H), 4.61 (s, 2 H), 3.55 (q, *J* = 7.0 Hz, 2 H), 3.15-3.11 (m, 2 H), 3.04-3.00 (m, 4 H), 2.66 (t, *J* = 5.7 Hz, 2 H), 2.31 (h, *J* = 6.8 Hz, 1 H), 1.81-1.69 (m, 4 H), 1.62 (p, *J* = 7.1 Hz, 2 H), 1.14 (t, *J* = 7.0 Hz, 3 H), 0.98 (d, *J* = 6.9 Hz, 6 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 176.3, 149.9, 149.3, 147.0, 137.6, 122.7, 105.4, 65.7, 63.3, 51.7, 36.6, 34.4, 32.5, 28.0, 23.3, 23.0, 22.4, 19.9, 15.3; MS (ESI) m/z 389 ($M + H$)⁺; Anal. Calcd for C₂₀H₃₂N₆O₂: C, 61.83; H, 8.30; N, 21.63; Found: C, 61.63; H, 8.23; N, 21.55.

Example 34

N-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide




5

A solution of *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.00 g, 3.18 mmol) in 15 mL of CH₂Cl₂ was placed under an atmosphere of nitrogen. The solution was chilled in an ice water bath and treated first with triethylamine (0.89 mL, 6.7 mmol) and then slowly with acetyl chloride (0.89 mL, 3.3 mmol). After 2 h, the reaction was warmed to ambient temperature, quenched with 15 mL of saturated NaHCO₃ solution and diluted with 30 mL of CHCl₃. The phases were separated and the organic portion was washed with H₂O (15 mL) and brine (15 mL). The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield a light yellow foam. The foam was crystallized from MeCN to give light yellow crystals. The crystals were pulverized and then triturated with Et₂O for 18 h and filtered off to yield *N*-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide (0.728 g) as a white solid. mp 178–181 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.44 (d, *J* = 8.1 Hz, 1 H), 7.82 (t, *J* = 5.1 Hz, 1 H), 7.59 (d, *J* = 8.2 Hz, 1 H), 7.45–7.42 (m, 1 H), 7.26–7.22 (m, 1 H), 6.96 (t, *J* = 5.6 Hz, 1 H), 6.59 (s, 2 H), 4.75 (s, 2 H), 3.61 (q, *J* = 7.0 Hz, 2 H), 3.20–3.14 (m, 4 H), 1.77 (s, 3 H), 1.71 (p, *J* = 7.0 Hz, 2 H), 1.17 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 169.4, 152.3, 149.5, 145.3, 132.4, 127.4, 126.1, 124.2, 121.2, 114.7, 65.8, 36.1, 50.2, 36.8, 27.9, 22.9, 15.3; MS (ESI) *m/z* 357 (M + H)⁺; Anal. Calcd for C₁₈H₂₄N₆O₂: C, 60.66; H, 6.79; N, 23.58; Found: C, 60.55; H, 6.84; N, 23.44.

20

Example 35

*N*¹-Isopropyl-2-propyl-7-(2-pyridin-3-ylethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

5 Part A

A suspension of 7-bromo-*N*¹-isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.71 g, 4.72 mmol) in 34 mL of 1-propanol was placed under an atmosphere of nitrogen. The mixture was treated with triethylamine (1.25 mL, 9.44 mmol) and then the headspace of the flask was evacuated and back-filled with nitrogen (3X). The reaction mixture was treated with potassium vinylfluoroborate (0.695 g, 5.19 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (0.069 g, 0.094 mmol) and the head-space purged and backfilled as before. The reaction was heated to 100 °C. After 16 h, the reaction mixture was cooled to ambient temperature and concentrated under reduced pressure. The residue was dissolved in 50 mL of CHCl₃ and washed with 5% Na₂CO₃ solution (15 mL). There was a thick emulsion that did not fully separate. The organic portion was washed with H₂O (15 mL) resulting in another slowly separating emulsion. The combined aqueous washes were back extracted with CHCl₃ (15 mL). The combined organic extracts were washed with brine (15 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield an orange solid. The solid was purified by chromatography (SiO₂, 20% (80:18:2 CHCl₃:MeOH:NH₄OH) in CHCl₃) to yield *N*¹-isopropyl-2-propyl-7-vinyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.92 g) as a tan solid.

Part B


A suspension of palladium (II) acetate (0.067 g, 0.30 mmol) and tri-*o*-tolylphosphine (0.271 g, 0.891 mmol) in 10 mL of MeCN was placed in a 50 mL heavy wall glass pressure flask and placed under an atmosphere of nitrogen. The mixture was stirred until homogeneous. The mixture was treated with triethylamine (1.58 mL, 11.9 mmol), 3-bromopyridine (0.430 mL, 4.46 mmol) and *N*¹-isopropyl-2-propyl-7-vinyl-1*H*-

imidazo[4,5-*c*]quinoline-1,4-diamine (0.92 g, 3.0 mmol). Nitrogen was bubbled through the mixture for 1 min and then the flask was sealed. The reaction was heated to 120 °C. After 15 h, the homogenous orange solution was cooled to ambient temperature and concentrated under reduced pressure. The residue was partitioned between 75 mL of CHCl₃ and 25 mL of 10% Na₂CO₃ solution. The phases were separated and the organic portion was washed with H₂O (25 mL). The combined aqueous portions were back-extracted with CHCl₃ (25 mL). The combined organic extracts were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to yield an orange solid. The solid was purified by chromatography (SiO₂, 20-50% (80:18:2 CHCl₃:MeOH:NH₄OH) in CHCl₃) to yield *N*¹-isopropyl-2-propyl-7-(2-pyridin-3-yl-vinyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.97 g) as a light yellow solid.

Part C

A solution of *N*¹-isopropyl-2-propyl-7-(2-pyridin-3-yl-vinyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.97 g, 2.5 mmol) in 15 mL of EtOH and 15 mL of MeOH was treated with palladium on carbon (0.10 g, 0.050 mmol, 5% w/w). The mixture was placed under an atmosphere of hydrogen (3.8 x 10⁵ Pa) and shaken at ambient temperature. After 3 d, the reaction mixture was filtered through a pad of CELITE filter agent and rinsed with a 1:1 MeOH:EtOH until the filtrate ran clear. The filtrate was concentrated under reduced pressure to yield a light green solid. The solid was purified by chromatography (SiO₂, 10-50% (80:18:2 CHCl₃:MeOH:NH₄OH) in CHCl₃) to yield a clear oil. The oil was twice crystallized from MeCN to give *N*¹-isopropyl-2-propyl-7-(2-pyridin-3-ylethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.224 g) as white crystals. mp 100–103 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.43 (d, *J* = 1.6 Hz, 1 H), 8.37 (dd, *J* = 3.8, 1.6 Hz, 1 H), 8.32 (d, *J* = 8.4 Hz, 1 H), 7.68-7.64 (m, 1 H), 7.37 (d, *J* = 1.5 Hz, 1 H), 7.31-7.26 (m, 1 H), 7.12 (dd, *J* = 8.4, 1.7 Hz, 1 H), 6.93 (d, *J* = 1.4 Hz, 1 H), 6.39 (s, 2 H), 3.52-3.44 (m, 1 H), 3.00 (s, 4 H), 2.89 (bs, 2 H), 1.91-1.79 (m, 2 H), 1.04-1.00 (m, 9 H); ¹³C NMR (125 MHz, DMSO-*d*₆) δ 154.2, 152.0, 150.1, 147.5, 145.0, 139.6, 137.2, 136.3, 132.8, 125.4, 123.9, 123.7, 122.0, 121.1, 113.1, 51.2, 37.0, 34.1, 28.2, 21.1, 20.6, 14.3; MS (APCI) *m/z* 389 (M + H)⁺; Anal. Calcd for C₂₃H₂₈N₆: C, 71.10; H, 7.26; N, 21.63; Found: C, 70.87; H, 7.54; N, 21.58.


Example 36

2-Ethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine

5 2-Ethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine was prepared in 5 steps from *tert*-butyl *N*-(3-aminoquinolin-4-yl)hydrazinecarboxylate in analogous fashion to the preparation of *N*¹-Isopropyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (Example 16). Triethyl orthopropionate was used in lieu of trimethyl orthobutyrate. The material was purified by chromatography (SiO₂, 20% (80:18:2 CHCl₃:MeOH:NH₄OH) in
10 CHCl₃) to yield an off white solid. The solid was crystallized from MeCN to yield 2-ethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.358 g) as light pink crystals. mp 245–247 °C; ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.47 (d, *J* = 8.1 Hz, 1 H), 7.57 (d, *J* = 8.0 Hz, 1 H), 7.39 (t, *J* = 7.5 Hz, 1 H), 7.22 (t, *J* = 7.5 Hz, 1 H), 6.95 (s, 1 H), 6.44 (s, 2 H), 3.54-3.46 (m, 1 H), 3.04-2.88 (m, 2 H), 1.37 (t, *J* = 7.1 Hz, 3 H), 1.01 (d, *J* =
15 5.8 Hz, 6 H); ¹³C NMR (125 MHz, DMSO-*d*₆) δ 155.6, 152.0, 145.0, 132.8, 126.8, 126.1, 124.1, 121.3, 120.9, 115.0, 51.2, 20.6, 19.7, 12.4; MS (APCI) *m/z* 270 (M + H)⁺; Anal. Calcd for C₁₅H₁₉N₅: C, 66.89; H, 7.11; N, 26.00; Found: C, 66.81; H, 7.22; N, 25.95.

Example 37

20 *N*-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide




25 A solution of *N*-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide (0.220 g, 0.617 mmol) in 15 mL of TFA was treated with platinum (IV) oxide (0.140 g, 0.617 mmol). The mixture was placed under an atmosphere of hydrogen (3.8 x 10⁵ Pa) and shaken at ambient temperature. After 18 h, the reaction mixture was diluted with CHCl₃ (15 mL) and filtered through a pad of CELITE filter agent

and rinsed with several portions of CHCl_3 (1% TFA). The filtrate was concentrated under reduced pressure to yield an off-white oil. The oil was suspended in a minimum amount of H_2O and treated with 50% NaOH solution until the pH of the liquid was 12-13. The mixture was extracted with CH_2Cl_2 (3 X 15 mL). The combined organic extracts were washed with brine (15 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield an off-white solid. The solid was purified by chromatography (SiO_2 , 15-20% (80:18:2 CHCl_3 :MeOH: NH_4OH) in CHCl_3) to yield a white solid. The solid was recrystallized from MeCN to yield *N*-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide (0.0633 g) as white crystals. mp 198–200 °C; ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.80 (t, J = 5.0 Hz, 1 H), 6.30 (t, J = 6.0 Hz, 1 H), 5.78 (s, 2 H), 4.61 s, 2 H), 3.55 (q, J = 7.0 Hz, 2 H), 3.12 (q, J = 6.7 Hz, 2 H), 3.05-3.01 (m, 4 H), 2.66 (t, J = 5.9 Hz, 2 H), 1.78-1.73 (m, 7 H), 1.62 (p, J = 7.1 Hz, 2 H), 1.14 (t, J = 7.0 Hz, 3 H); ^{13}C NMR (125 MHz $\text{DMSO}-d_6$) δ 169.4, 149.9, 149.3, 147.1, 137.6, 122.7, 105.5, 65.7, 63.4, 51.6, 36.8, 32.5, 27.9, 23.9, 23.0, 23.0, 22.5, 15.3; MS (APCI) m/z 361 ($\text{M} + \text{H}$) $^+$; Anal. Calcd for $\text{C}_{18}\text{H}_{28}\text{N}_6\text{O}_2$: C, 59.98; H, 7.83; N, 23.31; Found: C, 59.96; H, 8.04; N, 23.37.

Example 38

1-(3-{[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea




A solution of *N*¹-(3-aminopropyl)-2-ethoxymethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (1.42 g, 4.52 mmol) in 25 mL of CH_2Cl_2 was placed under an atmosphere of nitrogen and chilled in an ice water bath. The solution was slowly treated with ethyl isocyanate (0.375 mL, 4.74 mmol). Upon addition the reaction mixture became thick and additional CH_2Cl_2 had to be added to keep the mixture stirring. The reaction was allowed to slowly warm to ambient temperature. After 17 h, the reaction was concentrated under reduced pressure to yield an off-white solid. The solid was purified by chromatography (SiO_2 , 15-50% (80:18:2 CHCl_3 :MeOH: NH_4OH) in CHCl_3) to yield an off-white solid.

The solid was recrystallized from MeCN to give 1-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea (1.16 g) as a white solid. mp 138–141 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.45 (dd, *J* = 8.1, 1.0 Hz, 1 H), 7.58 (dd, *J* = 8.3, 0.9 Hz, 1 H), 7.46–7.41 (m, 1 H), 7.27–7.21 (m, 1 H), 6.96 (t, *J* = 5.6 Hz, 1 H), 6.60 (s, 2 H), 5.83 (t, *J* = 5.7 Hz, 1 H), 5.73 (t, *J* = 5.5 Hz, 1 H), 4.75 (s, 2 H), 3.61 (q, *J* = 7.0 Hz, 2 H), 3.20–3.08 (m, 4 H), 3.03–2.94 (m, 2 H), 1.68 (p, *J* = 6.9 Hz, 2 H), 1.17 (t, *J* = 7.0 Hz, 3 H), 0.96 (t, *J* = 7.1 Hz, 3 H); ¹³C NMR (125 MHz, DMSO-*d*₆) δ 158.4, 152.3, 149.5, 145.3, 132.4, 127.4, 126.1, 124.2, 121.3, 121.2, 114.8, 65.9, 63.1, 50.2, 37.6, 34.4, 28.9, 16.0, 15.4; MS (ESI) *m/z* 386 (M + H)⁺; Anal. Calcd for C₁₉H₂₇N₇O₂•0.5H₂O: C, 57.85; H, 7.15; N, 24.85; Found: C, 57.84; H, 7.31; N, 25.04.

Example 39

1-(3-{[4-Amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea



15


A solution of 1-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea (0.700 g, 1.82 mmol) in 20 mL of TFA was treated with platinum (IV) oxide (0.413 g, 1.82 mmol). The mixture was placed under an atmosphere of nitrogen (3.8 x 10⁵ Pa) and shaken at ambient temperature. After 22 h, the reaction mixture was diluted with 50 mL of CHCl₃ and 10 mL of MeOH and filtered through a pad of CELITE filter agent. The pad was rinsed with 20% MeOH in CHCl₃ until the filtrate ran clear. The filtrate was concentrated under reduced pressure to yield a clear, colorless oil. The oil was suspended in 25 mL of H₂O and treated with 50% NaOH until the pH of the liquid reached 13. A white solid precipitated. The solid was triturated in the basic solution for 15 min and then filtered to give a white solid. The solid was recrystallized from MeCN to yield 1-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea (0.400 g) as white crystals. mp 189–191 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ 6.31 (t, *J* = 5.9 Hz, 1 H), 5.83–5.79 (m, 3

25

H), 5.74 (t, $J = 5.6$ Hz, 1 H), 4.61 (s, 2 H), 3.55 (q, $J = 7.0$ Hz, 2 H), 3.11-2.94 (m, 8 H), 2.66 (t, $J = 5.8$ Hz, 2 H), 1.83-1.67 (m, 4 H), 1.59 (p, $J = 6.9$ Hz, 2 H), 1.14 (t, $J = 7.0$ Hz, 3 H), 0.97 (t, $J = 7.1$ Hz, 3 H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 158.4, 149.9, 149.3, 147.0, 137.6, 122.7, 105.5, 65.7, 63.3, 51.7, 37.5, 34.4, 32.5, 28.9, 23.2, 23.0, 22.4, 16.0, 15.3; MS (APCI) m/z 390 ($\text{M} + \text{H}$) $^+$; Anal. Calcd for $\text{C}_{19}\text{H}_{31}\text{N}_7\text{O}_2$: C, 58.59; H, 8.02; N, 25.17; Found: C, 58.35; H, 8.30; N, 25.31.

Example 40

N-{3-[(4-Amino-2-ethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}acetamide



10

A solution of *N*¹-(3-Aminopropyl)-2-ethyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (0.89 g, 3.1 mmol) in 30 mL of CH_2Cl_2 was placed under an atmosphere of nitrogen. The solution was treated first with triethylamine (0.92 mL, 6.6 mmol) and then slowly with acetyl chloride (0.234 mL, 3.29 mmol). After 16 h, another equivalent of acetyl chloride was added. After 3 h, the reaction was concentrated under reduced pressure to yield an orange oil. The oil was then treated with 20 mL of 1 M HCl in MeOH and stirred at ambient temperature to hydrolyze any bis-amide. After 16 h, the reaction was quenched with 25 mL of 10% Na_2CO_3 solution. The mixture was concentrated under reduced pressure to remove MeOH. The aqueous oily suspension was treated with 75 mL of 95:5 CHCl_3 :MeOH, transferred to a separatory funnel and separated. The organic portion was washed with brine (25 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield a yellow foam. The foam was purified by chromatography (40 g SiO_2 , 25% (80:18:2 CHCl_3 :MeOH: NH_4OH) in CHCl_3) to yield a light yellow solid. The solid was recrystallized from MeCN to give *N*-{3-[(4-amino-2-ethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}acetamide (0.67 g) as yellow crystals. mp 167–169 °C; ^1H NMR (300 MHz, DMSO- d_6) δ 8.42 (d, $J = 8.1$ Hz, 1 H), 7.85 (t, $J = 5.2$ Hz, 1 H), 7.59 (d, $J = 8.3$ Hz, 1 H), 7.44-7.38 (m, 1 H), 7.25-7.20 (m, 1 H), 6.96 (t, $J = 5.3$ Hz, 1 H), 6.51 (s, 2 H), 3.19-3.08 (m, 4 H), 2.97 (q, $J = 7.4$ Hz, 2 H), 1.78 (s, 3 H), 1.69 (p, $J = 7.0$ Hz, 2 H),


25

1.38 (t, $J = 7.5$ Hz, 3 H); ^{13}C NMR (125 MHz, $\text{DMSO-}d_6$) δ 169.5, 154.9, 152.0, 144.9, 132.1, 126.9, 126.1, 124.2, 121.1, 121.0, 114.9, 49.7, 36.8, 28.0, 23.0, 19.5, 12.4; MS (ESI) m/z 327 ($\text{M} + \text{H}$) $^+$; Anal. Calcd for $\text{C}_{17}\text{H}_{22}\text{N}_6\text{O}$: C, 62.56; H, 6.79; N, 25.75; Found: C, 62.40; H, 7.06; N, 25.82.

5

Example 41

N-{3-[(4-Amino-2-ethyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}acetamide



10


A solution of *N*-{3-[(4-amino-2-ethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}acetamide (0.22 g, 0.67 mmol) in 10 mL of TFA was treated with platinum(IV) oxide (0.15 g, 0.67 mmol). The mixture was placed under an atmosphere of hydrogen (3.8×10^5 Pa) and shaken at ambient temperature. After 44 h, the reaction mixture was diluted with 25 mL of 4:1 CHCl_3 :MeOH and filtered through a pad of CELITE filter agent. The filtrate was concentrated under reduced pressure to yield a clear oil. The oil was suspended in 15 mL of H_2O and treated with 50% NaOH solution until the pH of the liquid reached 13. The mixture was extracted with CH_2Cl_2 (3 x 15 mL). The combined organic extracts were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to yield a white foam. The foam was purified by chromatography (SiO_2 , 30% (80:18:2 CHCl_3 :MeOH: NH_4OH) in CHCl_3) to yield a clear colorless oil. The oil was crystallized from MeCN to give *N*-{3-[(4-amino-2-ethyl-6,7,8,9-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}acetamide (0.0601 g) as off-white crystals. mp 175–176 $^\circ\text{C}$; ^1H NMR (500 MHz, $\text{DMSO-}d_6$) δ 7.81 (t, $J = 5.2$ Hz, 1 H), 6.28 (t, $J = 5.7$ Hz, 1 H), 5.60 (s, 2 H), 3.12 (q, $J = 6.8$ Hz, 2 H), 3.01 (s, 2 H), 2.94 (q, $J = 6.1$ Hz, 2 H), 2.81 (q, $J = 7.5$ Hz, 2 H), 2.65 (t, $J = 5.8$ Hz, 2 H), 1.78 (s, 3 H), 1.77–1.72 (m, 4 H), 1.60 (p, $J = 7.1$ Hz, 2 H), 1.29 (t, $J = 7.5$ Hz, 3 H); ^{13}C NMR (125 MHz, $\text{DMSO-}d_6$) δ 168.9, 154.1, 148.8, 145.6, 137.0, 122.2, 105.0, 50.7, 36.3, 32.0, 27.5, 22.9, 22.6, 22.5, 22.1, 19.1, 11.8; MS

25

(APCI) m/z 331 ($M + H$)⁺; Anal. Calcd for C₁₇H₂₆N₆O: C, 61.79; H, 7.93; N, 25.43; Found: C, 61.80; H, 7.80; N, 25.67.

Example 42

5 [4-Amino-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol




A solution of 2-(ethoxymethyl)-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine (prepared as described in Example 5, 0.502 g, 1.68 mmol) in dichloromethane (15 mL) under a nitrogen atmosphere was cooled in an ice bath. A solution of boron tribromide in dichloromethane (1 M, 4.19 mL, 4.19 mmol) was added dropwise. The mixture was allowed to warm slowly to room temperature, then was stirred for 16 hours. The reaction was quenched with methanol (10 mL) and concentrated under reduced pressure to yield a white solid. The solid was treated with 6 M aqueous hydrochloric acid (20 mL) to form a white suspension that was stirred at 50 °C for 2 hours. The mixture was allowed to cool to room temperature and was adjusted to pH 7 with 50% aqueous sodium hydroxide solution. A white solid was isolated by filtration, washed with water, and dried under vacuum. The crude product was purified by chromatography (silica gel, eluted with 1:1 CHCl₃:(80:18:2 CHCl₃:MeOH:NH₄OH)) then was recrystallized from acetonitrile to afford 0.0577 g of [4-amino-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol as white crystals, mp 228–229 °C;

¹H NMR (300 MHz, DMSO-*d*₆) δ 8.61 (dd, *J* = 8.2, 1.2 Hz, 1 H), 7.57 (dd, *J* = 8.4, 0.9 Hz, 1 H), 7.44–7.38 (m, 1 H), 7.25–7.20 (m, 1 H), 6.87 (d, *J* = 2.2 Hz, 1 H), 6.56 (s, 2 H), 5.58 (t, *J* = 5.8 Hz, 1 H), 4.77 (d, *J* = 5.7 Hz, 2 H), 3.71–3.62 (m, 1 H), 1.03 (d, *J* = 6.1 Hz, 6 H);

MS (APCI) m/z 272 ($M + H$)⁺;
 Anal. Calcd for C₁₄H₁₇N₅O: C, 61.98; H, 6.32; N, 25.81; Found: C, 62.17; H, 6.25; N, 25.93.

Example 43


[4-Amino-1-(isopropylamino)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol



- 5 A mixture of [4-amino-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol (0.120 g, 0.442 mmol) and platinum (IV) oxide (0.100 g, 0.422 mmol) in trifluoroacetic acid (10 mL) was hydrogenated on a Parr apparatus at 50 psi (3.4×10^5 Pa) at room temperature for 18 hours. The mixture was diluted with chloroform (20 mL) and filtered through a pad of CELITE filter agent. The filter agent was rinsed with a 4:1
- 10 chloroform/methanol solution. The filtrate was concentrated under reduced pressure to yield an oil that was suspended in water (10 mL) and treated with 10% aqueous sodium hydroxide until pH 7 was reached. A white precipitate was isolated by filtration, rinsed with water, and dried overnight under vacuum to provide [4-amino-1-(isopropylamino)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol.
- 15 MS (APCI) m/z 276 ($M + H$)⁺.

Exemplary Compounds

- 20 Certain exemplary compounds, including some of those described above in the Examples, have the following Formula I-1d and the following R₁, R₂, and R₃ substituents, wherein each line of the table represents a specific compound.



I-1d

R ₁	R ₂	R ₃
isopropyl	hydrogen	2-(hydroxymethyl)phenyl

isopropyl	hydrogen	5-(hydroxymethyl)pyridin-3-yl
isopropyl	methyl	2-(hydroxymethyl)phenyl
isopropyl	methyl	5-(hydroxymethyl)pyridin-3-yl
isopropyl	ethyl	pyridin-3-yl
isopropyl	ethyl	benzyloxy
isopropyl	ethyl	2-methanesulfonylaminoethoxy
isopropyl	ethyl	3-methanesulfonylaminoethoxy
isopropyl	ethyl	2-(pyridin-3-yl)ethyl
isopropyl	ethyl	2-(hydroxymethyl)phenyl
isopropyl	ethyl	5-(hydroxymethyl)pyridin-3-yl
isopropyl	propyl	2-(hydroxymethyl)phenyl
isopropyl	propyl	5-(hydroxymethyl)pyridin-3-yl
isopropyl	butyl	2-(hydroxymethyl)phenyl
isopropyl	butyl	5-(hydroxymethyl)pyridin-3-yl
isopropyl	methoxymethyl	pyridin-3-yl
isopropyl	methoxymethyl	benzyloxy
isopropyl	methoxymethyl	2-methanesulfonylaminoethoxy
isopropyl	methoxymethyl	3-methanesulfonylaminoethoxy
isopropyl	methoxymethyl	2-(pyridin-3-yl)ethyl
isopropyl	methoxymethyl	2-(hydroxymethyl)phenyl
isopropyl	methoxymethyl	5-(hydroxymethyl)pyridin-3-yl
isopropyl	hydroxymethyl	pyridin-3-yl
isopropyl	hydroxymethyl	benzyloxy
isopropyl	hydroxymethyl	2-methanesulfonylaminoethoxy
isopropyl	hydroxymethyl	3-methanesulfonylaminoethoxy
isopropyl	hydroxymethyl	2-(pyridin-3-yl)ethyl
isopropyl	hydroxymethyl	2-(hydroxymethyl)phenyl
isopropyl	hydroxymethyl	5-(hydroxymethyl)pyridin-3-yl
isopropyl	2-methoxyethyl	2-(hydroxymethyl)phenyl
isopropyl	2-methoxyethyl	5-(hydroxymethyl)pyridin-3-yl
isopropyl	2-hydroxyethyl	pyridin-3-yl
isopropyl	2-hydroxyethyl	benzyloxy
isopropyl	2-hydroxyethyl	2-methanesulfonylaminoethoxy
isopropyl	2-hydroxyethyl	3-methanesulfonylaminoethoxy
isopropyl	2-hydroxyethyl	2-(pyridin-3-yl)ethyl
isopropyl	2-hydroxyethyl	2-(hydroxymethyl)phenyl
isopropyl	2-hydroxyethyl	5-(hydroxymethyl)pyridin-3-yl
isopropyl	ethoxymethyl	2-(hydroxymethyl)phenyl
isopropyl	ethoxymethyl	5-(hydroxymethyl)pyridin-3-yl
benzyl	hydrogen	2-(hydroxymethyl)phenyl
benzyl	hydrogen	5-(hydroxymethyl)pyridin-3-yl
benzyl	methyl	2-(hydroxymethyl)phenyl
benzyl	methyl	5-(hydroxymethyl)pyridin-3-yl
benzyl	ethyl	pyridin-3-yl
benzyl	ethyl	benzyloxy
benzyl	ethyl	2-methanesulfonylaminoethoxy
benzyl	ethyl	3-methanesulfonylaminoethoxy
benzyl	ethyl	2-(pyridin-3-yl)ethyl
benzyl	ethyl	2-(hydroxymethyl)phenyl
benzyl	ethyl	5-(hydroxymethyl)pyridin-3-yl

benzyl	propyl	2-(hydroxymethyl)phenyl
benzyl	propyl	5-(hydroxymethyl)pyridin-3-yl
benzyl	butyl	2-(hydroxymethyl)phenyl
benzyl	butyl	5-(hydroxymethyl)pyridin-3-yl
benzyl	methoxymethyl	pyridin-3-yl
benzyl	methoxymethyl	benzyloxy
benzyl	methoxymethyl	2-methanesulfonylaminoethoxy
benzyl	methoxymethyl	3-methanesulfonylaminoethoxy
benzyl	methoxymethyl	2-(pyridin-3-yl)ethyl
benzyl	methoxymethyl	2-(hydroxymethyl)phenyl
benzyl	methoxymethyl	5-(hydroxymethyl)pyridin-3-yl
benzyl	hydroxymethyl	pyridin-3-yl
benzyl	hydroxymethyl	benzyloxy
benzyl	hydroxymethyl	2-methanesulfonylaminoethoxy
benzyl	hydroxymethyl	3-methanesulfonylaminoethoxy
benzyl	hydroxymethyl	2-(pyridin-3-yl)ethyl
benzyl	hydroxymethyl	2-(hydroxymethyl)phenyl
benzyl	hydroxymethyl	5-(hydroxymethyl)pyridin-3-yl
benzyl	2-methoxyethyl	2-(hydroxymethyl)phenyl
benzyl	2-methoxyethyl	5-(hydroxymethyl)pyridin-3-yl
benzyl	2-hydroxyethyl	pyridin-3-yl
benzyl	2-hydroxyethyl	benzyloxy
benzyl	2-hydroxyethyl	2-methanesulfonylaminoethoxy
benzyl	2-hydroxyethyl	3-methanesulfonylaminoethoxy
benzyl	2-hydroxyethyl	2-(pyridin-3-yl)ethyl
benzyl	2-hydroxyethyl	2-(hydroxymethyl)phenyl
benzyl	2-hydroxyethyl	5-(hydroxymethyl)pyridin-3-yl
benzyl	ethoxymethyl	2-(hydroxymethyl)phenyl
benzyl	ethoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	hydrogen	2-(hydroxymethyl)phenyl
3-phenylpropyl	hydrogen	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	methyl	2-(hydroxymethyl)phenyl
3-phenylpropyl	methyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	ethyl	pyridin-3-yl
3-phenylpropyl	ethyl	benzyloxy
3-phenylpropyl	ethyl	2-methanesulfonylaminoethoxy
3-phenylpropyl	ethyl	3-methanesulfonylaminoethoxy
3-phenylpropyl	ethyl	2-(pyridin-3-yl)ethyl
3-phenylpropyl	ethyl	2-(hydroxymethyl)phenyl
3-phenylpropyl	ethyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	propyl	2-(hydroxymethyl)phenyl
3-phenylpropyl	propyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	butyl	2-(hydroxymethyl)phenyl
3-phenylpropyl	butyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	methoxymethyl	pyridin-3-yl
3-phenylpropyl	methoxymethyl	benzyloxy
3-phenylpropyl	methoxymethyl	2-methanesulfonylaminoethoxy
3-phenylpropyl	methoxymethyl	3-methanesulfonylaminoethoxy
3-phenylpropyl	methoxymethyl	2-(pyridin-3-yl)ethyl
3-phenylpropyl	methoxymethyl	2-(hydroxymethyl)phenyl

3-phenylpropyl	methoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	hydroxymethyl	pyridin-3-yl
3-phenylpropyl	hydroxymethyl	benzyloxy
3-phenylpropyl	hydroxymethyl	2-methanesulfonylaminoethoxy
3-phenylpropyl	hydroxymethyl	3-methanesulfonylaminoethoxy
3-phenylpropyl	hydroxymethyl	2-(pyridin-3-yl)ethyl
3-phenylpropyl	hydroxymethyl	2-(hydroxymethyl)phenyl
3-phenylpropyl	hydroxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	2-methoxyethyl	2-(hydroxymethyl)phenyl
3-phenylpropyl	2-methoxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	2-hydroxyethyl	pyridin-3-yl
3-phenylpropyl	2-hydroxyethyl	benzyloxy
3-phenylpropyl	2-hydroxyethyl	2-methanesulfonylaminoethoxy
3-phenylpropyl	2-hydroxyethyl	3-methanesulfonylaminoethoxy
3-phenylpropyl	2-hydroxyethyl	2-(pyridin-3-yl)ethyl
3-phenylpropyl	2-hydroxyethyl	2-(hydroxymethyl)phenyl
3-phenylpropyl	2-hydroxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-phenylpropyl	ethoxymethyl	2-(hydroxymethyl)phenyl
3-phenylpropyl	ethoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	hydrogen	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	hydrogen	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	methyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	methyl	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	ethyl	pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	ethyl	benzyloxy
3-[3-(isopropyl)ureido]propyl	ethyl	2-methanesulfonylaminoethoxy
3-[3-(isopropyl)ureido]propyl	ethyl	3-methanesulfonylaminoethoxy
3-[3-(isopropyl)ureido]propyl	ethyl	2-(pyridin-3-yl)ethyl
3-[3-(isopropyl)ureido]propyl	ethyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	ethyl	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	propyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	propyl	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	butyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	butyl	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	methoxymethyl	pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	methoxymethyl	benzyloxy
3-[3-(isopropyl)ureido]propyl	methoxymethyl	2-methanesulfonylaminoethoxy
3-[3-(isopropyl)ureido]propyl	methoxymethyl	3-methanesulfonylaminoethoxy
3-[3-(isopropyl)ureido]propyl	methoxymethyl	2-(pyridin-3-yl)ethyl
3-[3-(isopropyl)ureido]propyl	methoxymethyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	methoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	hydroxymethyl	pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	hydroxymethyl	benzyloxy
3-[3-(isopropyl)ureido]propyl	hydroxymethyl	2-methanesulfonylaminoethoxy
3-[3-(isopropyl)ureido]propyl	hydroxymethyl	3-methanesulfonylaminoethoxy
3-[3-(isopropyl)ureido]propyl	hydroxymethyl	2-(pyridin-3-yl)ethyl
3-[3-(isopropyl)ureido]propyl	hydroxymethyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	hydroxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	2-methoxyethyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	2-methoxyethyl	5-(hydroxymethyl)pyridin-3-yl

3-[3-(isopropyl)ureido]propyl	2-hydroxyethyl	pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	2-hydroxyethyl	benzyloxy
3-[3-(isopropyl)ureido]propyl	2-hydroxyethyl	2-methanesulfonylaminoethoxy
3-[3-(isopropyl)ureido]propyl	2-hydroxyethyl	3-methanesulfonylaminoethoxy
3-[3-(isopropyl)ureido]propyl	2-hydroxyethyl	2-(pyridin-3-yl)ethyl
3-[3-(isopropyl)ureido]propyl	2-hydroxyethyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	2-hydroxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-[3-(isopropyl)ureido]propyl	ethoxymethyl	2-(hydroxymethyl)phenyl
3-[3-(isopropyl)ureido]propyl	ethoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	hydrogen	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	hydrogen	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	methyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	methyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	ethyl	pyridin-3-yl
3-methanesulfonylaminoethyl	ethyl	benzyloxy
3-methanesulfonylaminoethyl	ethyl	2-methanesulfonylaminoethoxy
3-methanesulfonylaminoethyl	ethyl	3-methanesulfonylaminoethoxy
3-methanesulfonylaminoethyl	ethyl	2-(pyridin-3-yl)ethyl
3-methanesulfonylaminoethyl	ethyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	ethyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	propyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	propyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	butyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	butyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	methoxymethyl	pyridin-3-yl
3-methanesulfonylaminoethyl	methoxymethyl	benzyloxy
3-methanesulfonylaminoethyl	methoxymethyl	2-methanesulfonylaminoethoxy
3-methanesulfonylaminoethyl	methoxymethyl	3-methanesulfonylaminoethoxy
3-methanesulfonylaminoethyl	methoxymethyl	2-(pyridin-3-yl)ethyl
3-methanesulfonylaminoethyl	methoxymethyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	methoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	hydroxymethyl	pyridin-3-yl
3-methanesulfonylaminoethyl	hydroxymethyl	benzyloxy
3-methanesulfonylaminoethyl	hydroxymethyl	2-methanesulfonylaminoethoxy
3-methanesulfonylaminoethyl	hydroxymethyl	3-methanesulfonylaminoethoxy
3-methanesulfonylaminoethyl	hydroxymethyl	2-(pyridin-3-yl)ethyl
3-methanesulfonylaminoethyl	hydroxymethyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	hydroxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	2-methoxyethyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	2-methoxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	2-hydroxyethyl	pyridin-3-yl
3-methanesulfonylaminoethyl	2-hydroxyethyl	benzyloxy
3-methanesulfonylaminoethyl	2-hydroxyethyl	2-methanesulfonylaminoethoxy
3-methanesulfonylaminoethyl	2-hydroxyethyl	3-methanesulfonylaminoethoxy
3-methanesulfonylaminoethyl	2-hydroxyethyl	2-(pyridin-3-yl)ethyl
3-methanesulfonylaminoethyl	2-hydroxyethyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	2-hydroxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-methanesulfonylaminoethyl	ethoxymethyl	2-(hydroxymethyl)phenyl
3-methanesulfonylaminoethyl	ethoxymethyl	5-(hydroxymethyl)pyridin-3-yl

3-(morpholine-4-carbonylamino)propyl	hydrogen	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	hydrogen	benzyloxy
3-(morpholine-4-carbonylamino)propyl	hydrogen	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	hydrogen	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	hydrogen	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	hydrogen	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	hydrogen	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	methyl	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	methyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	methyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	methyl	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	methyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	methyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	methyl	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	ethyl	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	ethyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	ethyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	ethyl	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	ethyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	ethyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	ethyl	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	propyl	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	propyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	propyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	propyl	3-methanesulfonylaminopropoxy


3-(morpholine-4-carbonylamino)propyl	propyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	propyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	propyl	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	butyl	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	butyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	butyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	butyl	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	butyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	butyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	butyl	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	methoxymethyl	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	methoxymethyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	methoxymethyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	methoxymethyl	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	methoxymethyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	methoxymethyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	methoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	hydroxymethyl	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	hydroxymethyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	hydroxymethyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	hydroxymethyl	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	hydroxymethyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	hydroxymethyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	hydroxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	2-methoxyethyl	pyridin-3-yl

3-(morpholine-4-carbonylamino)propyl	2-methoxyethyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	2-methoxyethyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	2-methoxyethyl	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	2-methoxyethyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	2-methoxyethyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	2-methoxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	2-hydroxyethyl	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	2-hydroxyethyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	2-hydroxyethyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	2-hydroxyethyl	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	2-hydroxyethyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	2-hydroxyethyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	2-hydroxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	ethoxymethyl	pyridin-3-yl
3-(morpholine-4-carbonylamino)propyl	ethoxymethyl	benzyloxy
3-(morpholine-4-carbonylamino)propyl	ethoxymethyl	2-methanesulfonylaminoethoxy
3-(morpholine-4-carbonylamino)propyl	ethoxymethyl	3-methanesulfonylaminopropoxy
3-(morpholine-4-carbonylamino)propyl	ethoxymethyl	2-(pyridin-3-yl)ethyl
3-(morpholine-4-carbonylamino)propyl	ethoxymethyl	2-(hydroxymethyl)phenyl
3-(morpholine-4-carbonylamino)propyl	ethoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	hydrogen	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	hydrogen	benzyloxy
3-(isopropylcarbonylamino)propyl	hydrogen	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	hydrogen	3-methanesulfonylaminopropoxy
3-(isopropylcarbonylamino)propyl	hydrogen	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	hydrogen	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	hydrogen	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	methyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	methyl	benzyloxy
3-(isopropylcarbonylamino)propyl	methyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	methyl	3-methanesulfonylaminopropoxy


3-(isopropylcarbonylamino)propyl	methyl	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	methyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	methyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	ethyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	ethyl	benzyloxy
3-(isopropylcarbonylamino)propyl	ethyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	ethyl	3-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	ethyl	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	ethyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	ethyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	propyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	propyl	benzyloxy
3-(isopropylcarbonylamino)propyl	propyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	propyl	3-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	propyl	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	propyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	propyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	butyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	butyl	benzyloxy
3-(isopropylcarbonylamino)propyl	butyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	butyl	3-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	butyl	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	butyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	butyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	methoxymethyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	methoxymethyl	benzyloxy
3-(isopropylcarbonylamino)propyl	methoxymethyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	methoxymethyl	3-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	methoxymethyl	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	methoxymethyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	methoxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	hydroxymethyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	hydroxymethyl	benzyloxy
3-(isopropylcarbonylamino)propyl	hydroxymethyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	hydroxymethyl	3-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	hydroxymethyl	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	hydroxymethyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	hydroxymethyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	2-methoxyethyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	2-methoxyethyl	benzyloxy
3-(isopropylcarbonylamino)propyl	2-methoxyethyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	2-methoxyethyl	3-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	2-methoxyethyl	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	2-methoxyethyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	2-methoxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	2-hydroxyethyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	2-hydroxyethyl	benzyloxy
3-(isopropylcarbonylamino)propyl	2-hydroxyethyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	2-hydroxyethyl	3-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	2-hydroxyethyl	2-(pyridin-3-yl)ethyl

3-(isopropylcarbonylamino)propyl	2-hydroxyethyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	2-hydroxyethyl	5-(hydroxymethyl)pyridin-3-yl
3-(isopropylcarbonylamino)propyl	ethoxymethyl	pyridin-3-yl
3-(isopropylcarbonylamino)propyl	ethoxymethyl	benzyloxy
3-(isopropylcarbonylamino)propyl	ethoxymethyl	2-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	ethoxymethyl	3-methanesulfonylaminoethoxy
3-(isopropylcarbonylamino)propyl	ethoxymethyl	2-(pyridin-3-yl)ethyl
3-(isopropylcarbonylamino)propyl	ethoxymethyl	2-(hydroxymethyl)phenyl
3-(isopropylcarbonylamino)propyl	ethoxymethyl	5-(hydroxymethyl)pyridin-3-yl


Certain exemplary compounds, including some of those described above in the Examples, have the following Formulas (Ii, IIb, Ij, or Ik) and the following R₁ and R₂ substituents, wherein each line of the table is matched with Formula Ii, IIb, Ij, or Ik to represent a specific compound.




Ii



IIb



Ij



Ik

R ₁	R ₂
isopropyl	ethyl
isopropyl	methoxymethyl
isopropyl	hydroxymethyl
isopropyl	2-hydroxyethyl
benzyl	ethyl
benzyl	methoxymethyl
benzyl	hydroxymethyl
benzyl	2-hydroxyethyl
3-phenylpropyl	ethyl
3-phenylpropyl	methoxymethyl
3-phenylpropyl	hydroxymethyl
3-phenylpropyl	2-hydroxyethyl

3-[3-(isopropyl)ureido]propyl	ethyl
3-[3-(isopropyl)ureido]propyl	methoxymethyl
3-[3-(isopropyl)ureido]propyl	hydroxymethyl
3-[3-(isopropyl)ureido]propyl	2-hydroxyethyl
3-methanesulfonylamino	ethyl
3-methanesulfonylamino	methoxymethyl
3-methanesulfonylamino	hydroxymethyl
3-methanesulfonylamino	2-hydroxyethyl
3-(methylcarbonylamino)propyl /	hydrogen
3-(methylcarbonylamino)propyl	methyl
3-(methylcarbonylamino)propyl	ethyl
3-(methylcarbonylamino)propyl	propyl
3-(methylcarbonylamino)propyl	butyl
3-(methylcarbonylamino)propyl	methoxymethyl
3-(methylcarbonylamino)propyl	hydroxymethyl
3-(methylcarbonylamino)propyl	2-methoxyethyl
3-(methylcarbonylamino)propyl	2-hydroxyethyl
3-(methylcarbonylamino)propyl	ethoxymethyl
3-(isopropylcarbonylamino)propyl	hydrogen
3-(isopropylcarbonylamino)propyl	methyl
3-(isopropylcarbonylamino)propyl	ethyl
3-(isopropylcarbonylamino)propyl	propyl
3-(isopropylcarbonylamino)propyl	butyl
3-(isopropylcarbonylamino)propyl	methoxymethyl
3-(isopropylcarbonylamino)propyl	hydroxymethyl
3-(isopropylcarbonylamino)propyl	2-methoxyethyl
3-(isopropylcarbonylamino)propyl	2-hydroxyethyl
3-(isopropylcarbonylamino)propyl	ethoxymethyl
3-[(morpholin-4-carbonyl)amino]propyl	hydrogen
3-[(morpholin-4-carbonyl)amino]propyl	methyl
3-[(morpholin-4-carbonyl)amino]propyl	ethyl
3-[(morpholin-4-carbonyl)amino]propyl	propyl
3-[(morpholin-4-carbonyl)amino]propyl	butyl
3-[(morpholin-4-carbonyl)amino]propyl	methoxymethyl
3-[(morpholin-4-carbonyl)amino]propyl	hydroxymethyl
3-[(morpholin-4-carbonyl)amino]propyl	2-methoxyethyl
3-[(morpholin-4-carbonyl)amino]propyl	2-hydroxyethyl
3-[(morpholin-4-carbonyl)amino]propyl	ethoxymethyl

In the above tables, the term "ureido" has the same meaning as "aminocarbonylamino".

- 5 Compounds of the invention have been found to modulate cytokine biosynthesis by inducing the production of interferon α and/or tumor necrosis factor α in human cells when tested using the method described below.

CYTOKINE INDUCTION IN HUMAN CELLS

An in vitro human blood cell system is used to assess cytokine induction. Activity is based on the measurement of interferon and tumor necrosis factor (α) (IFN and TNF, respectively) secreted into culture media as described by Testerman et. al. in "Cytokine Induction by the Immunomodulators Imiquimod and S-27609", Journal of Leukocyte Biology, 58, 365-372 (September, 1995).

Blood Cell Preparation for Culture:

Whole blood from healthy human donors is collected by venipuncture into EDTA vacutainer tubes. Peripheral blood mononuclear cells (PBMC) are separated from whole blood by density gradient centrifugation using HISTOPAQUE-1077. Blood is diluted 1:1 with Dulbecco's Phosphate Buffered Saline (DPBS) or Hank's Balanced Salts Solution (HBSS). The PBMC layer is collected and washed twice with DPBS or HBSS and resuspended at 4×10^6 cells/mL in RPMI complete. The PBMC suspension is added to 48 well flat bottom sterile tissue culture plates (Costar, Cambridge, MA or Becton Dickinson Labware, Lincoln Park, NJ) containing an equal volume of RPMI complete media containing test compound.

Compound Preparation:

The compounds are solubilized in dimethyl sulfoxide (DMSO). The DMSO concentration should not exceed a final concentration of 1% for addition to the culture wells. The compounds are generally tested at concentrations ranging from 30-0.014 μ M.

Incubation:

The solution of test compound is added at 60 μ M to the first well containing RPMI complete and serial 3 fold dilutions are made in the wells. The PBMC suspension is then added to the wells in an equal volume, bringing the test compound concentrations to the desired range (30-0.014 μ M). The final concentration of PBMC suspension is 2×10^6 cells/mL. The plates are covered with sterile plastic lids, mixed gently and then incubated for 18 to 24 hours at 37°C in a 5% carbon dioxide atmosphere.

Separation:

Following incubation the plates are centrifuged for 10 minutes at 1000 rpm (~200 x g) at 4°C. The cell-free culture supernatant is removed with a sterile polypropylene pipet and transferred to sterile polypropylene tubes. Samples are maintained at -30 to -70°C until analysis. The samples are analyzed for interferon (α) by ELISA and for tumor necrosis factor (α) by ELISA or IGEN Assay.

Interferon (α) and Tumor Necrosis Factor (α) Analysis by ELISA:

Interferon (α) concentration is determined by ELISA using a Human Multi-Species kit from PBL Biomedical Laboratories, New Brunswick, NJ. Results are expressed in pg/mL.

Tumor necrosis factor (α) (TNF) concentration is determined using ELISA kits available from Biosource International, Camarillo, CA. Alternately, the TNF concentration can be determined by ORIGEN M-Series Immunoassay and read on an IGEN M-8 analyzer from IGEN International, Gaithersburg, MD. The immunoassay uses a human TNF capture and detection antibody pair from Biosource International, Camarillo, CA. Results are expressed in pg/mL.

Certain compounds of the invention may modulate cytokine biosynthesis by inhibiting production of tumor necrosis factor α (TNF- α) when tested using the method described below.

TNF- α INHIBITION IN MOUSE CELLS

The mouse macrophage cell line Raw 264.7 is used to assess the ability of compounds to inhibit tumor necrosis factor- α (TNF- α) production upon stimulation by lipopolysaccharide (LPS).

Single Concentration Assay:

Blood Cell Preparation for Culture

Raw cells (ATCC) are harvested by gentle scraping and then counted. The cell suspension is brought to 3×10^5 cells/mL in RPMI with 10 % fetal bovine serum (FBS). Cell suspension (100 μ L) is added to 96-well flat bottom sterile tissues culture plates

(Becton Dickinson Labware, Lincoln Park, NJ). The final concentration of cells is 3×10^4 cells/well. The plates are incubated for 3 hours. Prior to the addition of test compound the medium is replaced with colorless RPMI medium with 3 % FBS.

5 Compound Preparation

The compounds are solubilized in dimethyl sulfoxide (DMSO). The DMSO concentration should not exceed a final concentration of 1% for addition to the culture wells. Compounds are tested at 5 μ M. LPS (Lipopolysaccharide from *Salmonella typhimurium*, Sigma-Aldrich) is diluted with colorless RPMI to the EC₇₀ concentration as measured by a dose response assay.

Incubation

A solution of test compound (1 μ L) is added to each well. The plates are mixed on a microtiter plate shaker for 1 minute and then placed in an incubator. Twenty minutes later the solution of LPS (1 μ L, EC₇₀ concentration \sim 10 ng/ml) is added and the plates are mixed for 1 minute on a shaker. The plates are incubated for 18 to 24 hours at 37 °C in a 5 % carbon dioxide atmosphere.

TNF- α Analysis

Following the incubation the supernatant is removed with a pipet. TNF- α concentration is determined by ELISA using a mouse TNF- α kit (from Biosource International, Camarillo, CA). Results are expressed in pg/mL. TNF- α expression upon LPS stimulation alone is considered a 100% response.

25 Dose Response Assay:

Blood Cell Preparation for Culture

Raw cells (ATCC) are harvested by gentle scraping and then counted. The cell suspension is brought to 4×10^5 cells/mL in RPMI with 10 % FBS. Cell suspension (250 μ L) is added to 48-well flat bottom sterile tissues culture plates (Costar, Cambridge, MA). The final concentration of cells is 1×10^5 cells/well. The plates are incubated for 3 hours. Prior to the addition of test compound the medium is replaced with colorless RPMI medium with 3 % FBS.

Compound Preparation

The compounds are solubilized in dimethyl sulfoxide (DMSO). The DMSO concentration should not exceed a final concentration of 1% for addition to the culture wells. Compounds are tested at 0.03, 0.1, 0.3, 1, 3, 5 and 10 μ M. LPS (Lipopolysaccharide from *Salmonella typhimurium*, Sigma-Aldrich) is diluted with colorless RPMI to the EC₇₀ concentration as measured by dose response assay.

Incubation

A solution of test compound (200 μ l) is added to each well. The plates are mixed on a microtiter plate shaker for 1 minute and then placed in an incubator. Twenty minutes later the solution of LPS (200 μ L, EC₇₀ concentration \sim 10 ng/ml) is added and the plates are mixed for 1 minute on a shaker. The plates are incubated for 18 to 24 hours at 37 °C in a 5 % carbon dioxide atmosphere.

TNF- α Analysis

Following the incubation the supernatant is removed with a pipet. TNF- α concentration is determined by ELISA using a mouse TNF- α kit (from Biosource International, Camarillo, CA). Results are expressed in pg/mL. TNF- α expression upon LPS stimulation alone is considered a 100% response.

The complete disclosures of the patents, patent documents, and publications cited herein are incorporated by reference in their entirety as if each were individually incorporated. Various modifications and alterations to this invention will become apparent to those skilled in the art without departing from the scope and spirit of this invention. It should be understood that this invention is not intended to be unduly limited by the illustrative embodiments and examples set forth herein and that such examples and embodiments are presented by way of example only with the scope of the invention intended to be limited only by the claims set forth herein as follows.

WHAT IS CLAIMED IS:

1. A compound selected from the group consisting of:
 N -(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)cyclopentanecarboxamide;
5 1-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea;
 N -{3-[4-amino-1-(isopropylamino)-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-7-yl]phenyl}methanesulfonamide;
10 N^1 -isopropyl-7-[3-(morpholin-4-ylcarbonyl)phenyl]-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
 N -(3-{[4-amino-2-(ethoxymethyl)-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-7-yl]oxy}propyl)methanesulfonamide;
 N^1 -isopropyl-7-phenyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
15 N -(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide;
 N -(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;
 N^1 -isopropyl-2-propyl-7-(2-pyridin-3-ylethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
20 2-Ethyl- N^1 -isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
1-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea;
 N -{3-[(4-amino-2-ethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}acetamide;
[4-amino-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol;
25 1-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea;
 N -(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)morpholine-4-carboxamide;
 N -(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide;
30 N -(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;

- 1-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea;
N-{3-[(4-amino-2-ethyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}acetamide; and
 5 [4-amino-1-(isopropylamino)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol;
 or a pharmaceutically acceptable salt thereof.
2. The compound or salt of claim 1 wherein the compound or salt is selected from the
 10 group consisting of:
N-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)cyclopentanecarboxamide;
 1-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea;
 15 *N*-{3-[4-amino-1-(isopropylamino)-2-propyl-1*H*-imidazo[4,5-*c*]quinolin-7-yl]phenyl}methanesulfonamide;
*N*¹-isopropyl-7-[3-(morpholin-4-ylcarbonyl)phenyl]-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
N-(3-{[4-amino-2-(ethoxymethyl)-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-7-yl]oxy}propyl)methanesulfonamide;
 20 *N*¹-isopropyl-7-phenyl-2-propyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
N-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide;
N-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;
 25 *N*¹-isopropyl-2-propyl-7-(2-pyridin-3-ylethyl)-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
 2-ethyl-*N*¹-isopropyl-1*H*-imidazo[4,5-*c*]quinoline-1,4-diamine;
 1-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea;
 30 *N*-{3-[(4-amino-2-ethyl-1*H*-imidazo[4,5-*c*]quinolin-1-yl)amino]propyl}acetamide;
 [4-amino-1-(isopropylamino)-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol; and
 pharmaceutically acceptable salts thereof.

3. The compound or salt of claim 2 wherein the compound or salt is *N*-(3-{[4-amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide or a pharmaceutically acceptable salt thereof.

5

4. The compound or salt of claim 1 wherein the compound or salt is selected from the group consisting of:

1-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-isopropylurea;

10 *N*-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)morpholine-4-carboxamide;

N-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-2-methylpropanamide;


15 *N*-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;

1-(3-{[4-amino-2-(ethoxymethyl)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)-3-ethylurea;

N-(3-{[4-amino-2-ethyl-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-1-yl]amino}propyl)acetamide;

20 [4-amino-1-(isopropylamino)-6,7,8,9-tetrahydro-1*H*-imidazo[4,5-*c*]quinolin-2-yl]methanol; and pharmaceutically acceptable salts thereof.

5. A compound of the Formula (I-1):



I-1

25


wherein:

R_1' is selected from the group consisting of alkoxyalkylenyl and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R_1' is bonded;

5 R_1 is selected from the group consisting of:

- R_4 ,
 -Y- R_4 ,
 -X- R_5 ,
 -X-N(R_6)-Y- R_4 ,
 10 -X-C(R_7)-N(R_6)- R_4 ,
 -X-O-C(R_7)-N(R_6)- R_4 ,
 -X-S(O)₂-N(R_6)- R_4 , and
 -X-O- R_4 ;

or R_1' and R_1 together with the nitrogen atom to which they are bonded can join to
 15 form a group selected from the group consisting of:



R_2 is selected from the group consisting of:

-hydrogen,
 -alkyl,
 20 -alkenyl,
 -aryl,
 -heteroaryl,
 -heterocyclyl,
 -alkylene-Z-alkyl,
 -alkylene-Z-aryl,
 25 -alkylene-Z-alkenyl, and
 -alkyl or alkenyl substituted by one or more substituents selected from the group consisting of:

-OH,
 30 -halogen,

5 -N(R₆)₂,
 -C(R₇)-N(R₆)₂,
 -S(O)₂-N(R₆)₂,
 -N(R₆)-C(R₇)-C₁₋₁₀ alkyl,
 -N(R₆)-C(R₇)-aryl,
 -N(R₆)-S(O)₂-C₁₋₁₀ alkyl,
 -N(R₆)-S(O)₂-aryl,
 -C(O)-C₁₋₁₀ alkyl,
 10 -O-C(R₇)-C₁₋₁₀ alkyl,
 -O-C(R₇)-aryl,
 -O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl,
 -O-C(R₇)-N(R₆)-aryl;
 -C(O)-O-C₁₋₁₀ alkyl,
 15 -N₃,
 -aryl,
 -heteroaryl,
 -heterocyclyl,
 -C(O)-aryl, and
 -C(O)-heteroaryl;

20 R₃ is selected from the group consisting of:

-Z'-R₄',
 -Z'-X'-R₄',
 -Z'-X'-Y'-R₄', and
 -Z'-X'-R₅';

25 each R is independently selected from the group consisting of alkyl, alkenyl, alkoxy, halogen, fluoroalkyl, hydroxy, amino, alkylamino, and dialkylamino;


n is an integer from 0 to 4;

m is 0 or 1; with the proviso that when m is 1, then n is 0, 1, or 2;

30 R₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy,


halogen, nitro, hydroxy, mercapto, cyano, carboxy, formyl, aryl, aryloxy, arylalkoxy, heteroaryl, heteroaryloxy, heteroarylalkoxy, heterocyclyl, heterocyclalkylenyl, amino, alkylamino, (arylalkylenyl)amino, dialkylamino, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo, with the proviso that when R₄ is a substituted alkyl group and the substituent contains a hetero atom which bonds directly to the alkyl group then the alkyl group contains at least two carbons between the substituent and the nitrogen atom to which R₁ is bonded;

R₅ is selected from the group consisting of:



X is C₂₋₂₀ alkylene;

Y is selected from the group consisting of -C(R₇)-, -C(R₇)-O-, -S(O)₂-, -S(O)₂-N(R₆)-, and -C(R₇)-N(R₉)-; wherein R₉ is selected from the group consisting of hydrogen, alkyl, and arylalkylenyl; or R₉ and R₄ together with the nitrogen atom to which R₉ is bonded can join to form the group



Z is selected from the group consisting of -O- and -S(O)₀₋₂;


A is selected from the group consisting of -CH(R₆)-, -O-, -N(R₆)-, -N(Y-R₄)-, and -N(X-N(R₆)-Y-R₄)-;

a and b are independently integers from 1 to 4 with the proviso that when A is -O-, -N(R₆)-, -N(Y-R₄)-, or -N(X-N(R₆)-Y-R₄)- then a and b are independently integers from 2 to 4;

R₄' is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, arylalkylenyl, aryloxyalkylenyl, alkylarylenyl, heteroaryl, heteroarylalkylenyl, heteroaryloxyalkylenyl, alkylheteroarylenyl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, hydroxyalkyl, haloalkyl, haloalkoxy, halogen,


nitro, hydroxy, mercapto, cyano, aryl, aryloxy, arylalkyleneoxy, heteroaryl, heteroaryloxy, heteroarylalkyleneoxy, heterocyclyl, amino, alkylamino, dialkylamino, (dialkylamino)alkyleneoxy, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo;


5 R_5' is selected from the group consisting of:



X' is selected from the group consisting of alkylene, alkenylene, alkynylene, arylene, heteroarylene, and heterocyclylene wherein the alkylene, alkenylene, and alkynylene groups can be optionally interrupted or terminated by arylene, heteroarylene, or heterocyclylene and optionally interrupted by one or more -O- groups;

Y' is selected from the group consisting of:





Z' is a bond or -O-;

A' is selected from the group consisting of -CH₂-, -O-, -C(O)-, -S(O)₀₋₂-, and
 5 -N(R₄')-;

Q is selected from the group consisting of a bond, -C(R₇)-, -C(R₇)-C(R₇)-,
 -S(O)₂-, -C(R₇)-N(R₁₁)-W-, -S(O)₂-N(R₁₁)-, -C(R₇)-O-, and -C(R₇)-N(OR₁₂)-;

V is selected from the group consisting of -C(R₇)-, -O-C(R₇)-, -N(R₁₁)-C(R₇)-, and
 -S(O)₂-;

10 W is selected from the group consisting of a bond, -C(O)-, and -S(O)₂-;

c and d are independently integers from 1 to 6 with the proviso that c + d is ≤ 7,
 and when A' is -O- or -N(R₄')- then c and d are independently integers from 2 to 4;

each R₆ is independently selected from the group consisting of hydrogen, alkyl,
 and arylalkylenyl;

15 each R₇ is independently selected from the group consisting of =O and =S;

each R₈ is independently C₂₋₇ alkylene;

R₁₀ is C₃₋₈ alkylene;


each R₁₁ is independently selected from the group consisting of hydrogen,
 C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₁₋₁₀ alkoxyC₂₋₁₀ alkylenyl, and arylC₁₋₁₀ alkylenyl; and

20 R₁₂ is selected from the group consisting of hydrogen and alkyl;
 or a pharmaceutically acceptable salt thereof.

6. The compound or salt of claim 5 wherein n is 0.

25 7. The compound or salt of claim 6 wherein R₃ is selected from the group consisting
 of -Z'-R₄', -Z'-X'-R₄', and -Z'-X'-Y'-R₄'.

8. The compound or salt of claim 7 wherein R_3 is selected from the group consisting of 2-(pyridin-3-yl)ethyl, pyridinyl, (hydroxymethyl)pyridinyl, phenyl, (hydroxymethyl)phenyl, ethoxyphenyl, (morpholine-4-carbonyl)phenyl, 3-(methanesulfonylamino)phenyl, 2-(methanesulfonylamino)ethoxy, 3-(methanesulfonylamino)propoxy, and benzyloxy.
9. The compound or salt of claim 5 wherein m is 0 and n is 1.
10. The compound or salt of claim 9 wherein R is halogen or hydroxy.
11. The compound or salt of claim 5 wherein m and n are each 0.
12. A compound of the Formula (II-1):



II-1

wherein:

each R_A is independently selected from the group consisting of:

- halogen,
- hydroxy,
- alkyl,
- alkenyl,
- haloalkyl,
- alkoxy,
- alkylthio,
- $-NH_2$,
- $-NH(alkyl)$, and
- $-N(alkyl)_2$;


n is an integer from 0 to 4;

R₁' is selected from the group consisting of alkoxyalkylenyl and hydroxyalkylenyl wherein the alkylenyl group contains at least two carbon atoms between the hydroxy or alkoxy substituent and the nitrogen atom to which R₁' is bonded;

5 R₁ is selected from the group consisting of:

-R₄,
 -Y-R₄,
 -X-R₅,
 -X-N(R₆)-Y-R₄,
 10 -X-C(R₇)-N(R₆)-R₄,
 -X-O-C(R₇)-N(R₆)-R₄,
 -X-S(O)₂-N(R₆)-R₄, and
 -X-O-R₄;

or R₁' and R₁ together with the nitrogen atom to which they are bonded can join to
 15 form a group selected from the group consisting of:




R₂ is selected from the group consisting of:

-hydrogen,
 -alkyl,
 20 -alkenyl,
 -aryl,
 -heteroaryl,
 -heterocyclyl,
 -alkylene-Z-alkyl,
 -alkylene-Z-aryl,
 25 -alkylene-Z-alkenyl, and
 -alkyl or alkenyl substituted by one or more substituents selected from the group consisting of:

-OH,
 30 -halogen,

- 5
- N(R₆)₂,
 - C(R₇)-N(R₆)₂,
 - S(O)₂-N(R₆)₂,
 - N(R₆)-C(R₇)-C₁₋₁₀ alkyl,
 - N(R₆)-C(R₇)-aryl,
 - N(R₆)-S(O)₂-C₁₋₁₀ alkyl,
 - N(R₆)-S(O)₂-aryl,
 - C(O)-C₁₋₁₀ alkyl,
 - O-C(R₇)-C₁₋₁₀ alkyl,
 - 10 -O-C(R₇)-aryl,
 - O-C(R₇)-N(R₆)-C₁₋₁₀ alkyl,
 - O-C(R₇)-N(R₆)-aryl;
 - C(O)-O-C₁₋₁₀ alkyl,
 - N₃,
 - 15 -aryl,
 - heteroaryl,
 - heterocyclyl,
 - C(O)-aryl, and
 - C(O)-heteroaryl;
- 20 R₄ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl groups can be unsubstituted or substituted by one or more substituents independently selected from the group consisting of alkyl, alkoxy, haloalkyl, haloalkoxy, halogen, nitro, hydroxy, mercapto, cyano, carboxy, formyl, aryl, aryloxy, arylalkoxy,
- 25 heteroaryl, heteroaryloxy, heteroarylalkoxy, heterocyclyl, heterocyclylalkylenyl, amino, alkylamino, (arylalkylenyl)amino, dialkylamino, and in the case of alkyl, alkenyl, alkynyl, and heterocyclyl, oxo, with the proviso that when R₄ is a substituted alkyl group and the substituent contains a hetero atom which bonds directly to the alkyl group then the alkyl group contains at least two carbons between the substituent and the nitrogen atom to
- 30 which R₁ is bonded;

R₅ is selected from the group consisting of:



each R₆ is independently selected from the group consisting of hydrogen, alkyl, and arylalkylenyl;


each R₇ is independently selected from the group consisting of =O and =S;

5 R₈ is C₂₋₇ alkylene;

A is selected from the group consisting of -CH(R₆)-, -O-, -N(R₆)-, -N(Y-R₄)-, and -N(X-N(R₆)-Y-R₄)-;

X is C₂₋₂₀ alkylene;

Y is selected from the group consisting of -C(R₇)-, -C(R₇)-O-, -S(O)₂-,
 10 -S(O)₂-N(R₆)-, and -C(R₇)-N(R₉)-; wherein R₉ is selected from the group consisting of
 hydrogen, alkyl, and arylalkylenyl; or R₉ and R₄ together with the nitrogen atom to which
 R₉ is bonded can join to form the group



Z is selected from the group consisting of -O- and -S(O)₀₋₂-; and

15 a and b are independently integers from 1 to 4 with the proviso that when A is
-O-, -N(R₆)-, -N(Y-R₄)-, or -N(X-N(R₆)-Y-R₄)- then a and b are independently integers
from 2 to 4;

or a pharmaceutically acceptable salt thereof.

20 13. The compound or salt of claim 12 wherein n is 0.

14. The compound or salt of any one of claims 5 through 13 wherein R₁ is selected from the group consisting of -R₄, -Y-R₄, and -X-N(R₆)-Y-R₄ wherein Y is -C(R₇)-, -S(O)₂-, or -C(R₇)-N(R₉)-.

15. The compound or salt of any one of claims 5 through 14 wherein R₁ is selected from the group consisting of hydrogen, alkyl, alkenyl, arylalkylenyl, arylalkenylenyl, heteroarylalkylenyl, heteroarylalkenylenyl, aminoalkylenyl, alkoxyalkylenyl, acyl,

alkylsulfonylaminoalkylenyl, arylsulfonylaminoalkylenyl, alkylcarbonylaminoalkylenyl, alkylaminocarbonyl, alkylaminocarbonylaminoalkylenyl, arylaminocarbonyl, (arylalkylenyl)aminoalkylenyl, heterocyclylcarbonylaminoalkylenyl, arylaminocarbonylaminoalkylenyl, heteroarylcarbonylaminoalkylenyl, and
5 heteroarylaminocarbonylaminoalkylenyl.

16. The compound or salt of any one of claims 5 through 15 wherein R_1 is selected from the group consisting of hydrogen, methyl, isopropyl, butyl, 2-methylpropyl, 1-ethylpropyl, 3-methylbutyl, cyclohexyl, benzyl, 3-phenylpropyl, cinnamyl, furan-2-ylmethyl, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{-NHR}_{13}$, wherein R_{13} is selected from the group consisting of
10 methylcarbonyl, isopropylcarbonyl, cyclopentylcarbonyl, tetrahydropyran-4-ylcarbonyl, methanesulfonyl, phenylsulfonyl, benzyl, ethylaminocarbonyl, isopropylaminocarbonyl, morpholine-4-carbonyl, phenylaminocarbonyl, pyridin-3-ylcarbonyl, and pyridin-3-ylaminocarbonyl.

15

17. The compound or salt of any one of claims 5 through 16 wherein R_2 is selected from the group consisting of hydrogen, alkyl, alkoxyalkylenyl, and hydroxyalkylenyl.

18. The compound or salt of any one of claims 5 through 17 wherein R_2 is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl, n-butyl, methoxymethyl, 2-methoxyethyl, ethoxymethyl, hydroxymethyl, 2-hydroxyethyl, and 3-hydroxypropyl.
20

19. A pharmaceutical composition comprising a therapeutically effective amount of a compound or salt of any one of claims 1 through 18 and a pharmaceutically acceptable
25 carrier.

20. Use of a compound or salt of any one of claims 1 through 18 for the preparation of a medicament for inducing cytokine biosynthesis.

30 21. Use of a compound or salt of any one of claims 1 through 18 for the preparation of a medicament for treating a viral disease.

22. Use of a compound or salt of any one of claims 1 through 18 for the preparation of a medicament for treating a neoplastic disease.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/28754

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 314745, 31/5377; C07D 471/04; A61P 31/12, 35/00, 37/02

US CL : 514/293, 232.8; 546/82; 544/126

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/293, 232.8; 546/82; 544/126

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	US 2004/0176367 A1(GRIESGRABER et al) 09 September 2004 (09.09.2004), see entire document, especially the example compounds.	1-14
A	US 2004/0162309 A1 (GORDEN et al) 19 August 2004 (19.08.2004), pages 10-12, Table 1, especially compounds IRM 24 and IRM 60.	1-14

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

07 February 2005 (07.02.2005)

Date of mailing of the international search report

07 MAR 2005

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703) 305-3230

Authorized officer

Evelyn Huang

Telephone No. 703-308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/28754

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claims 15-22 are improper multiple dependent claims.
3. ☒ Claims Nos.: 15-22
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.